



9-21-00

A/BOX/seg

Express Mail No.: EK916750871US
Docket No.: 789CIP2C



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
PATENT APPLICATION TRANSMITTAL UNDER 37 CFR 1.53

BOX PATENT APPLICATION
Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Transmitted herewith for filing is the patent application of

Inventor(s): Y. Tom Tang, Chenghua Liu, Ping Zhou, Vinod Asundi, Feiyan Ren, Qing A. Zhao, Jie Zhang, Jian-Rui Wang, Tom Wehrman, Radoje T. Drmanac

Title: NOVEL NUCLEIC ACIDS AND POLYPEPTIDES

1. Type of application

- ☒ This is a new application for a
- ☒ Utility patent.
- ☐ Design patent.
- ☒ This is a continuation-in-part application of prior application no. 09/574,454 filed May 19, 2000, Attorney Docket No. 789CIP, which is a continuation-in-part application of prior application no. 09/519,705 filed March 07, 2000, Attorney Docket No. 789.

2. Application Papers Enclosed

- 1 Title Page
- 113 Pages of Specification (excluding Claims, Abstract, Drawings & Sequence Listing)
- 4 Page(s) of Claims
- 1 Page(s) of Abstract
- 0 Sheet(s) of Drawings (Figs. X-X) ☐ Formal ☐ Informal
- 102 Page(s) of Sequence Listing

CERTIFICATION UNDER 37 CFR 1.10

I hereby certify that this Patent Application Transmittal and the documents referred to as enclosed therewith are being deposited with the United States Postal Service on September 19, 2000, in an envelope addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231 utilizing the "Express Mail Post Office to Addressee" service of the United States Postal Service under Mailing Label No. EK916750871US


Leslie A. Moji

3. Oath or Declaration

- ☐ Enclosed
 - ☐ Executed by (check all applicable boxes)
 - ☐ Inventor(s)
 - ☐ Legal representative of inventors(s) (37 CFR 1.42 or 1.43)
 - ☐ Joint inventor or person showing a proprietary interest on behalf of inventor who refused to sign or cannot be reached
 - ☐ The petition required by 37 CFR 1.47 and the statement required by 37 CFR 1.47 are enclosed. See Item 5D below for fee.
- ☒ Unexecuted – the undersigned attorney or agent is authorized to file this application on behalf of the applicant(s). An executed declaration will follow.

4. Additional Papers Enclosed

- ☐ Preliminary Amendment
- ☐ Information Disclosure Statement
- ☐ Declaration of Biological Deposit
- ☒ Computer readable copy of sequence listing containing nucleotide and/or amino acid sequence
- ☒ Statement Under 37 CFR § 1.821
- ☒ Paper copy of sequence listing identical to computer copy (102 pages)
- ☐ Microfiche computer program
- ☒ Verified statement claiming small entity status under 37 CFR 1.9 and 1.27
- ☐ Associate Power of Attorney
- ☐ Verified translation of a non-English patent application
- ☒ Return receipt postcard
- ☐ Other _____

5. Priority Applications Under 35 USC 119

Certified copies of applications from which priority under 35 USC 119 is claimed are listed below and

- ☐ are attached.
- ☐ will follow.

6. **Filing Fee Calculation (37 CFR 1.16)**

A. ☒ **Utility Application**

CLAIMS AS FILED – INCLUDING PRELIMINARY AMENDMENT (IF ANY)						
			SMALL ENTITY		OTHER THAN A SMALL ENTITY	
	NO. FILED	NO. EXTRA	RATE	FEE	RATE	FEE
BASIC FEE				\$345.00		\$690.00
TOTAL	30-20	= 10	X 9 =	\$90.00	X 18 =	\$0.00
INDEP.	3-3	= 0	X 39 =	\$0.00	X 78 =	\$0.00
<input checked="" type="checkbox"/> First Presentation of Multiple Dependent Claim			+ 130 =	\$130.00	+ 260 =	\$0.00
FILING FEE:				\$565.00	OR	\$0.00

B. ☐ **Design Application (\$155.00/\$310.00) Filing Fee: \$**_____

C. ☐ **Plant Application (\$240.00/\$480.00) Filing Fee: \$**_____

D. **Other fees**

☐ **Recording Assignment [Fee -- \$40.00 per assignment] \$**_____

☐ **Other \$**_____

TOTAL FEES \$ **565.00**

7. Method of Payments of Fees

- ☐ Enclosed check
- ☒ Charge Deposit Account No. 501169. A duplicate copy of this transmittal is enclosed
- ☐ Not enclosed

8. Deposit Account and Refund Authorization

The Commissioner is hereby authorized to charge payment of any additional fees due or credit any overpayment to Deposit Account No. 501169. A duplicate copy of this transmittal is enclosed.


Please refund any overpayment to Hyseq, Inc. at the address below.

Please direct all future correspondence to Leslie A. Mooi at the address below.

Respectfully submitted,

Date: September 19, 2000

By:



Leslie A. Mooi
Attorney for Applicants
Registration No.: 37,047
HYSEQ, INC.
670 Almanor Avenue
Sunnyvale, CA 94085
(408) 524-8100
(408) 524-8145 (Telefacsimile)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) or Patentee(s): Y. Tom Tang, Chenghua Liu, Ping Zhou, Vinod Asundi, Feiyan Ren, Qing A. Zhao, Jie Zhang, Jian-Rui Wang, Tom Wehrman, Radoje T. Drmanac

Application No. or Patent No.: Not Yet Assigned

Filed or Issued: Herewith

For: NOVEL NUCLEIC ACIDS AND POLYPEPTIDES

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR § 1.9(f) AND 1.27(c)) - SMALL BUSINESS CONCERN**

I hereby declare that I am

- ☐ The owner of the small business concern identified below:
- ☒ An official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF CONCERN: HYSEQ, INC.
ADDRESS: 670 Almanor Avenue
Sunnyvale, CA 94085

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR § 121.12, and reproduced in 37 CFR § 1.9(d), for purposes of paying reduced fees under § 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to, and remain with, the small business concern identified above with regard to the invention, entitled NOVEL NUCLEIC ACIDS AND POLYPEPTIDES by inventors, Y. Tom Tang, Chenghua Liu, Ping Zhou, Vinod Asundi, Feiyan Ren, Qing A. Zhao, Jie Zhang, Jian-Rui Wang, Tom Wehrman, Radoje T. Drmanac, described in

- ☒ The specification filed herewith.
- ☐ Application Serial No. [], filed [Date].
- ☐ Patent No. [], issued [Date].

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR § 1.9(c), or by any concern which would not qualify as a small business concern under 37 CFR § 1.9(d) or a nonprofit organization under 37 CFR § 1.9(e).

Full Name: _____

Address: _____

() Individual () Small Business Concern () Nonprofit Organization

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate (37 CFR § 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under § 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Name of person signing: James N. Fletcher

Title of person
other than owner: Secretary

Address of person signing: HYSEQ, INC.
670 Almanor Avenue
Sunnyvale, CA 94085

Signature: _____

Date: _____

¹NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR § 1.27)

Our Ref. No.: 789CIP2C

NOVEL NUCLEIC ACIDS AND POLYPEPTIDES

Express Mail Label No.: EK916750871US

NOVEL NUCLEIC ACIDS AND POLYPEPTIDES

1. CROSS REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation-in-part application of U.S. Application Serial No. 09/574,454, filed May 19, 2000, Attorney Docket No. 789CIP, which in turn is a continuation-in-part application of U.S. Application Serial No. 09/519,705, filed March 07, 2000, Attorney Docket No. 789, both of which are incorporated herein by reference in their entirety.

2. BACKGROUND OF THE INVENTION

2.1 TECHNICAL FIELD

15 The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with uses for these polynucleotides and proteins, for example in therapeutic, diagnostic and research methods.

2.2 BACKGROUND

20 Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs, chemokines, and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of hybridization cloning; activity of the protein in the
25 case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization-based cloning techniques, have advanced the state of the art by making available large numbers of DNA/amino acid sequences for proteins that are
30 known to have biological activity, for example, by virtue of their secreted nature in the case of leader sequence cloning, by virtue of their cell or tissue source in the case of

PCR-based techniques, or by virtue of structural similarity to other genes of known biological activity.

Identified polynucleotide and polypeptide sequences have numerous applications in, for example, diagnostics, forensics, gene mapping; identification of mutations responsible for genetic disorders or other traits, to assess biodiversity, and to produce many other types of data and products dependent on DNA and amino acid sequences.

3. SUMMARY OF THE INVENTION

The compositions of the present invention include novel isolated polypeptides, novel isolated polynucleotides encoding such polypeptides, including recombinant DNA molecules, cloned genes or degenerate variants thereof, especially naturally occurring variants such as allelic variants, antisense polynucleotide molecules, and antibodies that specifically recognize one or more epitopes present on such polypeptides, as well as hybridomas producing such antibodies.

The compositions of the present invention additionally include vectors, including expression vectors, containing the polynucleotides of the invention, cells genetically engineered to contain such polynucleotides and cells genetically engineered to express such polynucleotides.

The present invention relates to a collection or library of at least one novel nucleic acid sequence assembled from expressed sequence tags (ESTs) isolated mainly by sequencing by hybridization (SBH), and in some cases, sequences obtained from one or more public databases. The invention relates also to the proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins. These nucleic acid sequences are designated as SEQ ID NO: 1 – 35 and are provided in the Sequence Listing. In the nucleic acids provided in the Sequence Listing, A is adenosine; C is cytosine; G is guanosine; T is thymine; and N is any of the four bases. In the amino acids provided in the Sequence Listing, * corresponds to the stop codon.

The nucleic acid sequences of the present invention also include, nucleic acid sequences that hybridize to the complement of SEQ ID NO: 1 – 35 under stringent hybridization conditions; nucleic acid sequences which are allelic variants or species

homologues of any of the nucleic acid sequences recited above, or nucleic acid sequences that encode a peptide comprising a specific domain or truncation of the peptides encoded by SEQ ID NO: 1 – 35. A polynucleotide comprising a nucleotide sequence having at least 90% identity to an identifying sequence of SEQ ID NO: 1 – 35 or a degenerate variant or fragment thereof. The identifying sequence can be 100 base pairs in length.

The nucleic acid sequences of the present invention also include the sequence information from the nucleic acid sequences of SEQ ID NO: 1 – 35. The sequence information can be a segment of any one of SEQ ID NO: 1 – 35 that uniquely identifies or represents the sequence information of SEQ ID NO: 1 – 35.

A collection as used in this application can be a collection of only one polynucleotide. The collection of sequence information or identifying information of each sequence can be provided on a nucleic acid array. In one embodiment, segments of sequence information is provided on a nucleic acid array to detect the polynucleotide that contains the segment. The array can be designed to detect full-match or mismatch to the polynucleotide that contains the segment. The collection can also be provided in a computer-readable format.

This invention also includes the reverse or direct complement of any of the nucleic acid sequences recited above; cloning or expression vectors containing the nucleic acid sequences; and host cells or organisms transformed with these expression vectors. Nucleic acid sequences (or their reverse or direct complements) according to the invention have numerous applications in a variety of techniques known to those skilled in the art of molecular biology, such as use as hybridization probes, use as primers for PCR, use in an array, use in computer-readable media, use in sequencing full-length genes, use for chromosome and gene mapping, use in the recombinant production of protein, and use in the generation of anti-sense DNA or RNA, their chemical analogs and the like.

In a preferred embodiment, the nucleic acid sequences of SEQ ID NO: 1-35 or novel segments or parts of the nucleic acids of the invention are used as primers in expression assays that are well known in the art. In a particularly preferred embodiment, the nucleic acid sequences of SEQ ID NO: 1-35 or novel segments or parts of the nucleic acids provided herein are used in diagnostics for identifying expressed genes or, as well known in

the art and exemplified by Vollrath et al., Science 258:52-59 (1992), as expressed sequence tags for physical mapping of the human genome.

The isolated polynucleotides of the invention include, but are not limited to, a polynucleotide comprising any one of the nucleotide sequences set forth in the SEQ ID NO: 1-35; a polynucleotide comprising any of the full length protein coding sequences of the SEQ ID NO: 1-35; and a polynucleotide comprising any of the nucleotide sequences of the mature protein coding sequences of the SEQ ID NO: 1-35. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent hybridization conditions to (a) the complement of any one of the nucleotide sequences set forth in the SEQ ID NO: 1-35; (b) a nucleotide sequence encoding any one of the amino acid sequences set forth in the Sequence Listing; (c) a polynucleotide which is an allelic variant of any polynucleotides recited above; (d) a polynucleotide which encodes a species homolog (e.g. orthologs) of any of the proteins recited above; or (e) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of any of the polypeptides comprising an amino acid sequence set forth in the Sequence Listing.

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising any of the amino acid sequences set forth in the Sequence Listing; or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides with biological activity that are encoded by (a) any of the polynucleotides having a nucleotide sequence set forth in the SEQ ID NO: 1-35; or (b) polynucleotides that hybridize to the complement of the polynucleotides of (a) under stringent hybridization conditions. Biologically or immunologically active variants of any of the polypeptide sequences in the Sequence Listing, and "substantial equivalents" thereof (e.g., with at least about 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% amino acid sequence identity) that preferably retain biological activity are also contemplated. The polypeptides of the invention may be wholly or partially chemically synthesized but are preferably produced by recombinant means using the genetically engineered cells (e.g. host cells) of the invention.

The invention also provides compositions comprising a polypeptide of the invention. Polypeptide compositions of the invention may further comprise an acceptable carrier, such as a hydrophilic, e.g., pharmaceutically acceptable, carrier.

The invention also provides host cells transformed or transfected with a polynucleotide of the invention.

The invention also relates to methods for producing a polypeptide of the invention comprising growing a culture of the host cells of the invention in a suitable culture medium under conditions permitting expression of the desired polypeptide, and purifying the polypeptide from the culture or from the host cells. Preferred embodiments include those in which the protein produced by such process is a mature form of the protein.

Polynucleotides according to the invention have numerous applications in a variety of techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use as oligomers, or primers, for PCR, use for chromosome and gene mapping, use in the recombinant production of protein, and use in generation of anti-sense DNA or RNA, their chemical analogs and the like. For example, when the expression of an mRNA is largely restricted to a particular cell or tissue type, polynucleotides of the invention can be used as hybridization probes to detect the presence of the particular cell or tissue mRNA in a sample using, *e.g.*, *in situ* hybridization.

In other exemplary embodiments, the polynucleotides are used in diagnostics as expressed sequence tags for identifying expressed genes or, as well known in the art and exemplified by Vollrath et al., Science 258:52-59 (1992), as expressed sequence tags for physical mapping of the human genome.

The polypeptides according to the invention can be used in a variety of conventional procedures and methods that are currently applied to other proteins. For example, a polypeptide of the invention can be used to generate an antibody that specifically binds the polypeptide. Such antibodies, particularly monoclonal antibodies, are useful for detecting or quantitating the polypeptide in tissue. The polypeptides of the invention can also be used as molecular weight markers, and as a food supplement.

Methods are also provided for preventing, treating, or ameliorating a medical condition which comprises the step of administering to a mammalian subject a therapeutically effective amount of a composition comprising a polypeptide of the present invention and a pharmaceutically acceptable carrier.

In particular, the polypeptides and polynucleotides of the invention can be utilized, for example, in methods for the prevention and/or treatment of disorders involving aberrant protein expression or biological activity.

The present invention further relates to methods for detecting the presence of the polynucleotides or polypeptides of the invention in a sample. Such methods can, for example, be utilized as part of prognostic and diagnostic evaluation of disorders as recited herein and for the identification of subjects exhibiting a predisposition to such conditions. The invention provides a method for detecting the polynucleotides of the invention in a sample, comprising contacting the sample with a compound that binds to and forms a complex with the polynucleotide of interest for a period sufficient to form the complex and under conditions sufficient to form a complex and detecting the complex such that if a complex is detected, the polynucleotide of interest is detected. The invention also provides a method for detecting the polypeptides of the invention in a sample comprising contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex and detecting the formation of the complex such that if a complex is formed, the polypeptide is detected.

The invention also provides kits comprising polynucleotide probes and/or monoclonal antibodies, and optionally quantitative standards, for carrying out methods of the invention. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and monitoring the progress of patients, involved in clinical trials for the treatment of disorders as recited above.

The invention also provides methods for the identification of compounds that modulate (i.e., increase or decrease) the expression or activity of the polynucleotides and/or polypeptides of the invention. Such methods can be utilized, for example, for the identification of compounds that can ameliorate symptoms of disorders as recited herein. Such methods can include, but are not limited to, assays for identifying compounds and other substances that interact with (e.g., bind to) the polypeptides of the invention. The invention provides a method for identifying a compound that binds to the polypeptides of the invention comprising contacting the compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound complex, wherein the

complex drives expression of a reporter gene sequence in the cell; and detecting the complex by detecting the reporter gene sequence expression such that if expression of the reporter gene is detected the compound the binds to a polypeptide of the invention is identified.

5 The methods of the invention also provides methods for treatment which involve the administration of the polynucleotides or polypeptides of the invention to individuals exhibiting symptoms or tendencies. In addition, the invention encompasses methods for treating diseases or disorders as recited herein comprising administering compounds and other substances that modulate the overall activity of the target gene products.

10 Compounds and other substances can effect such modulation either on the level of target gene/protein expression or target protein activity.

The polypeptides of the present invention and the polynucleotides encoding them are also useful for the same functions known to one of skill in the art as the polypeptides and polynucleotides to which they have homology (set forth in Table 1); for which they have a signature region (as set forth in Table 3); or for which they have homology to a gene family (as set forth in Table 4). If no homology is set forth for a sequence, then the polypeptides and polynucleotides of the present invention are useful for a variety of applications, as described herein, including use in arrays for detection.

20 4. DETAILED DESCRIPTION OF THE INVENTION

4.1 DEFINITIONS

It must be noted that as used herein and in the appended claims, the singular forms “a”, “an” and “the” include plural references unless the context clearly dictates otherwise.

25 The term "active" refers to those forms of the polypeptide which retain the biologic and/or immunologic activities of any naturally occurring polypeptide. According to the invention, the terms “biologically active” or “biological activity” refer to a protein or peptide having structural, regulatory or biochemical functions of a naturally occurring molecule. Likewise “immunologically active” or “immunological activity” refers to the

capability of the natural, recombinant or synthetic polypeptide to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "activated cells" as used in this application are those cells which are engaged in extracellular or intracellular membrane trafficking, including the export of secretory or enzymatic molecules as part of a normal or disease process.

The terms "complementary" or "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence 5'-AGT-3' binds to the complementary sequence 3'-TCA-5'. Complementarity between two single-stranded molecules may be "partial" such that only some of the nucleic acids bind or it may be "complete" such that total complementarity exists between the single stranded molecules. The degree of complementarity between the nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands.

The term "embryonic stem cells (ES)" refers to a cell that can give rise to many differentiated cell types in an embryo or an adult, including the germ cells. The term "germ line stem cells (GSCs)" refers to stem cells derived from primordial stem cells that provide a steady and continuous source of germ cells for the production of gametes. The term "primordial germ cells (PGCs)" refers to a small population of cells set aside from other cell lineages particularly from the yolk sac, mesenteries, or gonadal ridges during embryogenesis that have the potential to differentiate into germ cells and other cells. PGCs are the source from which GSCs and ES cells are derived. The PGCs, the GSCs and the ES cells are capable of self-renewal. Thus these cells not only populate the germ line and give rise to a plurality of terminally differentiated cells that comprise the adult specialized organs, but are able to regenerate themselves.

The term "expression modulating fragment," EMF, means a series of nucleotides which modulates the expression of an operably linked ORF or another EMF.

As used herein, a sequence is said to "modulate the expression of an operably linked sequence" when the expression of the sequence is altered by the presence of the EMF. EMFs include, but are not limited to, promoters, and promoter modulating sequences (inducible elements). One class of EMFs are nucleic acid fragments which induce the expression of an operably linked ORF in response to a specific regulatory factor or physiological event.

The terms "nucleotide sequence" or "nucleic acid" or "polynucleotide" or "oligonucleotide" are used interchangeably and refer to a heteropolymer of nucleotides or the sequence of these nucleotides. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA) or to any DNA-like or RNA-like material. It is contemplated that where the polynucleotide is RNA, the T (thymine) in the sequences provided herein is substituted with U (uracil). Generally, nucleic acid segments provided by this invention may be assembled from fragments of the genome and short oligonucleotide linkers, or from a series of oligonucleotides, or from individual nucleotides, to provide a synthetic nucleic acid which is capable of being expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial or viral operon, or a eukaryotic gene.

The terms "oligonucleotide fragment" or a "polynucleotide fragment", "portion," or "segment" or "probe" or "primer" are used interchangeable and refer to a sequence of nucleotide residues which are at least about 5 nucleotides, more preferably at least about 7 nucleotides, more preferably at least about 9 nucleotides, more preferably at least about 11 nucleotides and most preferably at least about 17 nucleotides. The fragment is preferably less than about 500 nucleotides, preferably less than about 200 nucleotides, more preferably less than about 100 nucleotides, more preferably less than about 50 nucleotides and most preferably less than 30 nucleotides. Preferably the probe is from about 6 nucleotides to about 200 nucleotides, preferably from about 15 to about 50 nucleotides, more preferably from about 17 to 30 nucleotides and most preferably from about 20 to 25 nucleotides. Preferably the fragments can be used in polymerase chain reaction (PCR), various hybridization procedures or microarray procedures to identify or amplify identical or related parts of mRNA or DNA molecules. A fragment or segment may uniquely identify each polynucleotide sequence of the present invention. Preferably the fragment comprises a sequence substantially similar to any one of SEQ ID NOs:1-35.

Probes may, for example, be used to determine whether specific mRNA molecules are present in a cell or tissue or to isolate similar nucleic acid sequences from chromosomal DNA as described by Walsh et al. (Walsh, P.S. et al., 1992, PCR Methods Appl 1:241-250). They may be labeled by nick translation, Klenow fill-in reaction, PCR,

or other methods well known in the art. Probes of the present invention, their preparation and/or labeling are elaborated in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, NY; or Ausubel, F.M. et al., 1989, Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, both of which are incorporated herein by reference in their entirety.

The nucleic acid sequences of the present invention also include the sequence information from the nucleic acid sequences of SEQ ID NOs: 1-35. The sequence information can be a segment of any one of SEQ ID NOs: 1-35 that uniquely identifies or represents the sequence information of that sequence of SEQ ID NO: 1-35. One such segment can be a twenty-mer nucleic acid sequence because the probability that a twenty-mer is fully matched in the human genome is 1 in 300. In the human genome, there are three billion base pairs in one set of chromosomes. Because 4^{20} possible twenty-mers exist, there are 300 times more twenty-mers than there are base pairs in a set of human chromosome. Using the same analysis, the probability for a seventeen-mer to be fully matched in the human genome is approximately 1 in 5. When these segments are used in arrays for expression studies, fifteen-mer segments can be used. The probability that the fifteen-mer is fully matched in the expressed sequences is also approximately one in five because expressed sequences comprise less than approximately 5% of the entire genome sequence.

Similarly, when using sequence information for detecting a single mismatch, a segment can be a twenty-five mer. The probability that the twenty-five mer would appear in a human genome with a single mismatch is calculated by multiplying the probability for a full match ($1/4^{25}$) times the increased probability for mismatch at each nucleotide position (3×25). The probability that an eighteen mer with a single mismatch can be detected in an array for expression studies is approximately one in five. The probability that a twenty-mer with a single mismatch can be detected in a human genome is approximately one in five.

The term "open reading frame," ORF, means a series of nucleotide triplets coding for amino acids without any termination codons and is a sequence translatable into protein.

The terms "operably linked" or "operably associated" refer to functionally related nucleic acid sequences. For example, a promoter is operably associated or operably

linked with a coding sequence if the promoter controls the transcription of the coding sequence. While operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements e.g. repressor genes are not contiguously linked to the coding sequence but still control transcription/translation of the coding sequence.

The term “pluripotent” refers to the capability of a cell to differentiate into a number of differentiated cell types that are present in an adult organism. A pluripotent cell is restricted in its differentiation capability in comparison to a totipotent cell.

The terms “polypeptide” or “peptide” or “amino acid sequence” refer to an oligopeptide, peptide, polypeptide or protein sequence or fragment thereof and to naturally occurring or synthetic molecules. A polypeptide “fragment,” “portion,” or “segment” is a stretch of amino acid residues of at least about 5 amino acids, preferably at least about 7 amino acids, more preferably at least about 9 amino acids and most preferably at least about 17 or more amino acids. The peptide preferably is not greater than about 200 amino acids, more preferably less than 150 amino acids and most preferably less than 100 amino acids. Preferably the peptide is from about 5 to about 200 amino acids. To be active, any polypeptide must have sufficient length to display biological and/or immunological activity.

The term “naturally occurring polypeptide” refers to polypeptides produced by cells that have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including, but not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

The term “translated protein coding portion” means a sequence which encodes for the full length protein which may include any leader sequence or any processing sequence.

The term “mature protein coding sequence” means a sequence which encodes a peptide or protein without a signal or leader sequence. The peptide may have been produced by processing in the cell which removes any leader/signal sequence. The peptide may be produced synthetically or the protein may have been produced using a polynucleotide only encoding for the mature protein coding sequence.

The term "derivative" refers to polypeptides chemically modified by such techniques as ubiquitination, labeling (e.g., with radionuclides or various enzymes), covalent polymer attachment such as pegylation (derivatization with polyethylene glycol) and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not normally occur in human proteins.

The term "variant"(or "analog") refers to any polypeptide differing from naturally occurring polypeptides by amino acid insertions, deletions, and substitutions, created using, *e.g.*, recombinant DNA techniques. Guidance in determining which amino acid residues may be replaced, added or deleted without abolishing activities of interest, may be found by comparing the sequence of the particular polypeptide with that of homologous peptides and minimizing the number of amino acid sequence changes made in regions of high homology (conserved regions) or by replacing amino acids with consensus sequence.

Alternatively, recombinant variants encoding these same or similar polypeptides may be synthesized or selected by making use of the "redundancy" in the genetic code. Various codon substitutions, such as the silent changes which produce various restriction sites, may be introduced to optimize cloning into a plasmid or viral vector or expression in a particular prokaryotic or eukaryotic system. Mutations in the polynucleotide sequence may be reflected in the polypeptide or domains of other peptides added to the polypeptide to modify the properties of any part of the polypeptide, to change characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate.

Preferably, amino acid "substitutions" are the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, *i.e.*, conservative amino acid replacements. "Conservative" amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino

acids include aspartic acid and glutamic acid. "Insertions" or "deletions" are preferably in the range of about 1 to 20 amino acids, more preferably 1 to 10 amino acids. The variation allowed may be experimentally determined by systematically making insertions, deletions, or substitutions of amino acids in a polypeptide molecule using recombinant DNA techniques and assaying the resulting recombinant variants for activity.

Alternatively, where alteration of function is desired, insertions, deletions or non-conservative alterations can be engineered to produce altered polypeptides. Such alterations can, for example, alter one or more of the biological functions or biochemical characteristics of the polypeptides of the invention. For example, such alterations may change polypeptide characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate. Further, such alterations can be selected so as to generate polypeptides that are better suited for expression, scale up and the like in the host cells chosen for expression. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

The terms "purified" or "substantially purified" as used herein denotes that the indicated nucleic acid or polypeptide is present in the substantial absence of other biological macromolecules, *e.g.*, polynucleotides, proteins, and the like. In one embodiment, the polynucleotide or polypeptide is purified such that it constitutes at least 95% by weight, more preferably at least 99% by weight, of the indicated biological macromolecules present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 1000 daltons, can be present).

The term "isolated" as used herein refers to a nucleic acid or polypeptide separated from at least one other component (*e.g.*, nucleic acid or polypeptide) present with the nucleic acid or polypeptide in its natural source. In one embodiment, the nucleic acid or polypeptide is found in the presence of (if anything) only a solvent, buffer, ion, or other component normally present in a solution of the same. The terms "isolated" and "purified" do not encompass nucleic acids or polypeptides present in their natural source.

The term "recombinant," when used herein to refer to a polypeptide or protein, means that a polypeptide or protein is derived from recombinant (*e.g.*, microbial, insect, or mammalian) expression systems. "Microbial" refers to recombinant polypeptides or proteins made in bacterial or fungal (*e.g.*, yeast) expression systems. As a product,

"recombinant microbial" defines a polypeptide or protein essentially free of native endogenous substances and unaccompanied by associated native glycosylation.

Polypeptides or proteins expressed in most bacterial cultures, *e.g.*, *E. coli*, will be free of glycosylation modifications; polypeptides or proteins expressed in yeast will have a glycosylation pattern in general different from those expressed in mammalian cells.

The term "recombinant expression vehicle or vector" refers to a plasmid or phage or virus or vector, for expressing a polypeptide from a DNA (RNA) sequence. An expression vehicle can comprise a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an amino terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

The term "recombinant expression system" means host cells which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit extrachromosomally. Recombinant expression systems as defined herein will express heterologous polypeptides or proteins upon induction of the regulatory elements linked to the DNA segment or synthetic gene to be expressed. This term also means host cells which have stably integrated a recombinant genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers. Recombinant expression systems as defined herein will express polypeptides or proteins endogenous to the cell upon induction of the regulatory elements linked to the endogenous DNA segment or gene to be expressed. The cells can be prokaryotic or eukaryotic.

The term "secreted" includes a protein that is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence when it is expressed in a suitable host cell. "Secreted" proteins include without limitation

proteins secreted wholly (*e.g.*, soluble proteins) or partially (*e.g.*, receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins that are transported across the membrane of the endoplasmic reticulum. "Secreted" proteins are also intended to include proteins containing non-typical signal sequences (e.g. Interleukin-1 Beta, see Krasney, P.A. and Young, P.R. (1992) Cytokine 4(2):134-143) and factors released from damaged cells (e.g. Interleukin-1 Receptor Antagonist, see Arend, W.P. et. al. (1998) Annu. Rev. Immunol. 16:27-55)

Where desired, an expression vector may be designed to contain a "signal or leader sequence" which will direct the polypeptide through the membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous protein sources by recombinant DNA techniques.

The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Stringent conditions can include highly stringent conditions (*i.e.*, hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1X SSC/0.1% SDS at 68°C), and moderately stringent conditions (*i.e.*, washing in 0.2X SSC/0.1% SDS at 42°C). Other exemplary hybridization conditions are described herein in the examples.

In instances of hybridization of deoxyoligonucleotides, additional exemplary stringent hybridization conditions include washing in 6X SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligonucleotides), 48°C (for 17-base oligos), 55°C (for 20-base oligonucleotides), and 60°C (for 23-base oligonucleotides).

As used herein, "substantially equivalent" can refer both to nucleotide and amino acid sequences, for example a mutant sequence, that varies from a reference sequence by one or more substitutions, deletions, or additions, the net effect of which does not result in an adverse functional dissimilarity between the reference and subject sequences. Typically, such a substantially equivalent sequence varies from one of those listed herein by no more than about 35% (*i.e.*, the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.35 or less). Such a sequence is said to have 65% sequence identity to the listed sequence. In one embodiment, a substantially

equivalent, *e.g.*, mutant, sequence of the invention varies from a listed sequence by no more than 30% (70% sequence identity); in a variation of this embodiment, by no more than 25% (75% sequence identity); and in a further variation of this embodiment, by no more than 20% (80% sequence identity) and in a further variation of this embodiment, by no more than 10% (90% sequence identity) and in a further variation of this embodiment, by no more than 5% (95% sequence identity). Substantially equivalent, *e.g.*, mutant, amino acid sequences according to the invention preferably have at least 80% sequence identity with a listed amino acid sequence, more preferably at least 90% sequence identity. Substantially equivalent nucleotide sequences of the invention can have lower percent sequence identities, taking into account, for example, the redundancy or degeneracy of the genetic code. Preferably, nucleotide sequence has at least about 65% identity, more preferably at least about 75% identity, and most preferably at least about 95% identity. For the purposes of the present invention, sequences having substantially equivalent biological activity and substantially equivalent expression characteristics are considered substantially equivalent. For the purposes of determining equivalence, truncation of the mature sequence (*e.g.*, via a mutation which creates a spurious stop codon) should be disregarded. Sequence identity may be determined, *e.g.*, using the Jotun Hein method (Hein, J. (1990) *Methods Enzymol.* 183:626-645). Identity between sequences can also be determined by other methods known in the art, *e.g.* by varying hybridization conditions.

The term "totipotent" refers to the capability of a cell to differentiate into all of the cell types of an adult organism.

The term "transformation" means introducing DNA into a suitable host cell so that the DNA is replicable, either as an extrachromosomal element, or by chromosomal integration. The term "transfection" refers to the taking up of an expression vector by a suitable host cell, whether or not any coding sequences are in fact expressed. The term "infection" refers to the introduction of nucleic acids into a suitable host cell by use of a virus or viral vector.

As used herein, an "uptake modulating fragment," UMF, means a series of nucleotides which mediate the uptake of a linked DNA fragment into a cell. UMFs can be readily identified using known UMFs as a target sequence or target motif with the

computer-based systems described below. The presence and activity of a UMF can be confirmed by attaching the suspected UMF to a marker sequence. The resulting nucleic acid molecule is then incubated with an appropriate host under appropriate conditions and the uptake of the marker sequence is determined. As described above, a UMF will increase the frequency of uptake of a linked marker sequence.

Each of the above terms is meant to encompass all that is described for each, unless the context dictates otherwise.

4.2 NUCLEIC ACIDS OF THE INVENTION

Nucleotide sequences of the invention are set forth in the Sequence Listing.

The isolated polynucleotides of the invention include a polynucleotide comprising the nucleotide sequences of the SEQ ID NO: 1 – 35; a polynucleotide encoding any one of the peptide sequences of SEQ ID NO:1 – 35; and a polynucleotide comprising the nucleotide sequence encoding the mature protein coding sequence of the polynucleotides of any one of SEQ ID NO: 1 - 35. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent conditions to (a) the complement of any of the nucleotides sequences of the SEQ ID NO: 1 – 35; (b) nucleotide sequences encoding any one of the amino acid sequences set forth in the Sequence Listing; (c) a polynucleotide which is an allelic variant of any polynucleotide recited above; (d) a polynucleotide which encodes a species homolog of any of the proteins recited above; or (e) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptides of SEQ ID NO: 1-35. Domains of interest may depend on the nature of the encoded polypeptide; e.g., domains in receptor-like polypeptides include ligand-binding, extracellular, transmembrane, or cytoplasmic domains, or combinations thereof; domains in immunoglobulin-like proteins include the variable immunoglobulin-like domains; domains in enzyme-like polypeptides include catalytic and substrate binding domains; and domains in ligand polypeptides include receptor-binding domains.

The polynucleotides of the invention include naturally occurring or wholly or partially synthetic DNA, e.g., cDNA and genomic DNA, and RNA, e.g., mRNA. The

polynucleotides may include all of the coding region of the cDNA or may represent a portion of the coding region of the cDNA.

The present invention also provides genes corresponding to the cDNA sequences disclosed herein. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. Further 5' and 3' sequence can be obtained using methods known in the art. For example, full length cDNA or genomic DNA that corresponds to any of the polynucleotides of the SEQ ID NO: 1 – 35 can be obtained by screening appropriate cDNA or genomic DNA libraries under suitable hybridization conditions using any of the polynucleotides of the SEQ ID NO: 1 - 35 or a portion thereof as a probe. Alternatively, the polynucleotides of the SEQ ID NO: 1 - 35 may be used as the basis for suitable primer(s) that allow identification and/or amplification of genes in appropriate genomic DNA or cDNA libraries.

The nucleic acid sequences of the invention can be assembled from ESTs and sequences (including cDNA and genomic sequences) obtained from one or more public databases, such as dbEST, gbpr, and UniGene. The EST sequences can provide identifying sequence information, representative fragment or segment information, or novel segment information for the full-length gene.

The polynucleotides of the invention also provide polynucleotides including nucleotide sequences that are substantially equivalent to the polynucleotides recited above. Polynucleotides according to the invention can have, *e.g.*, at least about 65%, at least about 70%, at least about 75%, at least about 80%, more typically at least about 90%, and even more typically at least about 95%, sequence identity to a polynucleotide recited above.

Included within the scope of the nucleic acid sequences of the invention are nucleic acid sequence fragments that hybridize under stringent conditions to any of the nucleotide sequences of the SEQ ID NO: 1 - 35, or complements thereof, which fragment is greater than about 5 nucleotides, preferably 7 nucleotides, more preferably greater than 9 nucleotides and most preferably greater than 17 nucleotides. Fragments of, *e.g.* 15, 17, or 20 nucleotides or more that are selective for (*i.e.* specifically hybridize to any one of

the polynucleotides of the invention) are contemplated. Probes capable of specifically hybridizing to a polynucleotide can differentiate polynucleotide sequences of the invention from other polynucleotide sequences in the same family of genes or can differentiate human genes from genes of other species, and are preferably based on unique nucleotide sequences.

The sequences falling within the scope of the present invention are not limited to these specific sequences, but also include allelic and species variations thereof. Allelic and species variations can be routinely determined by comparing the sequence provided in SEQ ID NO: 1 - 35, a representative fragment thereof, or a nucleotide sequence at least 90% identical, preferably 95% identical, to SEQ ID NOs: 1 - 35 with a sequence from another isolate of the same species. Furthermore, to accommodate codon variability, the invention includes nucleic acid molecules coding for the same amino acid sequences as do the specific ORFs disclosed herein. In other words, in the coding region of an ORF, substitution of one codon for another codon that encodes the same amino acid is expressly contemplated.

The nearest neighbor or homology result for the nucleic acids of the present invention, including SEQ ID NOs: 1 - 35, can be obtained by searching a database using an algorithm or a program. Preferably, a BLAST which stands for Basic Local Alignment Search Tool is used to search for local sequence alignments (Altshul, S.F. J Mol. Evol. 36 290-300 (1993) and Altschul S.F. et al. J. Mol. Biol. 21:403-410 (1990)). Alternatively a FASTA version 3 search against Genpept, using Fastxy algorithm.

Species homologs (or orthologs) of the disclosed polynucleotides and proteins are also provided by the present invention. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous or related to that encoded by the polynucleotides.

The nucleic acid sequences of the invention are further directed to sequences which encode variants of the described nucleic acids. These amino acid sequence variants may be prepared by methods known in the art by introducing appropriate

nucleotide changes into a native or variant polynucleotide. There are two variables in the construction of amino acid sequence variants: the location of the mutation and the nature of the mutation. Nucleic acids encoding the amino acid sequence variants are preferably constructed by mutating the polynucleotide to encode an amino acid sequence that does not occur in nature. These nucleic acid alterations can be made at sites that differ in the nucleic acids from different species (variable positions) or in highly conserved regions (constant regions). Sites at such locations will typically be modified in series, *e.g.*, by substituting first with conservative choices (*e.g.*, hydrophobic amino acid to a different hydrophobic amino acid) and then with more distant choices (*e.g.*, hydrophobic amino acid to a charged amino acid), and then deletions or insertions may be made at the target site. Amino acid sequence deletions generally range from about 1 to 30 residues, preferably about 1 to 10 residues, and are typically contiguous. Amino acid insertions include amino- and/or carboxyl-terminal fusions ranging in length from one to one hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions may range generally from about 1 to 10 amino residues, preferably from 1 to 5 residues. Examples of terminal insertions include the heterologous signal sequences necessary for secretion or for intracellular targeting in different host cells and sequences such as FLAG or poly-histidine sequences useful for purifying the expressed protein.

In a preferred method, polynucleotides encoding the novel amino acid sequences are changed via site-directed mutagenesis. This method uses oligonucleotide sequences to alter a polynucleotide to encode the desired amino acid variant, as well as sufficient adjacent nucleotides on both sides of the changed amino acid to form a stable duplex on either side of the site of being changed. In general, the techniques of site-directed mutagenesis are well known to those of skill in the art and this technique is exemplified by publications such as, Edelman et al., *DNA* 2:183 (1983). A versatile and efficient method for producing site-specific changes in a polynucleotide sequence was published by Zoller and Smith, *Nucleic Acids Res.* 10:6487-6500 (1982). PCR may also be used to create amino acid sequence variants of the novel nucleic acids. When small amounts of template DNA are used as starting material, primer(s) that differs slightly in sequence from the corresponding region in the template DNA can generate the desired amino acid

variant. PCR amplification results in a population of product DNA fragments that differ from the polynucleotide template encoding the polypeptide at the position specified by the primer. The product DNA fragments replace the corresponding region in the plasmid and this gives a polynucleotide encoding the desired amino acid variant.

5 A further technique for generating amino acid variants is the cassette mutagenesis technique described in Wells et al., *Gene* 34:315 (1985); and other mutagenesis techniques well known in the art, such as, for example, the techniques in Sambrook et al., supra, and *Current Protocols in Molecular Biology*, Ausubel et al. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the
10 same or a functionally equivalent amino acid sequence may be used in the practice of the invention for the cloning and expression of these novel nucleic acids. Such DNA sequences include those which are capable of hybridizing to the appropriate novel nucleic acid sequence under stringent conditions.

Polynucleotides encoding preferred polypeptide truncations of the invention can
15 be used to generate polynucleotides encoding chimeric or fusion proteins comprising one or more domains of the invention and heterologous protein sequences.

The polynucleotides of the invention additionally include the complement of any of the polynucleotides recited above. The polynucleotide can be DNA (genomic, cDNA, amplified, or synthetic) or RNA. Methods and algorithms for obtaining such
20 polynucleotides are well known to those of skill in the art and can include, for example, methods for determining hybridization conditions that can routinely isolate polynucleotides of the desired sequence identities.

In accordance with the invention, polynucleotide sequences comprising the mature protein coding sequences corresponding to any one of SEQ ID NO: 1-35, or
25 functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression of that nucleic acid, or a functional equivalent thereof, in appropriate host cells. Also included are the cDNA inserts of any of the clones identified herein.

A polynucleotide according to the invention can be joined to any of a variety of
30 other nucleotide sequences by well-established recombinant DNA techniques (see Sambrook J et al. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor

Laboratory, NY). Useful nucleotide sequences for joining to polynucleotides include an assortment of vectors, e.g., plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide.

- 5 In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable marker for the host cell.

Vectors according to the invention include expression vectors, replication vectors, probe generation vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a

- 10 multicellular organism.

The present invention further provides recombinant constructs comprising a nucleic acid having any of the nucleotide sequences of the SEQ ID NOs: 1 - 35 or a fragment thereof or any other polynucleotides of the invention. In one embodiment, the recombinant constructs of the present invention comprise a vector, such as a plasmid or
15 viral vector, into which a nucleic acid having any of the nucleotide sequences of the SEQ ID NOs: 1 - 35 or a fragment thereof is inserted, in a forward or reverse orientation. In the case of a vector comprising one of the ORFs of the present invention, the vector may further comprise regulatory sequences, including for example, a promoter, operably linked to the ORF. Large numbers of suitable vectors and promoters are known to those
20 of skill in the art and are commercially available for generating the recombinant constructs of the present invention. The following vectors are provided by way of example. Bacterial: pBs, phagescript, PsiX174, pBluescript SK, pBs KS, pNH8a, pNH16a, pNH18a, pNH46a (Stratagene); pTrc99A, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLneo, pSV2cat, pOG44, PXTI, pSG (Stratagene)
25 pSVK3, pBPV, pMSG, pSVL (Pharmacia).

The isolated polynucleotide of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art.

- 30 General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein

"operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

5 Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, and trc. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and
10 mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art. Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, *e.g.*, the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream
15 structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic
20 space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an amino terminal identification peptide imparting desired characteristics, *e.g.*, stabilization or simplified purification of expressed recombinant product. Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation
25 initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*,
30 *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM 1 (Promega Biotech, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced or derepressed by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Polynucleotides of the invention can also be used to induce immune responses. For example, as described in Fan et al., *Nat. Biotech.* 17:870-872 (1999), incorporated herein by reference, nucleic acid sequences encoding a polypeptide may be used to generate antibodies against the encoded polypeptide following topical administration of naked plasmid DNA or following injection, and preferably intramuscular injection of the DNA. The nucleic acid sequences are preferably inserted in a recombinant expression vector and may be in the form of naked DNA.

4.3 HOSTS

The present invention further provides host cells genetically engineered to contain the polynucleotides of the invention. For example, such host cells may contain nucleic acids of the invention introduced into the host cell using known transformation, transfection or infection methods. The present invention still further provides host cells genetically engineered to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell.

Knowledge of nucleic acid sequences allows for modification of cells to permit, or increase, expression of endogenous polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased polypeptide expression by replacing, in

whole or in part, the naturally occurring promoter with all or part of a heterologous promoter so that the cells express the polypeptide at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to the encoding sequences. See, for example, PCT International Publication No. WO94/12650, PCT International Publication No. WO92/20808, and PCT International Publication No. WO91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the desired protein coding sequences in the cells.

The host cell can be a higher eukaryotic host cell, such as a mammalian cell, a lower eukaryotic host cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the recombinant construct into the host cell can be effected by calcium phosphate transfection, DEAE, dextran mediated transfection, or electroporation (Davis, L. et al., *Basic Methods in Molecular Biology* (1986)). The host cells containing one of the polynucleotides of the invention, can be used in conventional manners to produce the gene product encoded by the isolated fragment (in the case of an ORF) or can be used to produce a heterologous protein under the control of the EMF.

Any host/vector system can be used to express one or more of the ORFs of the present invention. These include, but are not limited to, eukaryotic hosts such as HeLa cells, Cv-1 cell, COS cells, 293 cells, and Sf9 cells, as well as prokaryotic host such as *E. coli* and *B. subtilis*. The most preferred cells are those which do not normally express the particular polypeptide or protein or which expresses the polypeptide or protein at low natural level. Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, New York (1989), the disclosure of which is hereby incorporated by reference.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell 23:175 (1981). Other cell lines capable of expressing a compatible vector are, for example, the C127, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Recombinant polypeptides and proteins produced in bacterial culture are usually isolated by initial extraction from cell pellets, followed by one or more salting-out, aqueous ion exchange or size exclusion chromatography steps. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or insects or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequence include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the host cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result

in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (gpt) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by reference herein in its entirety.

4.4 POLYPEPTIDES OF THE INVENTION

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising: the amino acid sequences set forth as any one of SEQ ID NO: 1-35 or an amino acid sequence encoded by any one of the nucleotide sequences SEQ ID NOs: 1 - 35 or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides preferably with biological or immunological activity that are encoded by: (a) a polynucleotide having any one of the nucleotide sequences set forth in the SEQ ID NOs: 1 – 35 or (b) polynucleotides encoding any one of the amino acid sequences set forth as SEQ ID NO: 1-35 or (c) polynucleotides that hybridize to the complement of the polynucleotides of either (a) or (b) under stringent hybridization conditions. The invention also provides biologically active or immunologically active variants of any of the amino acid sequences set forth as SEQ ID NO: 1-35 or the corresponding full length or mature protein; and “substantial equivalents” thereof (e.g., with at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, typically at least about 95%, more typically at least about 98%, or most typically at least about 99% amino acid identity) that retain biological activity. Polypeptides encoded by allelic variants may have a similar, increased, or decreased activity compared to polypeptides comprising SEQ ID NO: 1-35.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for

example, as described in H. U. Saragovi, et al., Bio/Technology 10, 773-778 (1992) and in R. S. McDowell, et al., J. Amer. Chem. Soc. 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein

5 binding sites.

The present invention also provides both full-length and mature forms (for example, without a signal sequence or precursor sequence) of the disclosed proteins. The protein coding sequence is identified in the sequence listing by translation of the disclosed nucleotide sequences. The mature form of such protein may be obtained by
10 expression of a full-length polynucleotide in a suitable mammalian cell or other host cell. The sequence of the mature form of the protein is also determinable from the amino acid sequence of the full-length form. Where proteins of the present invention are membrane bound, soluble forms of the proteins are also provided. In such forms, part or all of the regions causing the proteins to be membrane bound are deleted so that the proteins are
15 fully secreted from the cell in which it is expressed.

Protein compositions of the present invention may further comprise an acceptable carrier, such as a hydrophilic, *e.g.*, pharmaceutically acceptable, carrier.

The present invention further provides isolated polypeptides encoded by the nucleic acid fragments of the present invention or by degenerate variants of the nucleic acid fragments of the present invention. By "degenerate variant" is intended nucleotide fragments which differ from a nucleic acid fragment of the present invention (*e.g.*, an ORF) by nucleotide sequence but, due to the degeneracy of the genetic code, encode an identical polypeptide sequence. Preferred nucleic acid fragments of the present invention are the ORFs that encode proteins.

25 A variety of methodologies known in the art can be utilized to obtain any one of the isolated polypeptides or proteins of the present invention. At the simplest level, the amino acid sequence can be synthesized using commercially available peptide synthesizers. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with
30 proteins may possess biological properties in common therewith, including protein activity. This technique is particularly useful in producing small peptides and fragments

of larger polypeptides. Fragments are useful, for example, in generating antibodies against the native polypeptide. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

5 The polypeptides and proteins of the present invention can alternatively be purified from cells which have been altered to express the desired polypeptide or protein. As used herein, a cell is said to be altered to express a desired polypeptide or protein when the cell, through genetic manipulation, is made to produce a polypeptide or protein which it normally does not produce or which the cell normally produces at a lower level.

10 One skilled in the art can readily adapt procedures for introducing and expressing either recombinant or synthetic sequences into eukaryotic or prokaryotic cells in order to generate a cell which produces one of the polypeptides or proteins of the present invention.

15 The invention also relates to methods for producing a polypeptide comprising growing a culture of host cells of the invention in a suitable culture medium, and purifying the protein from the cells or the culture in which the cells are grown. For example, the methods of the invention include a process for producing a polypeptide in which a host cell containing a suitable expression vector that includes a polynucleotide of the invention is cultured under conditions that allow expression of the encoded
20 polypeptide. The polypeptide can be recovered from the culture, conveniently from the culture medium, or from a lysate prepared from the host cells and further purified. Preferred embodiments include those in which the protein produced by such process is a full length or mature form of the protein.

25 In an alternative method, the polypeptide or protein is purified from bacterial cells which naturally produce the polypeptide or protein. One skilled in the art can readily follow known methods for isolating polypeptides and proteins in order to obtain one of the isolated polypeptides or proteins of the present invention. These include, but are not limited to, immunochromatography, HPLC, size-exclusion chromatography, ion-exchange chromatography, and immuno-affinity chromatography. See, *e.g.*, Scopes,
30 *Protein Purification: Principles and Practice*, Springer-Verlag (1994); Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*; Ausubel et al., *Current Protocols in*

Molecular Biology. Polypeptide fragments that retain biological/immunological activity include fragments comprising greater than about 100 amino acids, or greater than about 200 amino acids, and fragments that encode specific protein domains.

The purified polypeptides can be used in *in vitro* binding assays which are well known in the art to identify molecules which bind to the polypeptides. These molecules include but are not limited to, for e.g., small molecules, molecules from combinatorial libraries, antibodies or other proteins. The molecules identified in the binding assay are then tested for antagonist or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

In addition, the peptides of the invention or molecules capable of binding to the peptides may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to cells. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for SEQ ID NO: 1-35.

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications, in the peptide or DNA sequence, can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Pat. No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein. Regions of the protein that are important for the protein function can be determined by various methods known in

the art including the alanine-scanning method which involved systematic substitution of single or strings of amino acids with alanine, followed by testing the resulting alanine-containing variant for biological activity. This type of analysis determines the importance of the substituted amino acid(s) in biological activity. Regions of the protein that are important for protein function may be determined by the eMATRIX program.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and are useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are encompassed by the present invention.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, Calif., U.S.A. (the MaxBat™ kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl™ or Cibacrom blue 3GA Sepharose™; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin

(TRX), or as a His tag. Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, Mass.), Pharmacia (Piscataway, N.J.) and Invitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope.

- 5 One such epitope ("FLAG®") is commercially available from Kodak (New Haven, Conn.).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, *e.g.*, silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or
10 all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The polypeptides of the invention include analogs (variants). This embraces
15 fragments, as well as peptides in which one or more amino acids has been deleted, inserted, or substituted. Also, analogs of the polypeptides of the invention embrace fusions of the polypeptides or modifications of the polypeptides of the invention, wherein the polypeptide or analog is fused to another moiety or moieties, *e.g.*, targeting moiety or another therapeutic agent. Such analogs may exhibit improved properties such as activity
20 and/or stability. Examples of moieties which may be fused to the polypeptide or an analog include, for example, targeting moieties which provide for the delivery of polypeptide to pancreatic cells, *e.g.*, antibodies to pancreatic cells, antibodies to immune cells such as T-cells, monocytes, dendritic cells, granulocytes, etc., as well as receptor and ligands expressed on pancreatic or immune cells. Other moieties which may be
25 fused to the polypeptide include therapeutic agents which are used for treatment, for example, immunosuppressive drugs such as cyclosporin, SK506, azathioprine, CD3 antibodies and steroids. Also, polypeptides may be fused to immune modulators, and other cytokines such as alpha or beta interferon.

30 **4.4.1 DETERMINING POLYPEPTIDE AND POLYNUCLEOTIDE
IDENTITY AND SIMILARITY**

Preferred identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in computer programs including, but are not limited to, the GCG program package, including GAP (Devereux, J., et al., Nucleic Acids Research 12(1):387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, BLASTX, FASTA (Altschul, S.F. et al., J. Molec. Biol. 215:403-410 (1990), PSI-BLAST (Altschul S.F. et al., Nucleic Acids Res. vol. 25, pp. 3389-3402, herein incorporated by reference), eMatrix software (Wu et al., J. Comp. Biol., Vol. 6, pp. 219-235 (1999), herein incorporated by reference), eMotif software (Nevill-Manning et al, ISMB-97, Vol. 4, pp. 202-209, herein incorporated by reference), pFam software (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1), pp. 320-322 (1998), herein incorporated by reference) and the Kyte-Doolittle hydrophobicity prediction algorithm (J. Mol Biol, 157, pp. 105-31 (1982), incorporated herein by reference). The BLAST programs are publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul, S., et al. NCB NLM NIH Bethesda, MD 20894; Altschul, S., et al., J. Mol. Biol. 215:403-410 (1990).

4.5 GENE THERAPY

Mutations in the polynucleotides of the invention gene may result in loss of normal function of the encoded protein. The invention thus provides gene therapy to restore normal activity of the polypeptides of the invention; or to treat disease states involving polypeptides of the invention. Delivery of a functional gene encoding polypeptides of the invention to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly viral vectors (e.g., adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (e.g., liposomes or chemical treatments). See, for example, Anderson, Nature, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, Science, 244: 1275-1281 (1989); Verma, Scientific American: 68-84 (1990); and Miller, Nature, 357: 455-460 (1992). Introduction of any one of the nucleotides of the present invention or a gene encoding the polypeptides of the present invention can also be accomplished with extrachromosomal substrates (transient

expression) or artificial chromosomes (stable expression). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes. Alternatively, it is contemplated that in other human disease states, preventing the expression of or inhibiting the activity of polypeptides of the invention will be useful in treating the disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of polypeptides of the invention.

Other methods inhibiting expression of a protein include the introduction of antisense molecules to the nucleic acids of the present invention, their complements, or their translated RNA sequences, by methods known in the art. Further, the polypeptides of the present invention can be inhibited by using targeted deletion methods, or the insertion of a negative regulatory element such as a silencer, which is tissue specific.

The present invention still further provides cells genetically engineered *in vivo* to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell. These methods can be used to increase or decrease the expression of the polynucleotides of the present invention.

Knowledge of DNA sequences provided by the invention allows for modification of cells to permit, increase, or decrease, expression of endogenous polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased polypeptide expression by replacing, in whole or in part, the naturally occurring promoter with all or part of a heterologous promoter so that the cells express the protein at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to the desired protein encoding sequences. See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the desired

protein coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the desired protein coding sequences in the cells.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences.

Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequences include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting

sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (gpt) gene.

5 The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by
10 reference herein in its entirety.

4.6 TRANSGENIC ANIMALS

 In preferred methods to determine biological functions of the polypeptides of the invention in vivo, one or more genes provided by the invention are either over expressed
15 or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals,
20 preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference. Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals,
25 are produced using methods as described in U.S. Patent No 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

 Transgenic animals can be prepared wherein all or part of a promoter of the polynucleotides of the invention is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using
30 homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased

protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

The polynucleotides of the present invention also make possible the development, through, e.g., homologous recombination or knock out strategies, of animals that fail to express polypeptides of the invention or that express a variant polypeptide. Such animals are useful as models for studying the *in vivo* activities of polypeptide as well as for studying modulators of the polypeptides of the invention.

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference. Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of the polynucleotides of the invention promoter is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

4.7 USES AND BIOLOGICAL ACTIVITY

The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified herein. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or of

5 polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA). The mechanism underlying the particular condition or pathology will dictate whether the polypeptides of the invention, the polynucleotides of the invention or modulators (activators or inhibitors) thereof would be beneficial to the subject in need of treatment. Thus, "therapeutic compositions of the
10 invention" include compositions comprising isolated polynucleotides (including recombinant DNA molecules, cloned genes and degenerate variants thereof) or polypeptides of the invention (including full length protein, mature protein and truncations or domains thereof), or compounds and other substances that modulate the overall activity of the target gene products, either at the level of target gene/protein
15 expression or target protein activity. Such modulators include polypeptides, analogs, (variants), including fragments and fusion proteins, antibodies and other binding proteins; chemical compounds that directly or indirectly activate or inhibit the polypeptides of the invention (identified, e.g., via drug screening assays as described herein); antisense polynucleotides and polynucleotides suitable for triple helix formation; and in particular
20 antibodies or other binding partners that specifically recognize one or more epitopes of the polypeptides of the invention.

The polypeptides of the present invention may likewise be involved in cellular activation or in one of the other physiological pathways described herein.

25 **4.7.1 RESEARCH USES AND UTILITIES**

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either
30 constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on gels; as chromosome markers or tags (when

labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The polypeptides provided by the present invention can similarly be used in assays to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding polypeptide is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology:

Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

4.7.2 NUTRITIONAL USES

Polynucleotides and polypeptides of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the polypeptide or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the polypeptide or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

4.7.3 CYTOKINE AND CELL PROLIFERATION/DIFFERENTIATION

ACTIVITY

A polypeptide of the present invention may exhibit activity relating to cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of therapeutic compositions of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e, CMK, HUVEC, and Caco. Therapeutic compositions of the invention can be used in the following:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19;

Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., I. Immunol. 149:3778-3783, 1992; Bowman et al., I. Immunol. 152:1756-1761, 1994.

5 Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A. M. and Shevach, E. M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human interleukin- , Schreiber, R. D. In Current Protocols
10 in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

 Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L. S. and Lipsky, P. E. In Current
15 Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6--Nordan, R. In
Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley
20 and Sons, Toronto. 1991; Smith et al., Proc. Natl. Aced. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11--Bennett, F., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.15.1 John
Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin
9--Ciarletta, A., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in
25 Immunology. J. E. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

 Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in:
Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H.
30 Margulies, E. M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function; Chapter

6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

5

4.7.4 STEM CELL GROWTH FACTOR ACTIVITY

A polypeptide of the present invention may exhibit stem cell growth factor activity and be involved in the proliferation, differentiation and survival of pluripotent and totipotent stem cells including primordial germ cells, embryonic stem cells, hematopoietic stem cells and/or germ line stem cells. Administration of the polypeptide of the invention to stem cells *in vivo* or *ex vivo* is expected to maintain and expand cell populations in a totipotent or pluripotent state which would be useful for re-engineering damaged or diseased tissues, transplantation, manufacture of bio-pharmaceuticals and the development of bio-sensors. The ability to produce large quantities of human cells has important working applications for the production of human proteins which currently must be obtained from non-human sources or donors, implantation of cells to treat diseases such as Parkinson's, Alzheimer's and other neurodegenerative diseases; tissues for grafting such as bone marrow, skin, cartilage, tendons, bone, muscle (including cardiac muscle), blood vessels, cornea, neural cells, gastrointestinal cells and others; and organs for transplantation such as kidney, liver, pancreas (including islet cells), heart and lung.

It is contemplated that multiple different exogenous growth factors and/or cytokines may be administered in combination with the polypeptide of the invention to achieve the desired effect, including any of the growth factors listed herein, other stem cell maintenance factors, and specifically including stem cell factor (SCF), leukemia inhibitory factor (LIF), Flt-3 ligand (Flt-3L), any of the interleukins, recombinant soluble IL-6 receptor fused to IL-6, macrophage inflammatory protein 1-alpha (MIP-1-alpha), G-CSF, GM-CSF, thrombopoietin (TPO), platelet factor 4 (PF-4), platelet-derived growth factor (PDGF), neural growth factors and basic fibroblast growth factor (bFGF).

Since totipotent stem cells can give rise to virtually any mature cell type, expansion of these cells in culture will facilitate the production of large quantities of

5 mature cells. Techniques for culturing stem cells are known in the art and administration of polypeptides of the invention, optionally with other growth factors and/or cytokines, is expected to enhance the survival and proliferation of the stem cell populations. This can be accomplished by direct administration of the polypeptide of the invention to the culture medium. Alternatively, stroma cells transfected with a polynucleotide that encodes for the polypeptide of the invention can be used as a feeder layer for the stem cell populations in culture or in vivo. Stromal support cells for feeder layers may include embryonic bone marrow fibroblasts, bone marrow stromal cells, fetal liver cells, or cultured embryonic fibroblasts (see U.S. Patent No. 5,690,926).

10 Stem cells themselves can be transfected with a polynucleotide of the invention to induce autocrine expression of the polypeptide of the invention. This will allow for generation of undifferentiated totipotent/pluripotent stem cell lines that are useful as is or that can then be differentiated into the desired mature cell types. These stable cell lines can also serve as a source of undifferentiated totipotent/pluripotent mRNA to create cDNA libraries and templates for polymerase chain reaction experiments. These studies would allow for the isolation and identification of differentially expressed genes in stem cell populations that regulate stem cell proliferation and/or maintenance.

15 Expansion and maintenance of totipotent stem cell populations will be useful in the treatment of many pathological conditions. For example, polypeptides of the present invention may be used to manipulate stem cells in culture to give rise to neuroepithelial cells that can be used to augment or replace cells damaged by illness, autoimmune disease, accidental damage or genetic disorders. The polypeptide of the invention may be useful for inducing the proliferation of neural cells and for the regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders which involve degeneration, death or trauma to neural cells or nerve tissue. In addition, the expanded stem cell populations can also be genetically altered for gene therapy purposes and to decrease host rejection of replacement tissues after grafting or implantation.

25 Expression of the polypeptide of the invention and its effect on stem cells can also be manipulated to achieve controlled differentiation of the stem cells into more differentiated cell types. A broadly applicable method of obtaining pure populations of a

specific differentiated cell type from undifferentiated stem cell populations involves the use of a cell-type specific promoter driving a selectable marker. The selectable marker allows only cells of the desired type to survive. For example, stem cells can be induced to differentiate into cardiomyocytes (Wobus et al., Differentiation, 48: 173-182, (1991); Klug et al., J. Clin. Invest., 98(1): 216-224, (1998)) or skeletal muscle cells (Browder, L. W. In: *Principles of Tissue Engineering* eds. Lanza et al., Academic Press (1997)).

Alternatively, directed differentiation of stem cells can be accomplished by culturing the stem cells in the presence of a differentiation factor such as retinoic acid and an antagonist of the polypeptide of the invention which would inhibit the effects of endogenous stem cell factor activity and allow differentiation to proceed.

In vitro cultures of stem cells can be used to determine if the polypeptide of the invention exhibits stem cell growth factor activity. Stem cells are isolated from any one of various cell sources (including hematopoietic stem cells and embryonic stem cells) and cultured on a feeder layer, as described by Thompson et al. Proc. Natl. Acad. Sci, U.S.A., 92: 7844-7848 (1995), in the presence of the polypeptide of the invention alone or in combination with other growth factors or cytokines. The ability of the polypeptide of the invention to induce stem cells proliferation is determined by colony formation on semi-solid support e.g. as described by Bernstein et al., Blood, 77: 2316-2321 (1991).

4.7.5 HEMATOPOIESIS REGULATING ACTIVITY

A polypeptide of the present invention may be involved in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell disorders. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation

of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

Therapeutic compositions of the invention can be used in the following:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M. G. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I. K. and Briddell, R. A. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, N.Y. 1994; Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R. E. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, N.Y. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R. I.

Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, N.Y. 1994; Long term culture initiating cell assay, Sutherland, H. J. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, N.Y. 1994.

5 **4.7.6 TISSUE GROWTH ACTIVITY**

A polypeptide of the present invention also may be involved in bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as in wound healing and tissue repair and replacement, and in healing of burns, incisions and ulcers.

10 A polypeptide of the present invention which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Compositions of a polypeptide, antibody, binding partner, or other modulator of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an
15 osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

20 A polypeptide of this invention may also be involved in attracting bone-forming cells, stimulating growth of bone-forming cells, or inducing differentiation of progenitors of bone-forming cells. Treatment of osteoporosis, osteoarthritis, bone degenerative disorders, or periodontal disease, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes may also be possible using the composition of the invention.

25 Another category of tissue regeneration activity that may involve the polypeptide of the present invention is tendon/ligament formation. Induction of tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing
30 damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue.

De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The compositions of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a composition may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a composition of the invention.

Compositions of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

Compositions of the present invention may also be involved in the generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising

such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring may allow normal tissue to regenerate. A polypeptide of the present invention may also exhibit angiogenic activity.

A composition of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A composition of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

Therapeutic compositions of the invention can be used in the following:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, H. I. and Rovee, D. T., eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

4.7.7 IMMUNE STIMULATING OR SUPPRESSING ACTIVITY

A polypeptide of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A polynucleotide of the invention can encode a polypeptide exhibiting such activities. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpes

viruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, proteins of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

5 Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitis, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease.

10 Such a protein (or antagonists thereof, including antibodies) of the present invention may also to be useful in the treatment of allergic reactions and conditions (*e.g.*, anaphylaxis, serum sickness, drug reactions, food allergies, insect venom allergies, mastocytosis, allergic rhinitis, hypersensitivity pneumonitis, urticaria, angioedema, eczema, atopic dermatitis, allergic contact dermatitis, erythema multiforme, Stevens-Johnson syndrome, 15 allergic conjunctivitis, atopic keratoconjunctivitis, venereal keratoconjunctivitis, giant papillary conjunctivitis and contact allergies), such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein (or antagonists thereof) of the present invention. The therapeutic effects of the 20 polypeptides or antagonists thereof on allergic reactions can be evaluated by in vivo animals models such as the cumulative contact enhancement test (Lastbom et al., Toxicology 125: 59-66, 1998), skin prick test (Hoffmann et al., Allergy 54: 446-54, 1999), guinea pig skin sensitization test (Vohr et al., Arch. Toxicol. 73: 501-9), and murine local lymph node assay (Kimber et al., J. Toxicol. Environ. Health 53: 563-79).

25 Using the proteins of the invention it may also be possible to modulate immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both.

30 Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent.

Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a therapeutic composition of the invention may prevent cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, a lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular therapeutic compositions in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins in vivo as described in Lenschow et al., *Science* 257:789-792 (1992) and Turka et al., *Proc. Natl. Acad. Sci USA*, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul et al., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of therapeutic compositions of the invention on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms.

Administration of reagents which block stimulation of T cells can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (e.g., a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response may be useful in cases of viral infection, including systemic viral diseases such as influenza, the common cold, and encephalitis.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells in vitro with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the in vitro activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected

cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells in vivo.

A polypeptide of the present invention may provide the necessary stimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient mounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I alpha chain protein and β_2 microglobulin protein or an MHC class II alpha chain protein and an MHC class II beta chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bowman et al., J. Virology 61:1992-1998; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J. J. and Brunswick, M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

5

4.7.8 ACTIVIN/INHIBIN ACTIVITY

A polypeptide of the present invention may also exhibit activin- or inhibin-related activities. A polynucleotide of the invention may encode a polypeptide exhibiting such characteristics. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a polypeptide of the present invention, alone or in heterodimers with a member of the inhibin family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the polypeptide of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, U.S. Pat. No. 4,798,885. A polypeptide of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as, but not limited to, cows, sheep and pigs.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods.

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

30

4.7.9 CHEMOTACTIC/CHEMOKINETIC ACTIVITY

A polypeptide of the present invention may be involved in chemotactic or chemokinetic activity for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Chemotactic and chemokinetic receptor activation can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic compositions (e.g. proteins, antibodies, binding partners, or modulators of the invention) provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

Therapeutic compositions of the invention can be used in the following:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Marguiles, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25:1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153:1762-1768, 1994.

4.7.10 HEMOSTATIC AND THROMBOLYTIC ACTIVITY

A polypeptide of the invention may also be involved in hemostasis or thrombolysis or thrombosis. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Compositions may be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A composition of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

Therapeutic compositions of the invention can be used in the following:

Assays for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

4.7.11 CANCER DIAGNOSIS AND THERAPY

Polypeptides of the invention may be involved in cancer cell generation, proliferation or metastasis. Detection of the presence or amount of polynucleotides or polypeptides of the invention may be useful for the diagnosis and/or prognosis of one or more types of cancer. For example, the presence or increased expression of a polynucleotide/polypeptide of the invention may indicate a hereditary risk of cancer, a precancerous condition, or an ongoing malignancy. Conversely, a defect in the gene or absence of the polypeptide may be associated with a cancer condition. Identification of single nucleotide polymorphisms associated with cancer or a predisposition to cancer may also be useful for diagnosis or prognosis.

Cancer treatments promote tumor regression by inhibiting tumor cell proliferation, inhibiting angiogenesis (growth of new blood vessels that is necessary to support tumor growth) and/or prohibiting metastasis by reducing tumor cell motility or invasiveness. Therapeutic compositions of the invention may be effective in adult and pediatric oncology including in solid phase tumors/malignancies, locally advanced tumors, human soft tissue sarcomas, metastatic cancer, including lymphatic metastases,

blood cell malignancies including multiple myeloma, acute and chronic leukemias, and lymphomas, head and neck cancers including mouth cancer, larynx cancer and thyroid cancer, lung cancers including small cell carcinoma and non-small cell cancers, breast cancers including small cell carcinoma and ductal carcinoma, gastrointestinal cancers including esophageal cancer, stomach cancer, colon cancer, colorectal cancer and polyps associated with colorectal neoplasia, pancreatic cancers, liver cancer, urologic cancers including bladder cancer and prostate cancer, malignancies of the female genital tract including ovarian carcinoma, uterine (including endometrial) cancers, and solid tumor in the ovarian follicle, kidney cancers including renal cell carcinoma, brain cancers including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central nervous system, bone cancers including osteomas, skin cancers including malignant melanoma, tumor progression of human skin keratinocytes, squamous cell carcinoma, basal cell carcinoma, hemangiopericytoma and Karposi's sarcoma.

Polypeptides, polynucleotides, or modulators of polypeptides of the invention (including inhibitors and stimulators of the biological activity of the polypeptide of the invention) may be administered to treat cancer. Therapeutic compositions can be administered in therapeutically effective dosages alone or in combination with adjuvant cancer therapy such as surgery, chemotherapy, radiotherapy, thermotherapy, and laser therapy, and may provide a beneficial effect, e.g. reducing tumor size, slowing rate of tumor growth, inhibiting metastasis, or otherwise improving overall clinical condition, without necessarily eradicating the cancer.

The composition can also be administered in therapeutically effective amounts as a portion of an anti-cancer cocktail. An anti-cancer cocktail is a mixture of the polypeptide or modulator of the invention with one or more anti-cancer drugs in addition to a pharmaceutically acceptable carrier for delivery. The use of anti-cancer cocktails as a cancer treatment is routine. Anti-cancer drugs that are well known in the art and can be used as a treatment in combination with the polypeptide or modulator of the invention include: Actinomycin D, Aminoglutethimide, Asparaginase, Bleomycin, Busulfan, Carboplatin, Carmustine, Chlorambucil, Cisplatin (cis-DDP), Cyclophosphamide, Cytarabine HCl (Cytosine arabinoside), Dacarbazine, Dactinomycin, Daunorubicin HCl,

Doxorubicin HCl, Estramustine phosphate sodium, Etoposide (V16-213), Floxuridine, 5-Fluorouracil (5-Fu), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide, Interferon Alpha-2a, Interferon Alpha-2b, Leuprolide acetate (LHRH-releasing factor analog), Lomustine, Mechlorethamine HCl (nitrogen mustard), Melphalan, Mercaptopurine, Mesna, Methotrexate (MTX), Mitomycin, Mitoxantrone HCl, Octreotide, Plicamycin, Procarbazine HCl, Streptozocin, Tamoxifen citrate, Thioguanine, Thiotepa, Vinblastine sulfate, Vincristine sulfate, Amsacrine, Azacitidine, Hexamethylmelamine, Interleukin-2, Mitoguanzone, Pentostatin, Semustine, Teniposide, and Vindesine sulfate.

In addition, therapeutic compositions of the invention may be used for prophylactic treatment of cancer. There are hereditary conditions and/or environmental situations (e.g. exposure to carcinogens) known in the art that predispose an individual to developing cancers. Under these circumstances, it may be beneficial to treat these individuals with therapeutically effective doses of the polypeptide of the invention to reduce the risk of developing cancers.

In vitro models can be used to determine the effective doses of the polypeptide of the invention as a potential cancer treatment. These *in vitro* models include proliferation assays of cultured tumor cells, growth of cultured tumor cells in soft agar (see Freshney, (1987) Culture of Animal Cells: A Manual of Basic Technique, Wiley-Liss, New York, NY Ch 18 and Ch 21), tumor systems in nude mice as described in Giovanella et al., J. Natl. Can. Inst., 52: 921-30 (1974), mobility and invasive potential of tumor cells in Boyden Chamber assays as described in Pilkington et al., Anticancer Res., 17: 4107-9 (1997), and angiogenesis assays such as induction of vascularization of the chick chorioallantoic membrane or induction of vascular endothelial cell migration as described in Ribatta et al., Intl. J. Dev. Biol., 40: 1189-97 (1999) and Li et al., Clin. Exp. Metastasis, 17:423-9 (1999), respectively. Suitable tumor cells lines are available, e.g. from American Type Tissue Culture Collection catalogs.

4.7.12 RECEPTOR/LIGAND ACTIVITY

A polypeptide of the present invention may also demonstrate activity as receptor, receptor ligand or inhibitor or agonist of receptor/ligand interactions. A polynucleotide of the invention can encode a polypeptide exhibiting such characteristics. Examples of

such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses. Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley- Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1- 7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

By way of example, the polypeptides of the invention may be used as a receptor for a ligand(s) thereby transmitting the biological activity of that ligand(s). Ligands may be identified through binding assays, affinity chromatography, dihybrid screening assays, BIAcore assays, gel overlay assays, or other methods known in the art.

Studies characterizing drugs or proteins as agonist or antagonist or partial agonists or a partial antagonist require the use of other proteins as competing ligands. The polypeptides of the present invention or ligand(s) thereof may be labeled by being coupled to radioisotopes, colorimetric molecules or a toxin molecules by conventional methods. ("Guide to Protein Purification" Murray P. Deutscher (ed) Methods in Enzymology Vol. 182 (1990) Academic Press, Inc. San Diego). Examples of radioisotopes include, but are not limited to, tritium and carbon-14 . Examples of

colorimetric molecules include, but are not limited to, fluorescent molecules such as fluorescamine, or rhodamine or other colorimetric molecules. Examples of toxins include, but are not limited, to ricin.

5 **4.7.13 DRUG SCREENING**

 This invention is particularly useful for screening chemical compounds by using the novel polypeptides or binding fragments thereof in any of a variety of drug screening techniques. The polypeptides or fragments employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. One
10 method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or a fragment thereof. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One
15 may measure, for example, the formation of complexes between polypeptides of the invention or fragments and the agent being tested or examine the diminution in complex formation between the novel polypeptides and an appropriate cell line, which are well known in the art.

 Sources for test compounds that may be screened for ability to bind to or modulate (i.e., increase or decrease) the activity of polypeptides of the invention include
20 (1) inorganic and organic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of either random or mimetic peptides, oligonucleotides or organic molecules.

 Chemical libraries may be readily synthesized or purchased from a number of commercial sources, and may include structural analogs of known compounds or
25 compounds that are identified as "hits" or "leads" via natural product screening.

 The sources of natural product libraries are microorganisms (including bacteria and fungi), animals, plants or other vegetation, or marine organisms, and libraries of mixtures for screening may be created by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of the organisms themselves.
30 Natural product libraries include polyketides, non-ribosomal peptides, and (non-naturally occurring) variants thereof. For a review, see *Science* 282:63-68 (1998).

Combinatorial libraries are composed of large numbers of peptides, oligonucleotides or organic compounds and can be readily prepared by traditional automated synthesis methods, PCR, cloning or proprietary synthetic methods. Of particular interest are peptide and oligonucleotide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created therefrom, see Myers, *Curr. Opin. Biotechnol.* 8:701-707 (1997). For reviews and examples of peptidomimetic libraries, see Al-Obeidi et al., *Mol. Biotechnol.* 9(3):205-23 (1998); Hruby et al., *Curr Opin Chem Biol*, 1(1):114-19 (1997); Dorner et al., *Bioorg Med Chem*, 4(5):709-15 (1996) (alkylated dipeptides).

Identification of modulators through use of the various libraries described herein permits modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to bind a polypeptide of the invention. The molecules identified in the binding assay are then tested for antagonist or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

The binding molecules thus identified may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to cells such as radioisotopes. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for a polypeptide of the invention. Alternatively, the binding molecules may be complexed with imaging agents for targeting and imaging purposes.

4.7.14 ASSAY FOR RECEPTOR ACTIVITY

The invention also provides methods to detect specific binding of a polypeptide e.g. a ligand or a receptor. The art provides numerous assays particularly useful for identifying previously unknown binding partners for receptor polypeptides of the invention. For example, expression cloning using mammalian or bacterial cells, or dihybrid screening assays can be used to identify polynucleotides encoding binding partners. As another example, affinity chromatography with the appropriate immobilized polypeptide of the invention can be used to isolate polypeptides that recognize and bind

polypeptides of the invention. There are a number of different libraries used for the identification of compounds, and in particular small molecules, that modulate (*i.e.*, increase or decrease) biological activity of a polypeptide of the invention. Ligands for receptor polypeptides of the invention can also be identified by adding exogenous ligands, or cocktails of ligands to two cells populations that are genetically identical except for the expression of the receptor of the invention: one cell population expresses the receptor of the invention whereas the other does not. The response of the two cell populations to the addition of ligands(s) are then compared. Alternatively, an expression library can be co-expressed with the polypeptide of the invention in cells and assayed for an autocrine response to identify potential ligand(s). As still another example, BIAcore assays, gel overlay assays, or other methods known in the art can be used to identify binding partner polypeptides, including, (1) organic and inorganic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules.

The role of downstream intracellular signaling molecules in the signaling cascade of the polypeptide of the invention can be determined. For example, a chimeric protein in which the cytoplasmic domain of the polypeptide of the invention is fused to the extracellular portion of a protein, whose ligand has been identified, is produced in a host cell. The cell is then incubated with the ligand specific for the extracellular portion of the chimeric protein, thereby activating the chimeric receptor. Known downstream proteins involved in intracellular signaling can then be assayed for expected modifications *i.e.* phosphorylation. Other methods known to those in the art can also be used to identify signaling molecules involved in receptor activity.

4.7.15 ANTI-INFLAMMATORY ACTIVITY

Compositions of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or

promote an inflammatory response. Compositions with such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation intimation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Compositions of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material. Compositions of this invention may be utilized to prevent or treat conditions such as, but not limited to, sepsis, acute pancreatitis, endotoxin shock, cytokine induced shock, rheumatoid arthritis, chronic inflammatory arthritis, pancreatic cell damage from diabetes mellitus type 1, graft versus host disease, inflammatory bowel disease, inflammation associated with pulmonary disease, other autoimmune disease or inflammatory disease, an antiproliferative agent such as for acute or chronic myelogenous leukemia or in the prevention of premature labor secondary to intrauterine infections.

4.7.16 LEUKEMIAS

Leukemias and related disorders may be treated or prevented by administration of a therapeutic that promotes or inhibits function of the polynucleotides and/or polypeptides of the invention. Such leukemias and related disorders include but are not limited to acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia, chronic leukemia, chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia (for a review of such disorders, see Fishman et al., 1985, Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia).

4.7.17 NERVOUS SYSTEM DISORDERS

Nervous system disorders, involving cell types which can be tested for efficacy of intervention with compounds that modulate the activity of the polynucleotides and/or polypeptides of the invention, and which can be treated upon thus observing an indication of therapeutic utility, include but are not limited to nervous system injuries, and diseases

or disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention include but are not limited to the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems:

(i) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries;

(ii) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia;

(iii) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis;

(iv) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis;

(v) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration;

(vi) neurological lesions associated with systemic diseases including but not limited to diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis;

(vii) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and

(viii) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including but not limited to multiple

sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

Therapeutics which are useful according to the invention for treatment of a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, therapeutics which elicit any of the following effects may be useful according to the invention:

- (i) increased survival time of neurons in culture;
- (ii) increased sprouting of neurons in culture or *in vivo*;
- (iii) increased production of a neuron-associated molecule in culture or *in vivo*, *e.g.*, choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or
- (iv) decreased symptoms of neuron dysfunction *in vivo*.

Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may be measured by the method set forth in Arakawa et al. (1990, J. Neurosci. 10:3507-3515); increased sprouting of neurons may be detected by methods set forth in Pestronk et al. (1980, Exp. Neurol. 70:65-82) or Brown et al. (1981, Ann. Rev. Neurosci. 4:17-42); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, *etc.*, depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, *e.g.*, weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron disorders that may be treated according to the invention include but are not limited to disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including but not limited to progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive

bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

4.7.18 OTHER ACTIVITIES

5 A polypeptide of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue
10 pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, co-factors or other nutritional
15 factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case
20 of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an
25 immune response against such protein or another material or entity which is cross-reactive with such protein.

4.7.19 IDENTIFICATION OF POLYMORPHISMS

The demonstration of polymorphisms makes possible the identification of such polymorphisms in human subjects and the pharmacogenetic use of this information for
30 diagnosis and treatment. Such polymorphisms may be associated with, e.g., differential predisposition or susceptibility to various disease states (such as disorders involving

inflammation or immune response) or a differential response to drug administration, and this genetic information can be used to tailor preventive or therapeutic treatment appropriately. For example, the existence of a polymorphism associated with a predisposition to inflammation or autoimmune disease makes possible the diagnosis of this condition in humans by identifying the presence of the polymorphism.

Polymorphisms can be identified in a variety of ways known in the art which all generally involve obtaining a sample from a patient, analyzing DNA from the sample, optionally involving isolation or amplification of the DNA, and identifying the presence of the polymorphism in the DNA. For example, PCR may be used to amplify an appropriate fragment of genomic DNA which may then be sequenced. Alternatively, the DNA may be subjected to allele-specific oligonucleotide hybridization (in which appropriate oligonucleotides are hybridized to the DNA under conditions permitting detection of a single base mismatch) or to a single nucleotide extension assay (in which an oligonucleotide that hybridizes immediately adjacent to the position of the polymorphism is extended with one or more labeled nucleotides). In addition, traditional restriction fragment length polymorphism analysis (using restriction enzymes that provide differential digestion of the genomic DNA depending on the presence or absence of the polymorphism) may be performed. Arrays with nucleotide sequences of the present invention can be used to detect polymorphisms. The array can comprise modified nucleotide sequences of the present invention in order to detect the nucleotide sequences of the present invention. In the alternative, any one of the nucleotide sequences of the present invention can be placed on the array to detect changes from those sequences.

Alternatively a polymorphism resulting in a change in the amino acid sequence could also be detected by detecting a corresponding change in amino acid sequence of the protein, e.g., by an antibody specific to the variant sequence.

4.7.20 ARTHRITIS AND INFLAMMATION

The immunosuppressive effects of the compositions of the invention against rheumatoid arthritis is determined in an experimental animal model system. The experimental model system is adjuvant induced arthritis in rats, and the protocol is described by J. Holoshitz, et al., 1983, Science, 219:56, or by B. Waksman et al., 1963,

Int. Arch. Allergy Appl. Immunol., 23:129. Induction of the disease can be caused by a single injection, generally intradermally, of a suspension of killed Mycobacterium tuberculosis in complete Freund's adjuvant (CFA). The route of injection can vary, but rats may be injected at the base of the tail with an adjuvant mixture. The polypeptide is administered in phosphate buffered solution (PBS) at a dose of about 1-5 mg/kg. The control consists of administering PBS only.

The procedure for testing the effects of the test compound would consist of intradermally injecting killed Mycobacterium tuberculosis in CFA followed by immediately administering the test compound and subsequent treatment every other day until day 24. At 14, 15, 18, 20, 22, and 24 days after injection of Mycobacterium CFA, an overall arthritis score may be obtained as described by J. Holoskitz above. An analysis of the data would reveal that the test compound would have a dramatic affect on the swelling of the joints as measured by a decrease of the arthritis score.

4.8 THERAPEUTIC METHODS

The compositions (including polypeptide fragments, analogs, variants and antibodies or other binding partners or modulators including antisense polynucleotides) of the invention have numerous applications in a variety of therapeutic methods. Examples of therapeutic applications include, but are not limited to, those exemplified herein.

4.8.1 EXAMPLE

One embodiment of the invention is the administration of an effective amount of the polypeptides or other composition of the invention to individuals affected by a disease or disorder that can be modulated by regulating the peptides of the invention. While the mode of administration is not particularly important, parenteral administration is preferred. An exemplary mode of administration is to deliver an intravenous bolus. The dosage of the polypeptides or other composition of the invention will normally be determined by the prescribing physician. It is to be expected that the dosage will vary according to the age, weight, condition and response of the individual patient. Typically,

the amount of polypeptide administered per dose will be in the range of about 0.01µg/kg to 100 mg/kg of body weight, with the preferred dose being about 0.1µg/kg to 10 mg/kg of patient body weight. For parenteral administration, polypeptides of the invention will be formulated in an injectable form combined with a pharmaceutically acceptable parenteral vehicle. Such vehicles are well known in the art and examples include water, saline, Ringer's solution, dextrose solution, and solutions consisting of small amounts of the human serum albumin. The vehicle may contain minor amounts of additives that maintain the isotonicity and stability of the polypeptide or other active ingredient. The preparation of such solutions is within the skill of the art.

4.9 PHARMACEUTICAL FORMULATIONS AND ROUTES OF ADMINISTRATION

A protein or other composition of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources and including antibodies and other binding partners of the polypeptides of the invention) may be administered to a patient in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders. Such a composition may optionally contain (in addition to protein or other active ingredient and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the disease or disorder in question. These agents include various growth factors such as epidermal growth factor (EGF),

platelet-derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), insulin-like growth factor (IGF), as well as cytokines described herein.

The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or other active ingredient or complement its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein or other active ingredient of the invention, or to minimize side effects. Conversely, protein or other active ingredient of the present invention may be included in formulations of the particular clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent (such as IL-1Ra, IL-1 Hy1, IL-1 Hy2, anti-TNF, corticosteroids, immunosuppressive agents). A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

As an alternative to being included in a pharmaceutical composition of the invention including a first protein, a second protein or a therapeutic agent may be concurrently administered with the first protein (e.g., at the same time, or at differing times provided that therapeutic concentrations of the combination of agents is achieved at the treatment site). Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms, e.g., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When applied to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein or other active ingredient of the present invention is administered to a mammal having a condition to be treated. Protein or other active ingredient of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein or other active ingredient of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein or other active ingredient of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

4.9.1 ROUTES OF ADMINISTRATION

Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. Administration of protein or other active ingredient of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a arthritic joints or in fibrotic tissue, often in a depot or sustained release formulation. In order to prevent the scarring process frequently occurring as complication of glaucoma surgery, the compounds may be administered topically, for example, as eye drops. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome

coated with a specific antibody, targeting, for example, arthritic or fibrotic tissue. The liposomes will be targeted to and taken up selectively by the afflicted tissue.

The polypeptides of the invention are administered by any route that delivers an effective dosage to the desired site of action. The determination of a suitable route of administration and an effective dosage for a particular indication is within the level of skill in the art. Preferably for wound treatment, one administers the therapeutic compound directly to the site. Suitable dosage ranges for the polypeptides of the invention can be extrapolated from these dosages or from similar studies in appropriate animal models. Dosages can then be adjusted as necessary by the clinician to provide maximal therapeutic benefit.

4.9.2 COMPOSITIONS/FORMULATIONS

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. These pharmaceutical compositions may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of protein or other active ingredient of the present invention is administered orally, protein or other active ingredient of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein or other active ingredient of the present invention, and preferably from about 25 to 90% protein or other active ingredient of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol,

propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein or other active ingredient of the present invention, and preferably from about 1 to 50% protein or other active ingredient of the present invention.

5 When a therapeutically effective amount of protein or other active ingredient of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein or other active ingredient of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein or other active ingredient solutions, having due regard to
10 pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein or other active ingredient of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other
15 vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal
20 administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills,
25 dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained from a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including
30 lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose,

hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose,

5 concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

10 Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds
15 may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

20 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be
25 determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch. The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit
30 dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or

aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a co-solvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The co-solvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without

destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g. polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose. Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity.

Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various types of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein or other active ingredient stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Many of the active ingredients of the invention may be provided as salts with pharmaceutically compatible counter ions. Such pharmaceutically acceptable base addition salts are those salts which retain the biological effectiveness and properties of the free acids and which are obtained by reaction with inorganic or organic bases such as sodium hydroxide, magnesium hydroxide, ammonia, trialkylamine, dialkylamine, monoalkylamine, dibasic amino acids, sodium acetate, potassium benzoate, triethanol amine and the like.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) or other active ingredient(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T

cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with
5 co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a
10 liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithins, phospholipids, saponin, bile acids,
15 and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323, all of which are incorporated herein by reference.

The amount of protein or other active ingredient of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and
20 severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein or other active ingredient of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein or other active ingredient of the present invention and observe the patient's response.

25 Larger doses of protein or other active ingredient of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01 μ g to about 100 mg (preferably about 0.1 μ g to about 10 mg, more
30 preferably about 0.1 μ g to about 1 mg) of protein or other active ingredient of the present invention per kg body weight. For compositions of the present invention which are

useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein or other active ingredient of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing or other active ingredient-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability. Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering

agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt %, preferably 1-10 wt % based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells. In further compositions, proteins or other active ingredients of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins or other active ingredients of the present invention. The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final

composition, may also effect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either in vivo or ex vivo into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA). Cells may also be cultured ex vivo in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced in vivo for therapeutic purposes.

4.9.3 EFFECTIVE DOSAGE

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from appropriate in vitro assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that can be used to more accurately determine useful doses in humans. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of the protein's biological activity). Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical

procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀.

- 5 Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See, *e.g.*, Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1. Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the desired effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

- 20 Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

- 25 An exemplary dosage regimen for polypeptides or other compositions of the invention will be in the range of about 0.01 µg/kg to 100 mg/kg of body weight daily, with the preferred dose being about 0.1 µg/kg to 25 mg/kg of patient body weight daily, varying in adults and children. Dosing may be once daily, or equivalent doses may be delivered at longer or shorter intervals.

- 30 The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's age and weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

4.9.4 PACKAGING

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

4.10 ANTIBODIES

Another aspect of the invention is an antibody that specifically binds the polypeptide of the invention. Such antibodies include monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, and complementary determining region (CDR)-grafted antibodies, including compounds which include CDR and/or antigen-binding sequences, which specifically recognize a polypeptide of the invention. Preferred antibodies of the invention are human antibodies which are produced and identified according to methods described in WO93/11236, published June 20, 1993, which is incorporated herein by reference in its entirety. Antibody fragments, including Fab, Fab', F(ab')₂, and F_v, are also provided by the invention. The term "specific for" indicates that the variable regions of the antibodies of the invention recognize and bind polypeptides of the invention exclusively (*i.e.*, able to distinguish the polypeptide of the invention from other similar polypeptides despite sequence identity, homology, or similarity found in the family of polypeptides), but may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow et al. (Eds), *Antibodies A Laboratory Manual*; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988),

Chapter 6. Antibodies that recognize and bind fragments of the polypeptides of the invention are also contemplated, provided that the antibodies are first and foremost specific for, as defined above, full length polypeptides of the invention. As with antibodies that are specific for full length polypeptides of the invention, antibodies of the invention that recognize fragments are those which can distinguish polypeptides from the same family of polypeptides despite inherent sequence identity, homology, or similarity found in the family of proteins. Antibodies of the invention can be produced using any method well known and routinely practiced in the art.

Non-human antibodies may be humanized by any methods known in the art. In one method, the non-human CDRs are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

Antibodies of the invention are useful for, for example, therapeutic purposes (by modulating activity of a polypeptide of the invention), diagnostic purposes to detect or quantitate a polypeptide of the invention, as well as purification of a polypeptide of the invention. Kits comprising an antibody of the invention for any of the purposes described herein are also comprehended. In general, a kit of the invention also includes a control antigen for which the antibody is immunospecific. The invention further provides a hybridoma that produces an antibody according to the invention. Antibodies of the invention are useful for detection and/or purification of the polypeptides of the invention.

Polypeptides of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R. P. Merrifield, J. Amer. Chem. Soc. 85, 2149-2154 (1963); J. L. Krstenansky, et al., FEBS Lett. 211, 10 (1987).

Monoclonal antibodies binding to the protein of the invention may be useful diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions

associated with the protein and also in the treatment of some forms of cancer where abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein. In general, techniques for preparing polyclonal and monoclonal antibodies as well as hybridomas capable of producing the desired antibody are well known in the art (Campbell, A.M., Monoclonal Antibodies Technology: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1984); St. Groth et al., J. Immunol. 35:1-21 (1990); Kohler and Milstein, Nature 256:495-497 (1975)), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., Immunology Today 4:72 (1983); Cole et al., in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc. (1985), pp. 77-96).

Any animal (mouse, rabbit, *etc.*) which is known to produce antibodies can be immunized with a peptide or polypeptide of the invention. Methods for immunization are well known in the art. Such methods include subcutaneous or intraperitoneal injection of the polypeptide. One skilled in the art will recognize that the amount of the protein encoded by the ORF of the present invention used for immunization will vary based on the animal which is immunized, the antigenicity of the peptide and the site of injection. The protein that is used as an immunogen may be modified or administered in an adjuvant in order to increase the protein's antigenicity. Methods of increasing the antigenicity of a protein are well known in the art and include, but are not limited to, coupling the antigen with a heterologous protein (such as globulin or β -galactosidase) or through the inclusion of an adjuvant during immunization.

For monoclonal antibodies, spleen cells from the immunized animals are removed, fused with myeloma cells, such as SP2/0-Ag14 myeloma cells, and allowed to become monoclonal antibody producing hybridoma cells. Any one of a number of methods well known in the art can be used to identify the hybridoma cell which produces an antibody with the desired characteristics. These include screening the hybridomas with an ELISA assay, Western blot analysis, or radioimmunoassay (Lutz et al., Exp. Cell Research. 175:109-124 (1988)). Hybridomas secreting the desired antibodies are cloned and the class and subclass is determined using procedures known in the art (Campbell,

A.M., Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1984)).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to proteins of the present invention.

For polyclonal antibodies, antibody-containing antiserum is isolated from the immunized animal and is screened for the presence of antibodies with the desired specificity using one of the above-described procedures. The present invention further provides the above-described antibodies in delectably labeled form. Antibodies can be delectably labeled through the use of radioisotopes, affinity labels (such as biotin, avidin, etc.), enzymatic labels (such as horseradish peroxidase, alkaline phosphatase, etc.) fluorescent labels (such as FITC or rhodamine, etc.), paramagnetic atoms, etc. Procedures for accomplishing such labeling are well-known in the art, for example, see (Sternberger, L.A. et al., J. Histochem. Cytochem. 18:315 (1970); Bayer, E.A. et al., Meth. Enzym. 62:308 (1979); Engval, E. et al., Immunol. 109:129 (1972); Goding, J.W. J. Immunol. Meth. 13:215 (1976)).

The labeled antibodies of the present invention can be used for *in vitro*, *in vivo*, and *in situ* assays to identify cells or tissues in which a fragment of the polypeptide of interest is expressed. The antibodies may also be used directly in therapies or other diagnostics. The present invention further provides the above-described antibodies immobilized on a solid support. Examples of such solid supports include plastics such as polycarbonate, complex carbohydrates such as agarose and Sepharose®, acrylic resins and such as polyacrylamide and latex beads. Techniques for coupling antibodies to such solid supports are well known in the art (Weir, D.M. et al., "Handbook of Experimental Immunology" 4th Ed., Blackwell Scientific Publications, Oxford, England, Chapter 10 (1986); Jacoby, W.D. et al., Meth. Enzym. 34 Academic Press, N.Y. (1974)). The immobilized antibodies of the present invention can be used for *in vitro*, *in vivo*, and *in situ* assays as well as for immuno-affinity purification of the proteins of the present invention.

4.11 COMPUTER READABLE SEQUENCES

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium which can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence of the present invention. As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate manufactures comprising the nucleotide sequence information of the present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (*e.g.* text file or database) in order to obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing any of the nucleotide sequences SEQ ID NOs: 1 - 35 or a representative fragment thereof; or a nucleotide sequence at least 95% identical to any of the nucleotide sequences of the SEQ ID NOs: 1 - 35 in computer readable form, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence

information provided in a computer readable medium. The examples which follow demonstrate how software which implements the BLAST (Altschul et al., J. Mol. Biol. 215:403-410 (1990)) and BLAZE (Brutlag et al., Comp. Chem. 17:203-207 (1993)) search algorithms on a Sybase system is used to identify open reading frames (ORFs) within a nucleic acid sequence. Such ORFs may be protein encoding fragments and may be useful in producing commercially important proteins such as enzymes used in fermentation reactions and in the production of commercially useful metabolites.

As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are suitable for use in the present invention. As stated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention.

As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of a known sequence which match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, Smith-Waterman, MacPattern (EMBL), BLASTN and BLASTA (NPOLYPEPTIDEIA). A skilled artisan can readily recognize that any one of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present computer-based

systems. As used herein, a "target sequence" can be any nucleic acid or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 300 amino acids, more preferably from about 30 to 100 nucleotide residues. However, it is well recognized that searches for commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

4.12 TRIPLE HELIX FORMATION

In addition, the fragments of the present invention, as broadly described, can be used to control gene expression through triple helix formation or antisense DNA or RNA, both of which methods are based on the binding of a polynucleotide sequence to DNA or RNA. Polynucleotides suitable for use in these methods are preferably 20 to 40 bases in length and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 15241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Olmno, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide.

4.13 DIAGNOSTIC ASSAYS AND KITS

The present invention further provides methods to identify the presence or expression of one of the ORFs of the present invention, or homolog thereof, in a test sample, using a nucleic acid probe or antibodies of the present invention, optionally conjugated or otherwise associated with a suitable label.

In general, methods for detecting a polynucleotide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polynucleotide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polynucleotide of the invention is detected in the sample. Such methods can also comprise contacting a sample under stringent hybridization conditions with nucleic acid primers that anneal to a polynucleotide of the invention under such conditions, and amplifying annealed polynucleotides, so that if a polynucleotide is amplified, a polynucleotide of the invention is detected in the sample.

In general, methods for detecting a polypeptide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polypeptide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polypeptide of the invention is detected in the sample.

In detail, such methods comprise incubating a test sample with one or more of the antibodies or one or more of the nucleic acid probes of the present invention and assaying for binding of the nucleic acid probes or antibodies to components within the test sample.

Conditions for incubating a nucleic acid probe or antibody with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid probe or antibody used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or immunological assay formats can readily be adapted to employ the nucleic acid probes or antibodies of the present invention. Examples of such assays can be found in Chard, T., *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G.R. et al., *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of immunoassays*:

Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1985). The test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as sputum, blood, serum, plasma, or urine. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention. Specifically, the invention provides a compartment kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the probes or antibodies of the present invention; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound probe or antibody.

In detail, a compartment kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers or strips of plastic or paper. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the antibodies used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound antibody or probe. Types of detection reagents include labeled nucleic acid probes, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. One skilled in the art will readily recognize that the disclosed probes and antibodies of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

4.14 MEDICAL IMAGING

The novel polypeptides and binding partners of the invention are useful in medical imaging of sites expressing the molecules of the invention (e.g., where the polypeptide of the invention is involved in the immune response, for imaging sites of inflammation or infection). See, e.g., Kunkel et al., U.S. Pat. NO. 5,413,778. Such methods involve chemical attachment of a labeling or imaging agent, administration of the labeled polypeptide to a subject in a pharmaceutically acceptable carrier, and imaging the labeled polypeptide *in vivo* at the target site.

4.15 SCREENING ASSAYS

Using the isolated proteins and polynucleotides of the invention, the present invention further provides methods of obtaining and identifying agents which bind to a polypeptide encoded by an ORF corresponding to any of the nucleotide sequences set forth in the SEQ ID NOs: 1 - 35, or bind to a specific domain of the polypeptide encoded by the nucleic acid. In detail, said method comprises the steps of:

- (a) contacting an agent with an isolated protein encoded by an ORF of the present invention, or nucleic acid of the invention; and
- (b) determining whether the agent binds to said protein or said nucleic acid.

In general, therefore, such methods for identifying compounds that bind to a polynucleotide of the invention can comprise contacting a compound with a polynucleotide of the invention for a time sufficient to form a polynucleotide/compound complex, and detecting the complex, so that if a polynucleotide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

Likewise, in general, therefore, such methods for identifying compounds that bind to a polypeptide of the invention can comprise contacting a compound with a polypeptide of the invention for a time sufficient to form a polypeptide/compound complex, and detecting the complex, so that if a polypeptide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

Methods for identifying compounds that bind to a polypeptide of the invention can also comprise contacting a compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound complex, wherein the complex drives

expression of a receptor gene sequence in the cell, and detecting the complex by detecting reporter gene sequence expression, so that if a polypeptide/compound complex is detected, a compound that binds a polypeptide of the invention is identified.

Compounds identified via such methods can include compounds which modulate the activity of a polypeptide of the invention (that is, increase or decrease its activity, relative to activity observed in the absence of the compound). Alternatively, compounds identified via such methods can include compounds which modulate the expression of a polynucleotide of the invention (that is, increase or decrease expression relative to expression levels observed in the absence of the compound). Compounds, such as compounds identified via the methods of the invention, can be tested using standard assays well known to those of skill in the art for their ability to modulate activity/expression.

The agents screened in the above assay can be, but are not limited to, peptides, carbohydrates, vitamin derivatives, or other pharmaceutical agents. The agents can be selected and screened at random or rationally selected or designed using protein modeling techniques.

For random screening, agents such as peptides, carbohydrates, pharmaceutical agents and the like are selected at random and are assayed for their ability to bind to the protein encoded by the ORF of the present invention. Alternatively, agents may be rationally selected or designed. As used herein, an agent is said to be "rationally selected or designed" when the agent is chosen based on the configuration of the particular protein. For example, one skilled in the art can readily adapt currently available procedures to generate peptides, pharmaceutical agents and the like, capable of binding to a specific peptide sequence, in order to generate rationally designed antipeptide peptides, for example see Hurby et al., Application of Synthetic Peptides: Antisense Peptides," In Synthetic Peptides, A User's Guide, W.H. Freeman, NY (1992), pp. 289-307, and Kaspczak et al., Biochemistry 28:9230-8 (1989), or pharmaceutical agents, or the like.

In addition to the foregoing, one class of agents of the present invention, as broadly described, can be used to control gene expression through binding to one of the ORFs or EMFs of the present invention. As described above, such agents can be randomly screened or rationally designed/selected. Targeting the ORF or EMF allows a

skilled artisan to design sequence specific or element specific agents, modulating the expression of either a single ORF or multiple ORFs which rely on the same EMF for expression control. One class of DNA binding agents are agents which contain base residues which hybridize or form a triple helix formation by binding to DNA or RNA.

- 5 Such agents can be based on the classic phosphodiester, ribonucleic acid backbone, or can be a variety of sulfhydryl or polymeric derivatives which have base attachment capacity.

Agents suitable for use in these methods preferably contain 20 to 40 bases and are designed to be complementary to a region of the gene involved in transcription (triple
10 helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization
15 blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide and other DNA binding agents.

Agents which bind to a protein encoded by one of the ORFs of the present
20 invention can be used as a diagnostic agent. Agents which bind to a protein encoded by one of the ORFs of the present invention can be formulated using known techniques to generate a pharmaceutical composition.

4.16 USE OF NUCLEIC ACIDS AS PROBES

25 Another aspect of the subject invention is to provide for polypeptide-specific nucleic acid hybridization probes capable of hybridizing with naturally occurring nucleotide sequences. The hybridization probes of the subject invention may be derived from any of the nucleotide sequences SEQ ID NOs: 1 - 35. Because the corresponding gene is only expressed in a limited number of tissues, a hybridization probe derived from
30 of any of the nucleotide sequences SEQ ID NOs: 1 - 35 can be used as an indicator of the presence of RNA of cell type of such a tissue in a sample.

Any suitable hybridization technique can be employed, such as, for example, in situ hybridization. PCR as described in US Patents Nos. 4,683,195 and 4,965,188 provides additional uses for oligonucleotides based upon the nucleotide sequences. Such probes used in PCR may be of recombinant origin, may be chemically synthesized, or a mixture of both. The probe will comprise a discrete nucleotide sequence for the detection of identical sequences or a degenerate pool of possible sequences for identification of closely related genomic sequences.

Other means for producing specific hybridization probes for nucleic acids include the cloning of nucleic acid sequences into vectors for the production of mRNA probes.

Such vectors are known in the art and are commercially available and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides. The nucleotide sequences may be used to construct hybridization probes for mapping their respective genomic sequences. The nucleotide sequence provided herein may be mapped to a chromosome or specific regions of a chromosome using well known genetic and/or chromosomal mapping techniques. These techniques include in situ hybridization, linkage analysis against known chromosomal markers, hybridization screening with libraries or flow-sorted chromosomal preparations specific to known chromosomes, and the like. The technique of fluorescent in situ hybridization of chromosome spreads has been described, among other places, in Verma et al (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY.

Fluorescent *in situ* hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of a nucleic acid on a physical chromosomal map and a specific disease (or predisposition to a specific disease) may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals.

4.17 PREPARATION OF SUPPORT BOUND OLIGONUCLEOTIDES

Oligonucleotides, i.e., small nucleic acid segments, may be readily prepared by, for example, directly synthesizing the oligonucleotide by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer.

Support bound oligonucleotides may be prepared by any of the methods known to those of skill in the art using any suitable support such as glass, polystyrene or Teflon. One strategy is to precisely spot oligonucleotides synthesized by standard synthesizers. Immobilization can be achieved using passive adsorption (Inouye & Hondo, (1990) J. Clin. Microbiol. 28(6) 1469-72); using UV light (Nagata *et al.*, 1985; Dahlen *et al.*, 1987; Morrissey & Collins, (1989) Mol. Cell Probes 3(2) 189-207) or by covalent binding of base modified DNA (Keller *et al.*, 1988; 1989); all references being specifically incorporated herein.

Another strategy that may be employed is the use of the strong biotin-streptavidin interaction as a linker. For example, Broude *et al.* (1994) Proc. Natl. Acad. Sci. USA 91(8) 3072-6, describe the use of biotinylated probes, although these are duplex probes, that are immobilized on streptavidin-coated magnetic beads. Streptavidin-coated beads may be purchased from Dynal, Oslo. Of course, this same linking chemistry is applicable to coating any surface with streptavidin. Biotinylated probes may be purchased from various sources, such as, e.g., Operon Technologies (Alameda, CA).

Nunc Laboratories (Naperville, IL) is also selling suitable material that could be used. Nunc Laboratories have developed a method by which DNA can be covalently bound to the microwell surface termed CovaLink NH. CovaLink NH is a polystyrene surface grafted with secondary amino groups (>NH) that serve as bridge-heads for further covalent coupling. CovaLink Modules may be purchased from Nunc Laboratories. DNA molecules may be bound to CovaLink exclusively at the 5'-end by a phosphoramidate bond, allowing immobilization of more than 1 pmol of DNA (Rasmussen *et al.*, (1991) Anal. Biochem. 198(1) 138-42).

The use of CovaLink NH strips for covalent binding of DNA molecules at the 5'-end has been described (Rasmussen *et al.*, (1991). In this technology, a phosphoramidate bond is employed (Chu *et al.*, (1983) Nucleic Acids Res. 11(8) 6513-29). This is beneficial as immobilization using only a single covalent bond is preferred. The phosphoramidate bond

joins the DNA to the CovaLink NH secondary amino groups that are positioned at the end of spacer arms covalently grafted onto the polystyrene surface through a 2 nm long spacer arm. To link an oligonucleotide to CovaLink NH via an phosphoramidate bond, the oligonucleotide terminus must have a 5'-end phosphate group. It is, perhaps, even possible for biotin to be covalently bound to CovaLink and then streptavidin used to bind the probes.

More specifically, the linkage method includes dissolving DNA in water (7.5 ng/ul) and denaturing for 10 min. at 95°C and cooling on ice for 10 min. Ice-cold 0.1 M 1-methylimidazole, pH 7.0 (1-MeIm₇), is then added to a final concentration of 10 mM 1-MeIm₇. A ss DNA solution is then dispensed into CovaLink NH strips (75 ul/well) standing on ice.

Carbodiimide 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC), dissolved in 10 mM 1-MeIm₇, is made fresh and 25 ul added per well. The strips are incubated for 5 hours at 50°C. After incubation the strips are washed using, e.g., Nunc-Immuno Wash; first the wells are washed 3 times, then they are soaked with washing solution for 5 min., and finally they are washed 3 times (where in the washing solution is 0.4 N NaOH, 0.25% SDS heated to 50°C).

It is contemplated that a further suitable method for use with the present invention is that described in PCT Patent Application WO 90/03382 (Southern & Maskos), incorporated herein by reference. This method of preparing an oligonucleotide bound to a support involves attaching a nucleoside 3'-reagent through the phosphate group by a covalent phosphodiester link to aliphatic hydroxyl groups carried by the support. The oligonucleotide is then synthesized on the supported nucleoside and protecting groups removed from the synthetic oligonucleotide chain under standard conditions that do not cleave the oligonucleotide from the support. Suitable reagents include nucleoside phosphoramidite and nucleoside hydrogen phosphorate.

An on-chip strategy for the preparation of DNA probe for the preparation of DNA probe arrays may be employed. For example, addressable laser-activated photodeprotection may be employed in the chemical synthesis of oligonucleotides directly on a glass surface, as described by Fodor *et al.* (1991) Science 251(4995) 767-73, incorporated herein by reference. Probes may also be immobilized on nylon supports as described by Van Ness *et al.* (1991) Nucleic Acids Res. 19(12) 3345-50; or linked to Teflon using the method of

Duncan & Cavalier (1988) Anal. Biochem. 169(1) 104-8; all references being specifically incorporated herein.

To link an oligonucleotide to a nylon support, as described by Van Ness *et al.* (1991), requires activation of the nylon surface via alkylation and selective activation of the 5'-amine of oligonucleotides with cyanuric chloride.

One particular way to prepare support bound oligonucleotides is to utilize the light-generated synthesis described by Pease *et al.*, (1994) PNAS USA 91(11) 5022-6, incorporated herein by reference). These authors used current photolithographic techniques to generate arrays of immobilized oligonucleotide probes (DNA chips). These methods, in which light is used to direct the synthesis of oligonucleotide probes in high-density, miniaturized arrays, utilize photolabile 5'-protected *N*-acyl-deoxynucleoside phosphoramidites, surface linker chemistry and versatile combinatorial synthesis strategies. A matrix of 256 spatially defined oligonucleotide probes may be generated in this manner.

4.18 PREPARATION OF NUCLEIC ACID FRAGMENTS

The nucleic acids may be obtained from any appropriate source, such as cDNAs, genomic DNA, chromosomal DNA, microdissected chromosome bands, cosmid or YAC inserts, and RNA, including mRNA without any amplification steps. For example, Sambrook *et al.* (1989) describes three protocols for the isolation of high molecular weight DNA from mammalian cells (p. 9.14-9.23).

DNA fragments may be prepared as clones in M13, plasmid or lambda vectors and/or prepared directly from genomic DNA or cDNA by PCR or other amplification methods. Samples may be prepared or dispensed in multiwell plates. About 100-1000 ng of DNA samples may be prepared in 2-500 ml of final volume.

The nucleic acids would then be fragmented by any of the methods known to those of skill in the art including, for example, using restriction enzymes as described at 9.24-9.28 of Sambrook *et al.* (1989), shearing by ultrasound and NaOH treatment.

Low pressure shearing is also appropriate, as described by Schriefer *et al.* (1990) Nucleic Acids Res. 18(24) 7455-6, incorporated herein by reference). In this method, DNA samples are passed through a small French pressure cell at a variety of low to intermediate pressures. A lever device allows controlled application of low to intermediate pressures to

the cell. The results of these studies indicate that low-pressure shearing is a useful alternative to sonic and enzymatic DNA fragmentation methods.

One particularly suitable way for fragmenting DNA is contemplated to be that using the two base recognition endonuclease, *CviJI*, described by Fitzgerald *et al.* (1992) Nucleic Acids Res. 20(14) 3753-62. These authors described an approach for the rapid fragmentation and fractionation of DNA into particular sizes that they contemplated to be suitable for shotgun cloning and sequencing.

The restriction endonuclease *CviJI* normally cleaves the recognition sequence PuGCPy between the G and C to leave blunt ends. Atypical reaction conditions, which alter the specificity of this enzyme (*CviJI***), yield a quasi-random distribution of DNA fragments from the small molecule pUC19 (2688 base pairs). Fitzgerald *et al.* (1992) quantitatively evaluated the randomness of this fragmentation strategy, using a *CviJI*** digest of pUC19 that was size fractionated by a rapid gel filtration method and directly ligated, without end repair, to a lac Z minus M13 cloning vector. Sequence analysis of 76 clones showed that *CviJI*** restricts pyGCPy and PuGCPu, in addition to PuGCPy sites, and that new sequence data is accumulated at a rate consistent with random fragmentation.

As reported in the literature, advantages of this approach compared to sonication and agarose gel fractionation include: smaller amounts of DNA are required (0.2-0.5 ug instead of 2-5 ug); and fewer steps are involved (no preligation, end repair, chemical extraction, or agarose gel electrophoresis and elution are needed).

Irrespective of the manner in which the nucleic acid fragments are obtained or prepared, it is important to denature the DNA to give single stranded pieces available for hybridization. This is achieved by incubating the DNA solution for 2-5 minutes at 80-90°C. The solution is then cooled quickly to 2°C to prevent renaturation of the DNA fragments before they are contacted with the chip. Phosphate groups must also be removed from genomic DNA by methods known in the art.

4.19 PREPARATION OF DNA ARRAYS

Arrays may be prepared by spotting DNA samples on a support such as a nylon membrane. Spotting may be performed by using arrays of metal pins (the positions of which correspond to an array of wells in a microtiter plate) to repeated by transfer of about 20 nl of a DNA solution to a nylon membrane. By offset printing, a density of dots higher than the

density of the wells is achieved. One to 25 dots may be accommodated in 1 mm², depending on the type of label used. By avoiding spotting in some preselected number of rows and columns, separate subsets (subarrays) may be formed. Samples in one subarray may be the same genomic segment of DNA (or the same gene) from different individuals, or
5 may be different, overlapped genomic clones. Each of the subarrays may represent replica spotting of the same samples. In one example, a selected gene segment may be amplified from 64 patients. For each patient, the amplified gene segment may be in one 96-well plate (all 96 wells containing the same sample). A plate for each of the 64 patients is prepared. By using a 96-pin device, all samples may be spotted on one 8 x 12 cm membrane. Subarrays
10 may contain 64 samples, one from each patient. Where the 96 subarrays are identical, the dot span may be 1 mm² and there may be a 1 mm space between subarrays.

Another approach is to use membranes or plates (available from NUNC, Naperville, Illinois) which may be partitioned by physical spacers e.g. a plastic grid molded over the membrane, the grid being similar to the sort of membrane applied to the bottom of multiwell
15 plates, or hydrophobic strips. A fixed physical spacer is not preferred for imaging by exposure to flat phosphor-storage screens or x-ray films.

The present invention is illustrated in the following examples. Upon consideration of the present disclosure, one of skill in the art will appreciate that many other embodiments and variations may be made in the scope of the present invention. Accordingly, it is
20 intended that the broader aspects of the present invention not be limited to the disclosure of the following examples. The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and compositions and methods which are functionally equivalent are within the scope of the invention. Indeed, numerous modifications and variations in the practice of the
25 invention are expected to occur to those skilled in the art upon consideration of the present preferred embodiments. Consequently, the only limitations which should be placed upon the scope of the invention are those which appear in the appended claims.

All references cited within the body of the instant specification are hereby incorporated by reference in their entirety.

5.0 EXAMPLES

5.1 EXAMPLE 1

Novel Nucleic Acid Sequences Obtained From Various Libraries

A plurality of novel nucleic acids were obtained from cDNA libraries prepared from various human tissues and in some cases isolated from a genomic library derived from human chromosome using standard PCR, SBH sequence signature analysis and Sanger sequencing techniques. The inserts of the library were amplified with PCR using primers specific for the vector sequences which flank the inserts. Clones from cDNA libraries were spotted on nylon membrane filters and screened with oligonucleotide probes (e.g., 7-mers) to obtain signature sequences. The clones were clustered into groups of similar or identical sequences. Representative clones were selected for sequencing.

In some cases, the 5' sequence of the amplified inserts was then deduced using a typical Sanger sequencing protocol. PCR products were purified and subjected to fluorescent dye terminator cycle sequencing. Single pass gel sequencing was done using a 377 Applied Biosystems (ABI) sequencer to obtain the novel nucleic acid sequences. In some cases RACE (Random Amplification of cDNA Ends) was performed to further extend the sequence in the 5' direction.

5.2 EXAMPLE 2

Novel Nucleic Acids

The novel nucleic acids of the present invention of the invention were assembled from sequences that were obtained from a cDNA library by methods described in Example 1 above, and in some cases sequences obtained from one or more public databases. The nucleic acids were assembled using an EST sequence as a seed. Then a recursive algorithm was used to extend the seed EST into an extended assemblage, by pulling additional sequences from different databases (i.e., Hyseq's database containing EST sequences, dbEST version 114, gb pri 114, and UniGene version 101) that belong to this assemblage. The algorithm terminated when there was no additional sequences from the above databases that would extend the assemblage. Inclusion of component sequences into the assemblage was based on a BLASTN hit to the extending assemblage with BLAST score greater than 300 and percent identity greater than 95%.

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e., dbEST version 118, gb pri 118, UniGene version 118, Genepet release 118). Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide and amino acid sequences, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS: 1- 35.

Table 1 shows the various tissue sources of SEQ ID NO: 1-35.

The homology for SEQ ID NO: 1-35 were obtained by a BLASTP version 2.0a1 19MP-WashU search against Genpept release 118, using BLAST algorithm. The results showed homologues for SEQ ID NO: 1-35 from Genpept. The homologues with identifiable functions for SEQ ID NO: 1-35 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication " Identification of prokaryotic and

TABLE 1

TISSUE ORIGIN	RNA SOURCE	HYSEQ LIBRARY NAME	SEQ ID NOS:
adult brain	GIBCO	AB3001	1 5 11 19 23-24 35
adult brain	GIBCO	ABD003	3 8 11 15 18 20 23-24 26 34-35
adult brain	Clontech	ABR001	5-7 11 19
adult brain	Clontech	ABR006	6 10 12 15 18 20 27
adult brain	Clontech	ABR008	1-4 6-11 14 16 18-19 21-22 26-28 30-31 35
adult brain	BioChain	ABR012	10 35
adult brain	Invitrogen	ABR014	10
adult brain	Invitrogen	ABT004	2 6 16 21-22 24
cultured preadipocytes	Stratagene	ADP001	10 24 26 31 33
adrenal gland	Clontech	ADR002	5 7 11 24 28
adult heart	GIBCO	AHR001	1 3 5-8 10-11 14-16 25 33 35
adult kidney	GIBCO	AKD001	3 5 8 11 15-16 21-22 24 28 30 33
adult kidney	Invitrogen	AKT002	8 10 12 15 18 21-22 26 28
adult lung	GIBCO	ALG001	10 33 35
young liver	GIBCO	ALV001	2 14 17 21-22 24 26
adult liver	Invitrogen	ALV002	17-18 24 28
adult liver	Clontech	ALV003	17
adult ovary	Invitrogen	AOV001	1-2 5-6 8 10-11 14 16 18-22 24 26 28 30 33 35
adult placenta	Clontech	APL001	6
placenta	Invitrogen	APL002	10 24 35
adult spleen	GIBCO	ASP001	2 7 10-11 18 24-25 28 33
testis	GIBCO	ATS001	21-22 26 28 30
adult bladder	Invitrogen	BLD001	10
bone marrow	Clontech	BMD001	1 11 21-22 31 33 35
bone marrow	Clontech	BMD002	1 10-11 14 25-26 28
adult colon	Invitrogen	CLN001	5 11 16
Mixture of 16 tissues - mRNAs*	Various Vendors*	CTL016	1
Mixture of 16 tissues - mRNAs*	Various Vendors*	CTL021	17
Mixture of 16 tissues - mRNAs*	Various Vendors*	CTL028	10
adult cervix	BioChain	CVX001	1-2 10-11 14 18 28 33 35
endothelial cells	Stratagene	EDT001	6 8 10-11 21-22 24 26 33 35
fetal brain	Clontech	FBR004	2 20 26 30
fetal brain	Clontech	FBR006	2 6 8-9 11 16 18 21-22 27 30

* The 16 tissue-mRNAs and their vendor source, are as follows: 1) Normal adult brain mRNA (Invitrogen), 2) normal adult kidney mRNA (Invitrogen), 3) normal adult liver mRNA (Invitrogen), 4) normal fetal brain mRNA (Invitrogen), 5) normal fetal kidney mRNA (Invitrogen), 6) normal fetal liver mRNA (Invitrogen), 7) normal fetal skin mRNA (Invitrogen), 8) human adrenal gland mRNA (Clontech), 9) human bone marrow mRNA (Clontech), 10) human leukemia lymphoblastic mRNA (Clontech), 11) human thymus mRNA (Clontech), 12) human lymph node mRNA (Clontech), 13) human spinal cord mRNA (Clontech), 14) human thyroid mRNA (Clontech), 15) human esophagus mRNA (BioChain), 16) human conceptional umbilical cord mRNA (BioChain).

TABLE 1

TISSUE ORIGIN	RNA SOURCE	HYSEQ LIBRARY NAME	SEQ ID NOS:
			35
fetal brain	Clontech	FBRs03	21-22
fetal brain	Invitrogen	FBT002	2 8-10 18-19 24 26 30
fetal kidney	Clontech	FKD001	7 28
fetal lung	Clontech	FLG001	10
fetal lung	Invitrogen	FLG003	10 24 26
fetal liver-spleen	Columbia University	FLS001	2 6-8 10-14 17 21-22 24 26 28 31 34-35
fetal liver-spleen	Columbia University	FLS002	3 5-6 8 11 13 16 18 21-22 24-26 28 30 32-35
fetal liver-spleen	Columbia University	FLS003	10 17-18 26
fetal liver	Invitrogen	FLV001	14 17 24 33 35
fetal liver	Clontech	FLV004	10 14 28
fetal muscle	Invitrogen	FMS001	3 6 10 26 31
fetal muscle	Invitrogen	FMS002	10 25 30
fetal skin	Invitrogen	FSK001	3 5-6 10 16-17 21-23 25-26 28 30-31
fetal skin	Invitrogen	FSK002	10 18 26
fetal spleen	BioChain	FSP001	10
umbilical cord	BioChain	FUC001	3 10 20 24 26 30 33
fetal brain	GIBCO	HFB001	3-6 8 10-11 15 18 21-24 35
infant brain	Columbia University	IB2002	2 4 7-8 10 14 16 18-19 21-23 26 28 35
infant brain	Columbia University	IB2003	3 5 8-10 16 20 23 26 29 35
infant brain	Columbia University	IBM002	19
infant brain	Columbia University	IBS001	26 35
lung, fibroblast	Stratagene	LFB001	10 21-22 30 33
lung tumor	Invitrogen	LGT002	2 10-11 14-15 24 28 30 35
lymphocytes	ATCC	LPC001	1 3 11 18 28 35
leukocyte	GIBCO	LUC001	1-3 5-7 11 16 18 21-22 24-26 28 35
melanoma from cell line ATCC #CRL 1424	Clontech	MEL004	2 24 26
mammary gland	Invitrogen	MMG001	2 6 10 12 14 16 18-19 24 26 28 31 35
induced neuron cells	Stratagene	NTD001	15 23
neuronal cells	Stratagene	NTU001	10 26
placenta	Clontech	PLA003	33
prostate	Clontech	PRT001	7-8 10-11 14 16 21-22 24
rectum	Invitrogen	REC001	6 26 28
salivary gland	Clontech	SAL001	10 16 21-22 35
skin fibroblast	ATCC	SFB001	10
small intestine	Clontech	SIN001	3 5 10 14 24 30
skeletal muscle	Clontech	SKM001	10 15
spinal cord	Clontech	SPC001	18 26 28
adult spleen	Clontech	SPLc01	1 14 26
stomach	Clontech	STO001	16
thalamus	Clontech	THA002	2-3 24 26

TABLE 1

TISSUE ORIGIN	RNA SOURCE	HYSEQ LIBRARY NAME	SEQ ID NOS:
thymus	Clontech	THMc02	3 6-7 14 17 26 31-32 35
thyroid gland	Clontech	THR001	3 6 8 10-11 17-18 21-22 24 26 28 33
trachea	Clontech	TRC001	10 21-22 33
uterus	Clontech	UTR001	10-11 24

TABLE 2

SEQ ID NO:	CORRESPONDING SEQ ID NO. IN U.S.S.N 09/574,454	ACCESSION NUMBER	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
1	1621	AL049785	Homo sapiens hypothetical protein	761	100
2	1724	U20657	Homo sapiens ubiquitin protease	719	47
3	2106	Y14494	Homo sapiens aralar1	3462	99
4	2500	U00051	Caenorhabditis elegans coded for by C. elegans cDNA yk50b2.5; coded for by C. elegans cDNA CEESV26F; similar to lipases over a short region	414	33
5	2501	X62575	Drosophila melanogaster ubiquitin-conjugating enzyme	78	26
6	2507	AB017005	Homo sapiens PMS2L14	1552	99
7	2520	AF222766	Bos taurus ankyrin 1	835	31
8	2555	AB039670	Homo sapiens ALEX1	2332	100
9	3766	U51000	Mus musculus DLX-1	1319	98
10	3935	X66405	Mus musculus collagen alpha1 type VI-precursor	5011	89
11	4297	AF208856	Homo sapiens BM-014	1012	99
12	4333	X62677	Oryctolagus cuniculus retrovirus related reverse transcriptase	75	52
13	4449	M14912	Homo sapiens pol	132	86
14	4562	AB024028	Arabidopsis thaliana uridine kinase-like protein	1021	45
15	4591	L11275	Saccharomyces cerevisiae selected as a weak suppressor of a mutant of the subunit AC40 of DNA dependant RNA polymerase I and III	97	19
16	4614	AL110500	Caenorhabditis elegans Y87G2A.13	260	24
17	4711	A14829	Homo sapiens preproapolipoprotein	974	100

TABLE 2

SEQ ID NO:	CORRESPONDING SEQ ID NO. IN U.S.S.N 09/574,454	ACCESSION NUMBER	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
18	4726	AF151086	Homo sapiens HSPC252	1223	81
19	4737	AF083116	Homo sapiens paraneoplastic cancer-testis-brain antigen	122	42
20	4762	AC006963	Homo sapiens similar to Kelch proteins; similar to BAA77027 (PID:g4650844)	1226	43
21	4790	L39995	Ceratovacuna lanigera cytochrome oxidase I	76	28
22	4790	L39995	Ceratovacuna lanigera cytochrome oxidase I	76	28
23	4828	AB035266	Homo sapiens neurexin II	9034	100
24	4999	M37194	Rattus norvegicus clathrin-associated protein 17	704	91
25	5155	AF217516	Homo sapiens uncharacterized bone marrow protein BM040	669	100
26	5244	U35776	Rattus norvegicus ADP-ribosylation factor 1-directed GTPase activating protein	1849	84
27	5280	AF159852	Drosophila melanogaster RNA-binding protein Smaug	259	44
28	5502	Y13247	Homo sapiens FB19 protein	5157	99
29	5813	AF116638	Homo sapiens PRO1546	70	48
30	5868	L21013	Dictyostelium discoideum RabC	126	25
31	5890	AL031033	Homo sapiens C321D2.4 (novel protein)	960	100
32	6070	X58236	Homo sapiens 36/8-8 fusion protein with epitope for anti-lectin antibody	90	73
33	6245	AF095593	Homo sapiens caveolin-1	946	100
34	6277	AF133521	Libellula pulchella troponin T	45	43

TABLE 2

SEQ ID NO:	CORRESPONDING SEQ ID NO. IN U.S.S.N 09/574,454	ACCESSION NUMBER	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
35	6298	AF161491	Homo sapiens HSPC142	1379	100

TABLE 3

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
1	DM01117	2 kw TRANSPOSASE WITHIN TRANSPOSITION VASOTOCIN.	DM01117A 11.17 9.173e-06 52-70
2	BL00972	Ubiquitin carboxyl-terminal hydrolases family 2 proteins.	BL00972A 11.93 5.091e-19 498-516 BL00972D 22.55 4.000e-15 1148-1173 BL00972C 16.48 9.143e-12 642-657 BL00972E 20.72 4.462e-11 1182-1204 BL00972B 9.45 4.176e-09 584-594
3	BL00215	Mitochondrial energy transfer proteins.	BL00215A 15.82 1.600e-15 333-358 BL00215A 15.82 1.794e-14 433-458 BL00215A 15.82 4.441e-14 525-550 BL00215B 10.44 3.250e-10 569-582 BL00215B 10.44 1.000e-08 381-394
4	PF00756	Putative esterase.	PF00756C 14.12 2.084e-09 206-236
5	PR00701	60KD INNER MEMBRANE PROTEIN SIGNATURE	PR00701I 8.59 6.256e-06 19-43
6	BL00058	DNA mismatch repair proteins mutL / hexB / PMS1 proteins.	BL00058A 20.73 5.125e-33 108-159 BL00058B 16.83 3.721e-29 172-211 BL00058C 18.05 7.568e-16 240-258 BL00058C 18.05 8.235e-13 36-54
7	PF00023	Ank repeat proteins.	PF00023A 16.03 7.600e-13 177-193 PF00023A 16.03 8.286e-11 144-160 PF00023A 16.03 2.500e-10 760-776 PF00023B 14.20 5.000e-10 339-349 PF00023B 14.20 5.500e-10 239-249 PF00023B 14.20 6.500e-10 471-481 PF00023B 14.20 8.000e-10 823-833 PF00023A 16.03 8.875e-10 554-570 PF00023A 16.03 1.000e-09 45-61 PF00023B 14.20 2.227e-09 438-448 PF00023B 14.20 2.636e-09 272-282 PF00023B 14.20 2.636e-09 583-593 PF00023A 16.03 5.821e-09 587-603 PF00023B 14.20 7.955e-09 405-415 PF00023A 16.03 9.679e-09 210-226
8	DM00892	3 RETROVIRAL PROTEINASE.	DM00892B 9.78 4.767e-06 27-33
9	PR00031	LAMBDA AND OTHER REPRESSOR HELIX-TURN-HELIX SIGNATURE	PR00031B 16.29 1.643e-17 166-183 PR00031A 8.77 9.400e-09 157-167
10	PR00159	2FE-2S FERREDOXIN SIGNATURE	PR00159B 8.50 7.882e-09 604-612
11	PR00138	MATRIXIN SIGNATURE	PR00138D 16.56 4.517e-09 77-103
12	BL00597	Plant lipid transfer proteins.	BL00597B 12.41 7.955e-06 11-40
13	BL00366	Uricase proteins.	BL00366A 11.55 9.780e-06 25-39
15	PF00602	Influenza RNA-dependant RNA polymerase subunit PB1.	PF00602E 11.62 5.160e-06 95-142
16	DM01688	2 POLY-IG RECEPTOR.	DM01688K 17.19 5.938e-06 321-360

* Results include in order: accession number subtype; raw score; p-value; position of signature in amino acid sequence.

TABLE 3

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
17	PD02807	APOLIPOPROTEIN E PRECURSOR APO-E GLYCOPROTEIN PLAS.	PD02807D 7.99 5.534e-09 105-155
18	DM00303	6 LEA 11-MER REPEAT REPEAT.	DM00303B 21.87 9.173e-07 217-252
19	BL00048	Protamine P1 proteins.	BL00048 6.39 8.763e-09 106-133 BL00048 6.39 9.663e-09 108-135
20	PF00651	BTB (also known as BR-C/Ttk) domain proteins.	PF00651 15.00 9.182e-15 76-89
21	PR00701	60KD INNER MEMBRANE PROTEIN SIGNATURE	PR00701B 15.26 7.188e-07 177-199
22	PR00701	60KD INNER MEMBRANE PROTEIN SIGNATURE	PR00701B 15.26 7.188e-07 177-199
23	DM00060	338 kw NEUREXIN ALPHA III CYSTEINE.	DM00060 6.92 3.925e-10 210-220
24	PR00317	EPENDYMIN SIGNATURE	PR00317F 10.90 5.935e-09 117-132
25	PR00305	14-3-3 PROTEIN ZETA SIGNATURE	PR00305F 15.95 7.150e-06 226-256
26	PR00405	HIV REV INTERACTING PROTEIN SIGNATURE	PR00405A 17.71 2.286e-20 19-39 PR00405B 11.83 6.077e-15 38-56 PR00405C 19.41 4.000e-13 59-81
28	PR00334	HMW KININOGEN SIGNATURE	PR00334B 8.69 9.914e-09 867-891
29	PR00551	2-S GLOBULIN FAMILY SIGNATURE	PR00551E 10.27 6.211e-06 12-27
30	PR00300	ATP-DEPENDENT CLP PROTEASE ATP-BINDING SUBUNIT SIGNATURE	PR00300A 9.56 7.896e-09 6-25
31	PR00467	MAMMALIAN LIPOXYGENASE SIGNATURE	PR00467E 9.00 1.000e-05 27-47
33	BL01210	Caveolins proteins.	BL01210B 13.92 1.000e-40 91-141 BL01210A 17.61 6.000e-39 43-82 BL01210C 17.86 8.579e-34 141-173
34	PR00658	CD44 ANTIGEN PRECURSOR SIGNATURE	PR00658D 8.52 9.919e-06 12-32
35	PR00512	5-HYDROXYTRYPTAMINE 1A RECEPTOR SIGNATURE	PR00512E 10.80 4.884e-06 30-47

TABLE 4

SEQ ID NO:	pFAM NAME	DESCRIPTION	p-value	pFAM SCORE
2	zf-MYND	MYND finger	1.1e-06	35.6
3	efhand	EF hand	0.0016	25.1
6	DNA_mis_repair	DNA mismatch repair protein	8.2e-36	132.4
7	ank	Ank repeat	3.6e-197	668.4
9	homeobox	Homeobox domain	1.5e-29	111.6
10	Collagen	Collagen triple helix repeat (20 copies)	7.5e-46	165.8
14	PRK	Phosphoribulokinase	9.5e-06	-24.1
17	Apolipoprotein	Apolipoprotein A1/A4/E family	5.2e-113	388.8
18	BTB	BTB/POZ domain	2.6e-25	97.5
20	Kelch	Kelch motif	1.7e-22	88.2
23	EGF	EGF-like domain	3.5e-09	43.9
24	Clat_adaptor_s	Clathrin adaptor complex small chain	5.7e-88	305.7
26	ArfGap	Putative GTP-ase activating protein for Arf	6.3e-62	219.2
27	SAM	SAM domain (Sterile alpha motif)	0.003	20.7
30	ras	Ras family	0.0058	-103.2
33	Caveolin	Caveolin	3.7e-120	412.6

TABLE 5

SEQ ID NO:	POSITION OF SIGNAL IN AMINO ACID SEQUENCE	maxS (MAXIMUM SCORE)	meanS (MEAN SCORE)
10	1-19	0.968	0.899
12	1-23	0.960	0.861
16	1-28	0.983	0.763
17	1-18	0.966	0.921
23	1-28	0.963	0.881
29	1-20	0.940	0.677

CLAIMS

WHAT IS CLAIMED IS:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 1 – 35, a mature protein coding portion of SEQ ID NO: 1 – 35, an active domain of SEQ ID NO: 1 – 35, and complementary sequences thereof.
2. An isolated polynucleotide encoding a polypeptide with biological activity, wherein said polynucleotide hybridizes to the polynucleotide of claim 1 under stringent hybridization conditions.
3. An isolated polynucleotide encoding a polypeptide with biological activity, wherein said polynucleotide has greater than about 90% sequence identity with the polynucleotide of claim 1.
4. The polynucleotide of claim 1 wherein said polynucleotide is DNA.
5. An isolated polynucleotide of claim 1 wherein said polynucleotide comprises the complementary sequences.
6. A vector comprising the polynucleotide of claim 1.
7. An expression vector comprising the polynucleotide of claim 1.
8. A host cell genetically engineered to comprise the polynucleotide of claim 1.
9. A host cell genetically engineered to comprise the polynucleotide of claim 1 operatively associated with a regulatory sequence that modulates expression of the polynucleotide in the host cell.
10. An isolated polypeptide, wherein the polypeptide is selected from the group consisting of:
 - (a) a polypeptide encoded by any one of the polynucleotides of claim 1; and

(b) a polypeptide encoded by a polynucleotide hybridizing under stringent conditions with any one of SEQ ID NO: 1 – 35.

11. A composition comprising the polypeptide of claim 10 and a carrier.

12. An antibody directed against the polypeptide of claim 10.

13. A method for detecting the polynucleotide of claim 1 in a sample, comprising:

a) contacting the sample with a compound that binds to and forms a complex with the polynucleotide of claim 1 for a period sufficient to form the complex; and

b) detecting the complex, so that if a complex is detected, the polynucleotide of claim 1 is detected.

14. A method for detecting the polynucleotide of claim 1 in a sample, comprising:

a) contacting the sample under stringent hybridization conditions with nucleic acid primers that anneal to the polynucleotide of claim 1 under such conditions;

b) amplifying a product comprising at least a portion of the polynucleotide of claim 1; and

c) detecting said product and thereby the polynucleotide of claim 1 in the sample.

15. The method of claim 14, wherein the polynucleotide is an RNA molecule and the method further comprises reverse transcribing an annealed RNA molecule into a cDNA polynucleotide.

16. A method for detecting the polypeptide of claim 10 in a sample, comprising:

a) contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex; and

b) detecting formation of the complex, so that if a complex formation is detected, the polypeptide of claim 10 is detected.

17. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

a) contacting the compound with the polypeptide of claim 10 under conditions sufficient to form a polypeptide/compound complex; and

b) detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

18. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

a) contacting the compound with the polypeptide of claim 10, in a cell, under conditions sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a reporter gene sequence in the cell; and

b) detecting the complex by detecting reporter gene sequence expression, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

19. A method of producing the polypeptide of claim 10, comprising,

a) culturing a host cell comprising a polynucleotide sequence selected from the group consisting of a polynucleotide sequence of SEQ ID NO: 1-35, a mature protein coding portion of SEQ ID NO: 1-35, an active domain of SEQ ID NO: 1-35, complementary sequences thereof and a polynucleotide sequence hybridizing under stringent conditions to SEQ ID NO: 1-35, under conditions sufficient to express the polypeptide in said cell; and

b) isolating the polypeptide from the cell culture or cells of step (a).

20. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of any one of the polypeptides from the Sequence Listing, the mature protein portion thereof, or the active domain thereof.

21. The polypeptide of claim 20 wherein the polypeptide is provided on a polypeptide array.

22. A collection of polynucleotides, wherein the collection comprising the sequence information of at least one of SEQ ID NO: 1 – 35.

5 23. The collection of claim 22, wherein the collection is provided on a nucleic acid array.

24. The collection of claim 23, wherein the array detects full-matches to any one of the polynucleotides in the collection.

10 25. The collection of claim 23, wherein the array detects mismatches to any one of the polynucleotides in the collection.

15 26. The collection of claim 22, wherein the collection is provided in a computer-readable format.

27. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising a polypeptide of claim 10 or 20 and a pharmaceutically acceptable carrier.

20 28. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising an antibody that specifically binds to a polypeptide of claim 10 or 20 and a pharmaceutically acceptable carrier.

ABSTRACT OF THE INVENTION

The present invention provides novel nucleic acids, novel polypeptide sequences encoded by these nucleic acids and uses thereof.

5

789CIP2C

DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As [a] below named inventor(s), I/we hereby declare that:

**Y. Tom Tang, Chenghua Liu, Ping Zhou, Vinod Asundi, Feiyan Ren, Qing A. Zhao,
Jie Zhang, Jian-Rui Wang, Tom Wehrman, Radoje T. Drmanac**

My/our residence, post office address and citizenship is/are as stated below next to my/our name(s).

I/we believe I/we am/are an/the original, first and sole/joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled: NOVEL NUCLEIC ACIDS AND POLYPEPTIDES, the specification of which

 X is attached hereto.

 was filed on [date] as Application Serial Number []
and was amended on [date].

I/We hereby state that I/we have reviewed and understand the contents of the above-identified specification, including the claims as amended by any amendment referred to above.

I/We acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I/We hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate, listed below and so identified, and I/we have also identified below any foreign application for patent or inventor's certificate on this invention filed by me or my legal representatives or assigns and having a filing date before that of the application on which priority is claimed:

NUMBER	COUNTRY	DAY/MONTH/ YEAR FILED	PRIORITY CLAIMED - YES OR NO

I/We hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I/we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

SERIAL NUMBER	FILING DATE	STATUS
09/574,454	May 19, 2000	Pending
09/519,705	March 07, 2000	Pending

I/We hereby declare that all statements made herein of my/our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under § 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

I/We hereby appoint the following attorneys and agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the United States Patent and Trademark Office connected therewith and request that all correspondence and telephone calls with respect to this application be directed to Leslie A. Mooi, HYSEQ, INC., 670 Almanor Avenue, Sunnyvale, CA 94085, Telephone No. (408) 524-8100:

ATTORNEY	REGISTRATION NO.
Petrina S. Hsi	38,496
Leslie A. Mooi	37,047

Full name of
first joint inventor: Y. Tom Tang

Inventor's signature: _____ Date: _____

Residence and Post
Office Address: 4230 Ranwick Court, San Jose, CA 95118

Citizenship: United States of America

Full name of
second joint inventor: Chenghua Liu

Inventor's signature: _____ Date: _____

Residence and Post
Office Address: 1125 Ranchero Way, Apt. # 14, San Jose, CA 95117

Citizenship: People's Republic of China

Full name of
third joint inventor:

Ping Zhou

Inventor's signature:

Date: _____

Residence and Post
Office Address:

1461 Japaur Lane, San Jose, CA 95132

Citizenship:

People's Republic of China

Full name of
fourth joint inventor:

Vinod Asundi

Inventor's signature:

Date: _____

Residence and Post
Office Address:

709 Foster City Blvd., Foster City, CA 94404

Citizenship:

United States of America

Full name of
fifth joint inventor:

Feiyan Ren

Inventor's signature:

Date: _____

Residence and Post
Office Address:

20685 Garden Manor Court, Cupertino, CA 95014

Citizenship:

People's Republic of China

Full name of
sixth joint inventor:

Qing A. Zhao

Inventor's signature:

Date: _____

Residence and Post
Office Address:

1028 S. de Anza Blvd., Apt. B-210, San Jose, CA 95129

Citizenship:

People's Republic of China

Full name of
seventh joint inventor: Jie Zhang

Inventor's signature: _____ Date: _____

Residence and Post
Office Address: 20800 Homestead Road, #38B, Cupertino, CA 95014

Citizenship: People's Republic of China

Full name of
eighth joint inventor: Jian-Rui Wang

Inventor's signature: _____ Date: _____

Residence and Post
Office Address: 744 Stendhal Lane, Cupertino, CA 95014

Citizenship: People's Republic of China

Full name of
ninth joint inventor: Tom Wehrman

Inventor's signature: _____ Date: _____

Residence and Post
Office Address: 300 Pasteur Drive, Edwards R314, Stanford University
Medical Center, Stanford, CA 94305

Citizenship: United States of America

Full name of
tenth joint inventor: Radoje T. Drmanac

Inventor's signature: _____ Date: _____

Residence and Post
Office Address: 850 East Greenwich Place, Palo Alto, CA 94303

Citizenship: Yugoslavia

SEQUENCE LISTING

<110> Tang, Y. Tom
 Liu, Chenghua
 Zhou, Ping
 Asundi, Vinod
 Ren, Feiyan
 Zhao, Qing A.
 Zhang, Jie
 Wang, Jian-Rui
 Wehrman, Tom
 Drmanac, Radoje T.

<120> Novel Nucleic Acids and
 Polypeptides

<130> 789CIP2C

<140> To Be Assigned
 <141>

<150> 09/574,454
 <151> 2000-05-19

<150> 09/519,705
 <151> 2000-03-07

<160> 35

<170> pt_FL_genes Version 2.0

<210> 1
 <211> 2235
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (258)..(671)

<400> 1	
taagcttgcg gccgccccgg ccgccccgc gggggacacc tctcgccgc cacagcttta	60
ggaaacacct ctacctctg cgaggcctcc cgggctccgg gaaaactaca ctggccagac	120
aattgcagca tgactttccc agggccctga ttttcagcac ggatgatttt ttcttcaggg	180

aagatggtgc ctatgagttc aatcctgact tcctggagga agctcatgaa tggaacaaaa 240
aaagagcaag aaaagca atg agg aat ggc ata tcc ccc att att att gat 290
Met Arg Asn Gly Ile Ser Pro Ile Ile Ile Asp
1 5 10
aat acc aac ctc cac gcc tgg gaa atg aag ccc tat gca gtc atg gca 338
Asn Thr Asn Leu His Ala Trp Glu Met Lys Pro Tyr Ala Val Met Ala
15 20 25
ctt gaa aat aac tat gaa gtt ata ttc cga gaa cct gac act cgc tgg 386
Leu Glu Asn Asn Tyr Glu Val Ile Phe Arg Glu Pro Asp Thr Arg Trp
30 35 40
aaa ttc aac gtt caa gag tta gca aga aga aac att cat ggt gtc tca 434
Lys Phe Asn Val Gln Glu Leu Ala Arg Arg Asn Ile His Gly Val Ser
45 50 55
aga gaa aaa atc cac cga atg aaa gaa cgg tat gaa cac gat gtt act 482
Arg Glu Lys Ile His Arg Met Lys Glu Arg Tyr Glu His Asp Val Thr
60 65 70 75
ttt cac agt gtg ctt cat gca gaa aag cca agc aga atg aac aga aac 530
Phe His Ser Val Leu His Ala Glu Lys Pro Ser Arg Met Asn Arg Asn
80 85 90
cag gac agg aat aat gca ttg cct tcc aac aat gcc aga tac tgg aat 578
Gln Asp Arg Asn Asn Ala Leu Pro Ser Asn Asn Ala Arg Tyr Trp Asn
95 100 105
tcc tac aca gag ttt cca aac cgg agg gcc cac ggt gga ttt aca aat 626
Ser Tyr Thr Glu Phe Pro Asn Arg Arg Ala His Gly Gly Phe Thr Asn
110 115 120
gag agc tcc tat cac aga agg ggc ggt tgt cac cat gga tat tag agg 674
Glu Ser Ser Tyr His Arg Arg Gly Gly Cys His His Gly Tyr *
125 130 135
cctatcttac agccaggcag aattttccta agtcagtttc tacttcagtt tttgttat 734
tttgttgcat tttagtcaga gctccaattc cagtgtaaat agctgaactc aaaagtttct 794
gagcaaagtc attatattca ctttcttcac caaaatttgt taaagtgcct ctatatgcat 854
ggtctgatgc tgggaattct gcagatttga gtaaacagtc tctttctcta gggtagaat 914
ttgaaaccaa aacttgagaa cacaccaag aatatattta cataggttca tagatgaaat 974
aaagtgttta tattatatat aagcttcagt accatttgct ctgaagtgat ctatttat 1034
tttcaggaaa ttcattctcca tcggtaaagt tgggaagggtg gagagaagtg gtgggggggc 1094
attgctactt atcaaagtgc cattgctact ttgataatct atgtatctaa aaatgtgaga 1154
tgtgcgactc ttatgatact gattttcctt taatgttaat atgccagaaa gcatacatct 1214

aagggaacat	tgtccttcaa	agtagacact	ttgggaagtt	atctctttat	tttaatgatg	1274
tatcattggt	aaaaatgctg	tcaaatcctt	aatagctaca	ggagctactg	agggaaatca	1334
gtgtcattat	ttaaagtcac	gccttgtgtt	tttactactt	tattcagcag	gattaaacct	1394
gaataacttt	tggctgttgt	gctaatagtg	taaataaaat	aagcctgcct	tcataaaaca	1454
ctaactttta	aaaggaataa	acgacttcta	aaattatgcc	tattaacatg	tgtaattagt	1514
cggcagctca	aatgtttggg	agtgcaagaa	attaggcacc	ccaggatata	ggtcatacag	1574
ggatatataa	aagccatgct	cattacaaaa	tgagcagttg	atgttttatg	tggcattaag	1634
acaatcaagt	cctcacaact	ctggaatgtc	ttcttatact	gatgctgaat	ttatgaatcc	1694
aaattaattt	ccaacagggt	ggaatcagat	ttaatgtgag	atcatgatag	acaagaccac	1754
agaggacgta	tgctctattt	cttgttggcc	aacagcttct	ttctaattgt	ctgtgaaaaa	1814
ttattttaag	tgtcttatat	aatgggtgctt	ttatggttat	taaaaattgt	aatgggtatc	1874
acatttatat	ggatttgtca	ttggatcttt	ttttggttca	acaataaaaa	aatttaatta	1934
cctaaatgcc	aagaaactca	acaatatacc	agtttttctg	tatcacaggc	ttatttacca	1994
gtcttttttt	taataaatag	gaatcgtaaa	ggtaatgaca	aaagcagcct	tataatttag	2054
ttgcttatat	atttgatctg	tgtacatgag	actgttttaa	cgttatctga	cactactgaa	2114
acctgctcga	catctccatg	actaccaaca	ccatgtgtaa	tgttttcttc	actaacattt	2174
taaaaactgg	tatctccttt	gagtaagttt	ggctgacaat	agtaaattccc	aatgaatcta	2234
a						2235

<210> 2
 <211> 4600
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (349)..(4305)

ttcaaggtca	cgggcggtgc	cagtccttac	ccctagtgcc	ccgcagcgtg	ccagcccact	60
gatgccggca	ggccgttacg	ctatgcctgt	ggcagcctcc	cacctgggag	aggactggcg	120
gccacgaggg	caacttcgtg	ggagaggtgg	tggcgggctc	ggggttcaga	ggctgaagct	180
gggtcgcagc	tcgaccgggg	cgctgatctg	tggtgtccg	ttccttgctg	gagaatttgg	240

ccacaaagag ctgccaagat agctgggcca ggaagaaagc gccgcagccc tgacccagac 300

gctgttgccg accccggggc actctggctg tcgaccaagc ggctcaag atg tct ggc 357
Met Ser Gly
1

ggg gcc agt gcc aca ggc cca agg aga ggg ccc cca gga ctg gag gac 405
Gly Ala Ser Ala Thr Gly Pro Arg Arg Gly Pro Pro Gly Leu Glu Asp
5 10 15

acc act agt aag aag aag cag aag gat cga gca aac cag gag agc aag 453
Thr Thr Ser Lys Lys Lys Gln Lys Asp Arg Ala Asn Gln Glu Ser Lys
20 25 30 35

gat gga gat cct agg aaa gag aca ggg tct cga tat gtt gcc cag gct 501
Asp Gly Asp Pro Arg Lys Glu Thr Gly Ser Arg Tyr Val Ala Gln Ala
40 45 50

ggg ctt gaa cct ctg gcc tca ggt gat cct tct gcc tca gcc tcc cat 549
Gly Leu Glu Pro Leu Ala Ser Gly Asp Pro Ser Ala Ser Ala Ser His
55 60 65

gca gct ggg atc aca ggc tca cgc cac cgt acc cgg ctg ttc ttt cct 597
Ala Ala Gly Ile Thr Gly Ser Arg His Arg Thr Arg Leu Phe Phe Pro
70 75 80

tca tcg tca ggg tca gca tcc act cct caa gag gag cag acc aaa gag 645
Ser Ser Ser Gly Ser Ala Ser Thr Pro Gln Glu Glu Gln Thr Lys Glu
85 90 95

gga gct tgt gaa gac cct cat gat ctc ttg gct act ccc act cca gag 693
Gly Ala Cys Glu Asp Pro His Asp Leu Leu Ala Thr Pro Thr Pro Glu
100 105 110 115

ttg ttg ctc gat tgg agg cag agt gca gaa gag gtg att gtc aag ctt 741
Leu Leu Leu Asp Trp Arg Gln Ser Ala Glu Glu Val Ile Val Lys Leu
120 125 130

cgt gtg gga gta ggt ccc ctg cag ctg gag gat gta gat gct gct ttc 789
Arg Val Gly Val Gly Pro Leu Gln Leu Glu Asp Val Asp Ala Ala Phe
135 140 145

aca gat aca gac tgt gtg gtg cgg ttt gca ggt ggt cag cag tgg ggt 837
Thr Asp Thr Asp Cys Val Val Arg Phe Ala Gly Gly Gln Gln Trp Gly
150 155 160

ggg gtc ttc tat gct gag ata aaa agc tct tgt gct aaa gtg caa acc 885
Gly Val Phe Tyr Ala Glu Ile Lys Ser Ser Cys Ala Lys Val Gln Thr
165 170 175

cgc aag ggc agt ctc ctg cac ctg aca ctg ccc aaa aag gtg cct atg 933
Arg Lys Gly Ser Leu Leu His Leu Thr Leu Pro Lys Lys Val Pro Met
180 185 190 195

ctc acg tgg ccc tcc ctc ctg gtt gag gct gat gaa cag ctt tgc ata 981
Leu Thr Trp Pro Ser Leu Leu Val Glu Ala Asp Glu Gln Leu Cys Ile
200 205 210

cca ccg ctg aac tcc caa acc tgc ctc ctg ggc tca gag gag aat tta	1029
Pro Pro Leu Asn Ser Gln Thr Cys Leu Leu Gly Ser Glu Glu Asn Leu	
215 220 225	
gcc cct ttg gca gga gag aaa gca gtg cct ccc ggg aat gac cca gtc	1077
Ala Pro Leu Ala Gly Glu Lys Ala Val Pro Pro Gly Asn Asp Pro Val	
230 235 240	
tct cca gcc atg gtc cgg agc aga aac cct ggg aaa gat gac tgt gcc	1125
Ser Pro Ala Met Val Arg Ser Arg Asn Pro Gly Lys Asp Asp Cys Ala	
245 250 255	
aag gag gag atg gca gtg gca gca gat gct gca acc ttg gtg gat gag	1173
Lys Glu Glu Met Ala Val Ala Ala Asp Ala Ala Thr Leu Val Asp Glu	
260 265 270 275	
ccc gag tcg atg gtg aac ctg gcg ttt gtc aag aat gac tcg tat gag	1221
Pro Glu Ser Met Val Asn Leu Ala Phe Val Lys Asn Asp Ser Tyr Glu	
280 285 290	
aag ggc ccg gat tca gtg gtg gtg cac gtg tac gtg aag gag atc tgc	1269
Lys Gly Pro Asp Ser Val Val Val His Val Tyr Val Lys Glu Ile Cys	
295 300 305	
agg gac acc tca aga gta ctt ttc cgt gag cag gac ttc acg ctc atc	1317
Arg Asp Thr Ser Arg Val Leu Phe Arg Glu Gln Asp Phe Thr Leu Ile	
310 315 320	
ttc cag acc agg gat gga aac ttc ctg agg ctg cac ccg ggc tgt ggg	1365
Phe Gln Thr Arg Asp Gly Asn Phe Leu Arg Leu His Pro Gly Cys Gly	
325 330 335	
ccc cac acc acc ttc cgt tgg cag gtg aag ctc agg aat ctg att gag	1413
Pro His Thr Thr Phe Arg Trp Gln Val Lys Leu Arg Asn Leu Ile Glu	
340 345 350 355	
cca gag cag tgc acc ttc tgt ttc acg gct tct cgc atc gac atc tgc	1461
Pro Glu Gln Cys Thr Phe Cys Phe Thr Ala Ser Arg Ile Asp Ile Cys	
360 365 370	
ctt cgt aag agg cag agt cag cgc tgg ggg ggc ctg gag gcc ccg gct	1509
Leu Arg Lys Arg Gln Ser Gln Arg Trp Gly Gly Leu Glu Ala Pro Ala	
375 380 385	
gca cga gtg ggt ggt gca aag gtt gcc gtg ccg aca ggt cca acc cct	1557
Ala Arg Val Gly Gly Ala Lys Val Ala Val Pro Thr Gly Pro Thr Pro	
390 395 400	
ctg gat tca acc cca cca gga ggt gct ccc cac ccc ctg aca ggc cag	1605
Leu Asp Ser Thr Pro Pro Gly Gly Ala Pro His Pro Leu Thr Gly Gln	
405 410 415	
gag gag gcc cgg gct gtg gag aag gat aaa tcc aag gca cga tct gag	1653
Glu Glu Ala Arg Ala Val Glu Lys Asp Lys Ser Lys Ala Arg Ser Glu	
420 425 430 435	

gac aca ggg cta gac agt gtg gca acc cgc aca ccc atg gag cat gta	1701
Asp Thr Gly Leu Asp Ser Val Ala Thr Arg Thr Pro Met Glu His Val	
440 445 450	
acc cca aag cca gag aca cac ctg gcc tcg ccc aag cct aca tgc atg	1749
Thr Pro Lys Pro Glu Thr His Leu Ala Ser Pro Lys Pro Thr Cys Met	
455 460 465	
gtg cct ccc atg ccc cac agc cca gtt agt gga gac agc gtg gag gag	1797
Val Pro Pro Met Pro His Ser Pro Val Ser Gly Asp Ser Val Glu Glu	
470 475 480	
gag gaa gag gaa gag aag aag gtg tgt ctg cca ggc ttc act ggc ctt	1845
Glu Glu Glu Glu Glu Lys Lys Val Cys Leu Pro Gly Phe Thr Gly Leu	
485 490 495	
gtc aat tta ggc aac acc tgc ttc atg aac agc gtc att cag tct ctg	1893
Val Asn Leu Gly Asn Thr Cys Phe Met Asn Ser Val Ile Gln Ser Leu	
500 505 510 515	
tcc aac act cgg gaa ctc cgg gac ttc ttc cat gac cgc tcc ttt gag	1941
Ser Asn Thr Arg Glu Leu Arg Asp Phe Phe His Asp Arg Ser Phe Glu	
520 525 530	
gct gag atc aac tac aac aac cca cta ggg act ggt ggg cgt ctg gcc	1989
Ala Glu Ile Asn Tyr Asn Asn Pro Leu Gly Thr Gly Gly Arg Leu Ala	
535 540 545	
att ggc ttt gcc gtg ctg ctt cgg gcg ctg tgg aag ggc acc cac cat	2037
Ile Gly Phe Ala Val Leu Leu Arg Ala Leu Trp Lys Gly Thr His His	
550 555 560	
gcc ttc cag cct tcc aag ttg aag gcc att gtg gcg agt aag gcc agc	2085
Ala Phe Gln Pro Ser Lys Leu Lys Ala Ile Val Ala Ser Lys Ala Ser	
565 570 575	
cag ttc aca ggc tat gca cag cat gat gcc cag gag ttc atg gct ttc	2133
Gln Phe Thr Gly Tyr Ala Gln His Asp Ala Gln Glu Phe Met Ala Phe	
580 585 590 595	
ctg ctg gat ggg ctg cac gag gac ctg aat cgc att cag aac aag ccc	2181
Leu Leu Asp Gly Leu His Glu Asp Leu Asn Arg Ile Gln Asn Lys Pro	
600 605 610	
tac aca gag acc gtg gat tca gat ggg cgg ccc gat gag gtg gta gct	2229
Tyr Thr Glu Thr Val Asp Ser Asp Gly Arg Pro Asp Glu Val Val Ala	
615 620 625	
gag gaa gca tgg cag cgg cac aag atg agg aat gac tct ttc atc gtg	2277
Glu Glu Ala Trp Gln Arg His Lys Met Arg Asn Asp Ser Phe Ile Val	
630 635 640	
gac cta ttt cag ggg cag tac aag tcg aag ctg gtg tgc cct gtg tgt	2325
Asp Leu Phe Gln Gly Gln Tyr Lys Ser Lys Leu Val Cys Pro Val Cys	
645 650 655	
gcc aag gtc tcc atc act ttt gac ccg ttt ctt tat ctg ccg gtg ccc	2373

Ala Lys Val Ser Ile Thr Phe Asp Pro Phe Leu Tyr Leu Pro Val Pro	
660 665 670 675	
ttg cca caa aag caa aag gtt ctc cct gtc ttt tat ttt gcc cga gag	2421
Leu Pro Gln Lys Gln Lys Val Leu Pro Val Phe Tyr Phe Ala Arg Glu	
680 685 690	
ccc cac agc aag ccc atc aag ttc ctg gtg agc gtc agc aag gag aac	2469
Pro His Ser Lys Pro Ile Lys Phe Leu Val Ser Val Ser Lys Glu Asn	
695 700 705	
tcc act gcg agc gaa gta ttg gac tcc ctc tct cag agt gtt cat gtg	2517
Ser Thr Ala Ser Glu Val Leu Asp Ser Leu Ser Gln Ser Val His Val	
710 715 720	
aag cct gag aac ctg cgt ttg gcg gag gta att aag aat cgt ttt cat	2565
Lys Pro Glu Asn Leu Arg Leu Ala Glu Val Ile Lys Asn Arg Phe His	
725 730 735	
cgt gtg ttc cta ccc tcc cac tca ctg gac act gtg tcc cca tct gat	2613
Arg Val Phe Leu Pro Ser His Ser Leu Asp Thr Val Ser Pro Ser Asp	
740 745 750 755	
acg ctc ctc tgc ttt gag ctg cta tcc tca gag ttg gct aag gag cgg	2661
Thr Leu Leu Cys Phe Glu Leu Leu Ser Ser Glu Leu Ala Lys Glu Arg	
760 765 770	
gta gtg gtg cta gag gtg caa cag cgc ccc cag gtg ccc agc gtc ccc	2709
Val Val Val Leu Glu Val Gln Gln Arg Pro Gln Val Pro Ser Val Pro	
775 780 785	
atc tcc aag tgt gca gcc tgc cag cgg aag caa cag tcg gag gat gaa	2757
Ile Ser Lys Cys Ala Ala Cys Gln Arg Lys Gln Gln Ser Glu Asp Glu	
790 795 800	
aag ctg aag cgc tgt acc cgg tgc tac cgt gtg ggc tac tgc aac cag	2805
Lys Leu Lys Arg Cys Thr Arg Cys Tyr Arg Val Gly Tyr Cys Asn Gln	
805 810 815	
ctc tgc cag aaa acc cac tgg cct gac cac aag ggc ctc tgc cga cct	2853
Leu Cys Gln Lys Thr His Trp Pro Asp His Lys Gly Leu Cys Arg Pro	
820 825 830 835	
gag aac att ggc tac ccc ttc ctg gtc agt gta cct gcc tca cgc ctc	2901
Glu Asn Ile Gly Tyr Pro Phe Leu Val Ser Val Pro Ala Ser Arg Leu	
840 845 850	
act tat gcc cgc ctc gct cag ttg cta gag ggc tat gcc cgg tac tct	2949
Thr Tyr Ala Arg Leu Ala Gln Leu Leu Glu Gly Tyr Ala Arg Tyr Ser	
855 860 865	
gtg agt gta ttc cag cca ccc ttt cag cca ggc cgc atg gcc ttg gag	2997
Val Ser Val Phe Gln Pro Pro Phe Gln Pro Gly Arg Met Ala Leu Glu	
870 875 880	
tct cag agc cct ggc tgc acc aca ctg ctc tcc aca ggt tcc ctg gag	3045
Ser Gln Ser Pro Gly Cys Thr Thr Leu Leu Ser Thr Gly Ser Leu Glu	

885	890	895	
gct ggg gac agc gag aga gac ccc att cag cca cct gag ctc cag ctg			3093
Ala Gly Asp Ser Glu Arg Asp Pro Ile Gln Pro Pro Glu Leu Gln Leu			
900	905	910	915
gtg acc cct atg gct gag ggg gac aca ggg ctt ccc cgg gtg tgg gca			3141
Val Thr Pro Met Ala Glu Gly Asp Thr Gly Leu Pro Arg Val Trp Ala			
	920	925	930
gcc cct gac cgg ggt cct gtg ccc agc acc agt gga att tct tct gag			3189
Ala Pro Asp Arg Gly Pro Val Pro Ser Thr Ser Gly Ile Ser Ser Glu			
	935	940	945
atg ctg gcc agt ggg ccc att gag gtt ggc tcc ttg cca gct ggc gag			3237
Met Leu Ala Ser Gly Pro Ile Glu Val Gly Ser Leu Pro Ala Gly Glu			
	950	955	960
agg gtg tcc cga ccc gaa gct gct gtg cct ggg tac cag cat cca agt			3285
Arg Val Ser Arg Pro Glu Ala Ala Val Pro Gly Tyr Gln His Pro Ser			
	965	970	975
gaa gct atg aat gcc cac aca ccc cag ttc ttc atc tat aaa att gat			3333
Glu Ala Met Asn Ala His Thr Pro Gln Phe Phe Ile Tyr Lys Ile Asp			
	980	985	990
tca tcc aac cga gag cag cgg cta gag gac aaa gga gac acc cca ctg			3381
Ser Ser Asn Arg Glu Gln Arg Leu Glu Asp Lys Gly Asp Thr Pro Leu			
	1000	1005	1010
gag ctg ggt gac gac tgt agc ctg gct ctc gtc tgg cgg aac aat gag			3429
Glu Leu Gly Asp Asp Cys Ser Leu Ala Leu Val Trp Arg Asn Asn Glu			
	1015	1020	1025
cgc ttg cag gag ttt gtg ttg gta gcc tcc aag gag ctg gaa tgt gct			3477
Arg Leu Gln Glu Phe Val Leu Val Ala Ser Lys Glu Leu Glu Cys Ala			
	1030	1035	1040
gag gat cca ggc tct gcc ggt gag gct gcc cgg gcc ggc cac ttc acc			3525
Glu Asp Pro Gly Ser Ala Gly Glu Ala Ala Arg Ala Gly His Phe Thr			
	1045	1050	1055
ctg gac cag tgc ctc aac ctc ttc aca cgg cct gag gtg ctg gca ccc			3573
Leu Asp Gln Cys Leu Asn Leu Phe Thr Arg Pro Glu Val Leu Ala Pro			
1060	1065	1070	1075
gag gag gcc tgg tac tgc cca cag tgc aaa cag cac cgt gag gcc tcc			3621
Glu Glu Ala Trp Tyr Cys Pro Gln Cys Lys Gln His Arg Glu Ala Ser			
	1080	1085	1090
aag cag ctg ttg cta tgg cgc ctg cca aat gtt ctc atc gtg cag ctc			3669
Lys Gln Leu Leu Leu Trp Arg Leu Pro Asn Val Leu Ile Val Gln Leu			
	1095	1100	1105
aag cgc ttc tcc ttt cgt agt ttt atc tgg cgt gac aag atc aat gac			3717
Lys Arg Phe Ser Phe Arg Ser Phe Ile Trp Arg Asp Lys Ile Asn Asp			
	1110	1115	1120

ttg gtg gag ttc cct gtt agg aac ctg gac ctg agc aag ttc tgc att 3765
 Leu Val Glu Phe Pro Val Arg Asn Leu Asp Leu Ser Lys Phe Cys Ile
 1125 1130 1135

ggt cag aaa gag gag cag ctg ccc agc tac gat cta tat gct gtc atc 3813
 Gly Gln Lys Glu Glu Gln Leu Pro Ser Tyr Asp Leu Tyr Ala Val Ile
 1140 1145 1150 1155

aac cac tat gga ggc atg att ggt ggc cac tac act gcc tgt gca cgc 3861
 Asn His Tyr Gly Gly Met Ile Gly Gly His Tyr Thr Ala Cys Ala Arg
 1160 1165 1170

ctg ccc aat gat cgt agc agt cag cgc agt gac gtg ggc tgg cgc ttg 3909
 Leu Pro Asn Asp Arg Ser Ser Gln Arg Ser Asp Val Gly Trp Arg Leu
 1175 1180 1185

ttt gat gac agc aca gtg aca acg gta gac gag agc cag gtt gtg acg 3957
 Phe Asp Asp Ser Thr Val Thr Val Asp Glu Ser Gln Val Val Thr
 1190 1195 1200

cgt tat gcc tat gta ctc ttc tac cgc cgg cgg aac tct cct gtg gag 4005
 Arg Tyr Ala Tyr Val Leu Phe Tyr Arg Arg Arg Asn Ser Pro Val Glu
 1205 1210 1215

agg ccc ccc agg gca ggt cac tct gag cac cac cca gac cta ggc cct 4053
 Arg Pro Pro Arg Ala Gly His Ser Glu His His Pro Asp Leu Gly Pro
 1220 1225 1230 1235

gca gct gag gct gct gcc agc cag gct tcc cgg att tgg cag gag ctg 4101
 Ala Ala Glu Ala Ala Ala Ser Gln Ala Ser Arg Ile Trp Gln Glu Leu
 1240 1245 1250

gag gct gag gag gag ccg gtg cct gag ggg tct ggg ccc ctg ggt ccc 4149
 Glu Ala Glu Glu Glu Pro Val Pro Glu Gly Ser Gly Pro Leu Gly Pro
 1255 1260 1265

tgg ggg ccc caa gac tgg gtg ggc ccc cta cca cgt ggc cct acc aca 4197
 Trp Gly Pro Gln Asp Trp Val Gly Pro Leu Pro Arg Gly Pro Thr Thr
 1270 1275 1280

cca gat gag ggc tgc ctc cgg tac ttt gtc ctg ggc acc gtg gcg gct 4245
 Pro Asp Glu Gly Cys Leu Arg Tyr Phe Val Leu Gly Thr Val Ala Ala
 1285 1290 1295

ttg gtg gcc ctc gtg ctc aac gtg ttc tat cct ctg gta tcc cag agt 4293
 Leu Val Ala Leu Val Leu Asn Val Phe Tyr Pro Leu Val Ser Gln Ser
 1300 1305 1310 1315

cgc tgg aga tga gct cgccctgcagg cagctgctgt gagctggcct acctgcctgc 4348
 Arg Trp Arg *

cccaggccat gcctgccttt gttgtgggga acacctctgg gctttgggcc tcagcttatg 4408

catctgggtgg gagaggggtgg ggaggttgtg gccctgcag gggcagagta tcctaggggtg 4468

tgtatccatc tggctgtctg tccattcatc ctgctgctct gacccttggc ctcaggcttg 4528
 gccctgcccc agctacttcc tgtacttaaa agtggttaata aaaccagact attcaggccc 4588
 aaaaaaaaaa aa 4600

<210> 3
 <211> 2663
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> CDS
 <222> (115)..(2157)

<400> 3
 cggaattccc gggtcgacga tttcgtcgcg gccggctgag tcctctccag ccgcgagagg 60
 cgtttctcca tccgcggctc ggcgcgtcgc tctgagcccc cgcgcccagg tggg atg 117
 Met
 1
 gaa gaa gcc tgt cag gtg cag aca act aag cga ggg gat cct cat gag 165
 Glu Glu Ala Cys Gln Val Gln Thr Thr Lys Arg Gly Asp Pro His Glu
 5 10 15
 tta aga aac ata ttt cta cag tat gcc agt act gag gtt gat gga gag 213
 Leu Arg Asn Ile Phe Leu Gln Tyr Ala Ser Thr Glu Val Asp Gly Glu
 20 25 30
 cgt tac atg acc cca gaa gac ttt gtt cag cgc tat ctt gga ctg tat 261
 Arg Tyr Met Thr Pro Glu Asp Phe Val Gln Arg Tyr Leu Gly Leu Tyr
 35 40 45
 aat gat cca aat agt aac cca aag atc gtg cag ctc ttg gca gga gta 309
 Asn Asp Pro Asn Ser Asn Pro Lys Ile Val Gln Leu Leu Ala Gly Val
 50 55 60 65
 gct gat caa acc aag gat ggg ttg atc tcc tat caa gag ttt ttg gca 357
 Ala Asp Gln Thr Lys Asp Gly Leu Ile Ser Tyr Gln Glu Phe Leu Ala
 70 75 80
 ttt gaa tct gtt tta tgt gct cca gat tcc atg ttc ata gtg gct ttc 405
 Phe Glu Ser Val Leu Cys Ala Pro Asp Ser Met Phe Ile Val Ala Phe
 85 90 95
 cag ttg ttt gac aag agt gga aat gga gag gtg aca ttt gaa aat gtc 453
 Gln Leu Phe Asp Lys Ser Gly Asn Gly Glu Val Thr Phe Glu Asn Val
 100 105 110
 aaa gaa att ttt gga cag act att att cat cat cat atc cct ttt aac 501
 Lys Glu Ile Phe Gly Gln Thr Ile Ile His His His Ile Pro Phe Asn
 115 120 125

tgg gat tgt gaa ttt atc cga ctg cat ttt ggg cat aac cgg aag aag	549
Trp Asp Cys Glu Phe Ile Arg Leu His Phe Gly His Asn Arg Lys Lys	
130 135 140 145	
cat ctt aac tac aca gaa ttc acg cag ttt ctc cag gag ctg caa ttg	597
His Leu Asn Tyr Thr Glu Phe Thr Gln Phe Leu Gln Glu Leu Gln Leu	
150 155 160	
gaa cat gca aga caa gcc ttt gca ctc aaa gac aaa agc aaa agt ggc	645
Glu His Ala Arg Gln Ala Phe Ala Leu Lys Asp Lys Ser Lys Ser Gly	
165 170 175	
atg att tct ggt ctg gat ttc agt gac atc atg gtt acc att aga tct	693
Met Ile Ser Gly Leu Asp Phe Ser Asp Ile Met Val Thr Ile Arg Ser	
180 185 190	
cac atg ctt act cct ttt gtg gag gag aac tta gtt tca gca gct gga	741
His Met Leu Thr Pro Phe Val Glu Glu Asn Leu Val Ser Ala Ala Gly	
195 200 205	
gga agt atc tca cac cag gtt agc ttc tcc tac ttc aat gca ttt aac	789
Gly Ser Ile Ser His Gln Val Ser Phe Ser Tyr Phe Asn Ala Phe Asn	
210 215 220 225	
tcg tta ctg aat aac atg gag ctt gtt cgt aag ata tat agc act cta	837
Ser Leu Leu Asn Asn Met Glu Leu Val Arg Lys Ile Tyr Ser Thr Leu	
230 235 240	
gct ggc aca agg aaa gat gtt gaa gtc aca aag gag gaa ttt gcc cag	885
Ala Gly Thr Arg Lys Asp Val Glu Val Thr Lys Glu Glu Phe Ala Gln	
245 250 255	
agt gcc ata cgc tat gga caa gtc aca cca cta gaa att gat att cta	933
Ser Ala Ile Arg Tyr Gly Gln Val Thr Pro Leu Glu Ile Asp Ile Leu	
260 265 270	
tat cag ctt gca gac tta tat aat gct tca ggg cgc ttg act ttg gca	981
Tyr Gln Leu Ala Asp Leu Tyr Asn Ala Ser Gly Arg Leu Thr Leu Ala	
275 280 285	
gat att gag aga ata gcc cca ttg gct gag ggg gcc tta cct tac aac	1029
Asp Ile Glu Arg Ile Ala Pro Leu Ala Glu Gly Ala Leu Pro Tyr Asn	
290 295 300 305	
ctg gca gaa ctt cag aga cag cag tct cct ggg tta ggc agg cct atc	1077
Leu Ala Glu Leu Gln Arg Gln Gln Ser Pro Gly Leu Gly Arg Pro Ile	
310 315 320	
tgg ctc cag att gcc gag tct gct tac aga ttc act ctg ggc tca gtt	1125
Trp Leu Gln Ile Ala Glu Ser Ala Tyr Arg Phe Thr Leu Gly Ser Val	
325 330 335	
gct gga gct gtg gga gcc act gca gtg tat cct ata gat ctg gtg aag	1173
Ala Gly Ala Val Gly Ala Thr Ala Val Tyr Pro Ile Asp Leu Val Lys	
340 345 350	
acc cga atg caa aac cag cgt ggc tct ggc tct gtt gtt ggg gag cta	1221

Thr Arg Met Gln Asn Gln Arg Gly Ser Gly Ser Val Val Gly Glu Leu
 355 360 365

atg tac aaa aac agc ttt gac tgt ttt aag aaa gtc ttg cgt tat gag 1269
 Met Tyr Lys Asn Ser Phe Asp Cys Phe Lys Lys Val Leu Arg Tyr Glu
 370 375 380 385

ggc ttc ttt gga ctc tac agg ggt ctg ata cca caa ctt ata ggg gtt 1317
 Gly Phe Phe Gly Leu Tyr Arg Gly Leu Ile Pro Gln Leu Ile Gly Val
 390 395 400

gct cca gaa aag gcc att aaa ctg act gtt aat gat ttt gtt cgg gac 1365
 Ala Pro Glu Lys Ala Ile Lys Leu Thr Val Asn Asp Phe Val Arg Asp
 405 410 415

aaa ttt acc aga aga gat ggc tct gtt cca ctt cca gca gaa gtt ctt 1413
 Lys Phe Thr Arg Arg Asp Gly Ser Val Pro Leu Pro Ala Glu Val Leu
 420 425 430

gct gga ggc tgt gct gga ggc tct cag gtc att ttt acc aac cca ttg 1461
 Ala Gly Gly Cys Ala Gly Gly Ser Gln Val Ile Phe Thr Asn Pro Leu
 435 440 445

gag ata gtg aag att cgt ctg caa gta gct gga gag atc acc acg gga 1509
 Glu Ile Val Lys Ile Arg Leu Gln Val Ala Gly Glu Ile Thr Thr Gly
 450 455 460 465

ccc aga gtc agc gcc ctg aat gtg ctc cgg gac ttg gga att ttt ggt 1557
 Pro Arg Val Ser Ala Leu Asn Val Leu Arg Asp Leu Gly Ile Phe Gly
 470 475 480

ctg tat aag ggt gcc aaa gcg tgt ttc ctc cga gac att ccc ttc tct 1605
 Leu Tyr Lys Gly Ala Lys Ala Cys Phe Leu Arg Asp Ile Pro Phe Ser
 485 490 495

gca atc tat ttt cct gtt tat gct cat tgc aaa cta ctt ctg gct gat 1653
 Ala Ile Tyr Phe Pro Val Tyr Ala His Cys Lys Leu Leu Leu Ala Asp
 500 505 510

gaa aat gga cac gtg gga ggt tta aat ctt ctt gca gct gga gcc atg 1701
 Glu Asn Gly His Val Gly Gly Leu Asn Leu Leu Ala Ala Gly Ala Met
 515 520 525

gca ggt gtc cca gct gca tct ctg gtg acc cct gct gat gtc atc aag 1749
 Ala Gly Val Pro Ala Ala Ser Leu Val Thr Pro Ala Asp Val Ile Lys
 530 535 540 545

aca aga ctg cag gtg gct gcc cgc gct ggc cag acg aca tac agt ggt 1797
 Thr Arg Leu Gln Val Ala Ala Arg Ala Gly Gln Thr Thr Tyr Ser Gly
 550 555 560

gtc atc gac tgt ttc agg aag att ctc cgg gaa gaa ggg ccc tca gca 1845
 Val Ile Asp Cys Phe Arg Lys Ile Leu Arg Glu Glu Gly Pro Ser Ala
 565 570 575

ttt tgg aaa ggg act gca gct cga gtg ttt cga tcc tct ccc cag ttt 1893
 Phe Trp Lys Gly Thr Ala Ala Arg Val Phe Arg Ser Ser Pro Gln Phe

<223> n = a,t,c or g

<400> 4

cgcgaggcaa canatgac	atg ttg gcc ttc ctg tct ggg atg ccg gtg acc	51
	Met Leu Ala Phe Leu Ser Gly Met Pro Val Thr	
	1 5 10	
aga aac acc aag tac ctc gac ctc aag aat tca caa gag atg ctc cgc	99	
Arg Asn Thr Lys Tyr Leu Asp Leu Lys Asn Ser Gln Glu Met Leu Arg		
	15 20 25	
tac aaa gag gtc tgc tac tac atg ctc ttt gcc ctg gct gcc tac ggg	147	
Tyr Lys Glu Val Cys Tyr Tyr Met Leu Phe Ala Leu Ala Ala Tyr Gly		
	30 35 40	
tgg ccc atg tac ctg atg cgg aag ccc gcc tgc ggc ctc tgc caa ctg	195	
Trp Pro Met Tyr Leu Met Arg Lys Pro Ala Cys Gly Leu Cys Gln Leu		
	45 50 55	
gct cgg tcc tgc tgc tgt tgc ctg tgt cct gcg agg ccg cgg ttc gcc	243	
Ala Arg Ser Cys Ser Cys Cys Leu Cys Pro Ala Arg Pro Arg Phe Ala		
	60 65 70 75	
cct gga gtc acc atc gag gaa gac aac tgc tgt ggc tgt aat gcc att	291	
Pro Gly Val Thr Ile Glu Glu Asp Asn Cys Cys Gly Cys Asn Ala Ile		
	80 85 90	
gcc atc cgg cgc cac ttc ctg gac gag aac atg act gcg gtg gac atc	339	
Ala Ile Arg Arg His Phe Leu Asp Glu Asn Met Thr Ala Val Asp Ile		
	95 100 105	
gtc tat acc tcc tgc cat gat gcg gtc tat gaa acg ccc ttc tac gtg	387	
Val Tyr Thr Ser Cys His Asp Ala Val Tyr Glu Thr Pro Phe Tyr Val		
	110 115 120	
gcg gtg gac cat gac aag aag aaa gtg gtg atc agt atc cgg ggg acc	435	
Ala Val Asp His Asp Lys Lys Lys Val Val Ile Ser Ile Arg Gly Thr		
	125 130 135	
ctg tcc ccc aag gat gcc ctg act gac ctg acg ggt gat gct gag cgc	483	
Leu Ser Pro Lys Asp Ala Leu Thr Asp Leu Thr Gly Asp Ala Glu Arg		
	140 145 150 155	
ctc ccc gtg gag ggg cac cac ggc acc tgg ctg ggc cac aag ggt atg	531	
Leu Pro Val Glu Gly His His Gly Thr Trp Leu Gly His Lys Gly Met		
	160 165 170	
gtc ctc tca gct gag tac atc aag aag aaa ctg gag cag gag atg gtc	579	
Val Leu Ser Ala Glu Tyr Ile Lys Lys Lys Leu Glu Gln Glu Met Val		
	175 180 185	
ctg tcc cag gcc ttt ggg cga gac ctg ggc cgc gga acc aaa cac tac	627	
Leu Ser Gln Ala Phe Gly Arg Asp Leu Gly Arg Gly Thr Lys His Tyr		
	190 195 200	
ggc ctg att gtg gtg ggc cac tcc ctg ggc gcg ggc act gct gcc atc	675	
Gly Leu Ile Val Val Gly His Ser Leu Gly Ala Gly Thr Ala Ala Ile		

atg ccc act ggc ctt gcc ctg gag ctg ccg act gca gac cac cgc aac	1395
Met Pro Thr Gly Leu Ala Leu Glu Leu Pro Thr Ala Asp His Arg Asn	
445 450 455	
agc agc gtc agg agc aag tcc cag tct gag atg agc ctg gag ggc ttc	1443
Ser Ser Val Arg Ser Lys Ser Gln Ser Glu Met Ser Leu Glu Gly Phe	
460 465 470 475	
tgc gag ggg cgg ctg ctg tgc cca gtg gtt gcg gcg gcg gcc cgc cag	1491
Ser Glu Gly Arg Leu Leu Ser Pro Val Val Ala Ala Ala Ala Arg Gln	
480 485 490	
gac ccg gtg gag ctg ctg ctg ctg tct acc cag gag cgg ctg gca gcg	1539
Asp Pro Val Glu Leu Leu Leu Leu Ser Thr Gln Glu Arg Leu Ala Ala	
495 500 505	
gag ctg cag gcc cgg cgg gca cca ctg gcc acc atg gag agc ctc tgc	1587
Glu Leu Gln Ala Arg Arg Ala Pro Leu Ala Thr Met Glu Ser Leu Ser	
510 515 520	
gac act gag tcc ctg tac agc ttc gac tcg cgc cgc tcc tca ggc ttc	1635
Asp Thr Glu Ser Leu Tyr Ser Phe Asp Ser Arg Arg Ser Ser Gly Phe	
525 530 535	
cgc agc atc cgg ggc tcc ccc agc ctc cac gct gtg ctg gag cgt gat	1683
Arg Ser Ile Arg Gly Ser Pro Ser Leu His Ala Val Leu Glu Arg Asp	
540 545 550 555	
gaa ggc cac ctc ttc tac att gac cct gcc atc ccc gag gaa aac cca	1731
Glu Gly His Leu Phe Tyr Ile Asp Pro Ala Ile Pro Glu Glu Asn Pro	
560 565 570	
tcc ctg agc tcg cgc act gag ctg ctg gcg gcc gac agc ctg tcc aag	1779
Ser Leu Ser Ser Arg Thr Glu Leu Leu Ala Ala Asp Ser Leu Ser Lys	
575 580 585	
cac tca cag gac acg cag ccc ctg gag gcg gcc ctg ggc agt ggc ggc	1827
His Ser Gln Asp Thr Gln Pro Leu Glu Ala Ala Leu Gly Ser Gly Gly	
590 595 600	
gtc act cct gag cgg ccc ccc agt gct gcg gcc aat gac gag gag gaa	1875
Val Thr Pro Glu Arg Pro Pro Ser Ala Ala Ala Asn Asp Glu Glu Glu	
605 610 615	
gag gtt ggc ggt ggg ggt ggc ggg ccg gcc tcc cgc ggg gag ctg gcg	1923
Glu Val Gly Gly Gly Gly Gly Gly Pro Ala Ser Arg Gly Glu Leu Ala	
620 625 630 635	
ctg cac aat ggg cgc ctg ggg gac tcg ccc agt cct cag gtg ctg gaa	1971
Leu His Asn Gly Arg Leu Gly Asp Ser Pro Ser Pro Gln Val Leu Glu	
640 645 650	
ttc gcc gag ttc atc gac agc ctc ttc aac ctg gac agc aag agc agc	2019
Phe Ala Glu Phe Ile Asp Ser Leu Phe Asn Leu Asp Ser Lys Ser Ser	
655 660 665	

tcc ttc caa gac ctc tac tgc atg gtg gtg ccc gag agc ccc acc agt Ser Phe Gln Asp Leu Tyr Cys Met Val Val Pro Glu Ser Pro Thr Ser 670 675 680	2067
gac tac gct gag ggc ccc aag tcc ccc agc cag caa gag atc ctg ctc Asp Tyr Ala Glu Gly Pro Lys Ser Pro Ser Gln Glu Ile Leu Leu 685 690 695	2115
cgt gcc cag ttc gag ccc aac ctg gtg ccc aag ccc cca cgg ctc ttt Arg Ala Gln Phe Glu Pro Asn Leu Val Pro Lys Pro Pro Arg Leu Phe 700 705 710 715	2163
gcc ggc tca gcc gac ccc tcc tcg ggc atc tca ctc tcg ccc tcc ttc Ala Gly Ser Ala Asp Pro Ser Ser Gly Ile Ser Leu Ser Pro Ser Phe 720 725 730	2211
ccg ctc agc tcc tcg ggt gag ctc atg gac ctg acg ccc acg ggc ctc Pro Leu Ser Ser Ser Gly Glu Leu Met Asp Leu Thr Pro Thr Gly Leu 735 740 745	2259
agt agc cag gaa tgc ctg gcg gct gac aag atc cgg act tct acc ccc Ser Ser Gln Glu Cys Leu Ala Ala Asp Lys Ile Arg Thr Ser Thr Pro 750 755 760	2307
act ggc cac gga gcc agc ccc gcc aag caa gat gag ctg gtc atc tca Thr Gly His Gly Ala Ser Pro Ala Lys Gln Asp Glu Leu Val Ile Ser 765 770 775	2355
gca cgc tag cccccca gttgcgtggc cagccgggcc caggcaggag caggtggccc Ala Arg * 780	2411
tgtgggcacc tgggtgcctgc cccctgccgg gcagctttaa ggacagaccc ccaggggcag	2471
tttagcctca ggcacaggca tcgctgctga gctgggggtc cgcatecccta cctcagctta	2531
ggacccccag agccaagggtg gctgggatct ggccccacag atggggaaag atggggaagg	2591
gtgtggagtg gggaggagcc tgggcagcct gctgggtggg ccacactcag cctgactgcc	2651
ctccatgggg gcattctggc accccctgct ccaggacagg ccatgggcaa gctgcctccc	2711
atcactgcct gctggctgct ctcccagggg ccagggtggag agcagtgcc cccgacacat	2771
gtattctcat ctgtggtcca ggccggcatc gtccctggcca cccccagat ctggtgcctg	2831
ctggccggcc ccctgggggtg cccctgccga ggtggcctgc agtgctgtac atgtttacag	2891
aagctgctgg gcttggtcca ggatgtgttc tgggcttgca agccccccgc ccaatcatgt	2951
gttcagtagc catcctctga gcagggccca aggagccag gggcctggag gggccagagg	3011
aggggtgggt cagggccgcc ccttctctgc cttgtgcctc tcatgctgcc tcctctgcc	3071
atgggtcctg ggcaccacagg cctgccctgc ctgctggcta cttoctggct taccttctac	3131
ccccaaggat cctcaccacc caaagggtgg tgggcactgc tgtgaccacc ccagctgcag	3191

tgcttaccca cagtgtttct cttggccgta tcagggcacc gctgtgcctc cgcttttctca 6671
tcttgaagat tactgggtccc cagggtaggt cagtgccctt aagcttaggg ggcttgttga 6731
gcatgttctg tggttctgtg tgcaaggcct gaaccatgac agctctggcc cagcgtggcc 6791
tggtcctggt ccctggcaca tccagtgggg cccgccccac ct 6833

<210> 5
<211> 1720
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (197)..(619)

<400> 5
atgatcgctt tcggaaccgg cccggaattc ccgggtcgac ccacgcgtcc gctctttggc 60
actagttcag aatgggtgatg tgtcggcccc ctctgccata ctcagaacac cagaaagcac 120
aaaaccgggt cctgtttgtc agccaccagt gagtcagagc cgctccctgt tttcttctgt 180
cccgccaag ccacca atg tct ctg gag cct caa aat ggg acg tat gca 229
Met Ser Leu Glu Pro Gln Asn Gly Thr Tyr Ala
1 5 10
gga cca gcg cca gca ttc cag cca ttt ttc ttc act gga gca ttt cca 277
Gly Pro Ala Pro Ala Phe Gln Pro Phe Phe Phe Thr Gly Ala Phe Pro
15 20 25
ttt aat atg caa gag ctg gta ctc aag gtg aga att cag aac cca tct 325
Phe Asn Met Gln Glu Leu Val Leu Lys Val Arg Ile Gln Asn Pro Ser
30 35 40
ctt cga gaa aat gat ttc att gaa att gaa ctg gac cga cag gag ctc 373
Leu Arg Glu Asn Asp Phe Ile Glu Ile Glu Leu Asp Arg Gln Glu Leu
45 50 55
acc tac caa gag ttg ctc aaa gtg tgt tgc tgt gag ctg ggt gtt aat 421
Thr Tyr Gln Glu Leu Leu Lys Val Cys Cys Cys Glu Leu Gly Val Asn
60 65 70 75
cca gat caa gtg gag aag atc aga aag tta ccc aat act ctg tta agg 469
Pro Asp Gln Val Glu Lys Ile Arg Lys Leu Pro Asn Thr Leu Leu Arg
80 85 90
aag gac aag gat gtt gct cga ctc caa gat ttc cag gag ctg gaa ctg 517
Lys Asp Lys Asp Val Ala Arg Leu Gln Asp Phe Gln Glu Leu Glu Leu
95 100 105
gtt ctg atg ata agt gaa aat aat ttt ctg ttc aga aat gct gca tcc 565

Val	Leu	Met	Ile	Ser	Glu	Asn	Asn	Phe	Leu	Phe	Arg	Asn	Ala	Ala	Ser			
	110						115					120						
aca	ctg	act	gaa	agg	cct	tgc	tat	aac	agg	aga	gct	tca	aaa	ctg	act	613		
Thr	Leu	Thr	Glu	Arg	Pro	Cys	Tyr	Asn	Arg	Arg	Ala	Ser	Lys	Leu	Thr			
	125					130				135								
tac	taa	tgc	agc	agg	act	ttttat	tca	ctg	agt	tatta	tgac	agt	gtg	cat	cac	ctct		
Tyr	*															669		
140																		
ggg	cca	agga	caag	ccattg	atc	taa	atgc	ctc	agat	gcc	cgg	gag	ggg	gcc	tct	ggt	gcca	
																729		
ctg	cat	agta	tata	cta	aaca	tcatt	ctgcc	aag	gtag	gaa	gcc	cct	gacc	cca	agc	agt		
																789		
ggt	gcc	actc	ttc	caag	cct	ctt	ggt	gcac	agta	aac	ccta	ttg	ctt	gaag	ctt	gaac	ag	
																849		
ctg	aga	agt	gtc	tgg	agag	gcaga	agctg	aag	gtt	ctat	atc	cag	tgt	gtg	ttt	atg	tcc	
																909		
aga	at	gta	ag	ag	ttgt	tct	aag	cag	cagc	tgag	agag	agg	agc	ctat	ttc	tag	ccac	
																969		
tcct	gtt	gac	agt	gcac	ctg	aagg	gct	ggg	atgc	gttt	ctt	ggt	gtt	g	cat	gct	caca	
																1029		
act	ctg	ctga	catt	ggga	aac	ttat	gag	aga	gga	agact	cg	ggaa	agc	aca	gata	ctg	ggac	
																1089		
agat	ggatt	c	tggt	gtg	ggg	aaag	cac	aga	tact	ggac	ag	atg	gatt	tota	gtg	tga	cttg	
																1149		
tgact	gtg	ag	gttt	cct	tata	acata	tttat	aaat	gtt	tact	cagg	gtta	aaaa	gtatt	tta	aga		
																1209		
ata	cag	ttaa	cta	att	gtaa	atat	gct	gtt	aac	caaaa	aga	gctt	tcct	c	ctc	act	tttt	
																1269		
tcct	ttt	gtaa	acact	cat	ga	ctg	ctt	ctct	gtc	atct	ctg	cat	taact	ccc	ct			
																1329		
tcgt	ggt	cac	tag	aggg	ctc	tct	gat	gc	ct	ttaa	agaa	caact	gctt	ttt	aca	aat	gc	
																1389		
cccc	cacccc	cacccc	gccc	atag	agac	ag	ggt	ctc	acta	tgt	ggccc	ag	gct	ggt	ctc	a		
																1449		
aact	tct	ggc	ctta	agt	gat	gct	cct	gtcc	ttg	gcct	ccc	aaag	tgt	gg	gatt	ac	agg	t
																1509		
gtg	agcc	act	gcac	ccag	cc	cact	tttttt	ttact	ctg	aa	gtatt	ccag	tcata	tgt	gt			
																1569		
gtg	taact	gc	atc	ctat	gag	aag	agc	acaa	atatt	gct	gt	tccat	gtt	ct	ccact	ttcat		
																1629		
ttt	ccact	ac	aaat	gaaa	ag	caatt	tttt	ga	gact	gaat	ct	gtt	gct	at	ttt	t	aaag	gttat
																1689		
tgt	ggg	aaac	tgag	ctaa	ag	gag	ttag	cat	c									
																1720		

<210> 6
 <211> 1658
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> CDS

<222> (523) .. (1479)

<400> 6

ctagtaggcc tgtggaacgc ctgcaggtac cgggtccggaa ttcccgggtc gacccacgcg	60
tccgcccacg cgtccgctct gtctcaaata ataataataa taataataat aataataata	120
ataataataa tgtaggggac ttgatgaagg gaaaggatca gatagattct gaaaagaagt	180
acagaacctg ctaaggccat catacctatt gatcggaagt cagtccatca gatttgctct	240
ggaccggtgg tactgagtct aatcactgcg tgtaagaaga tagtaggaaa cattctggat	300
gctaggtgcc agtaatatg atctaaagct taaggactat ggaatggatc tcattgaagt	360
ttcaggcaat ggatgtgggg tagaagaaga aaacttcgaa ggcttaatat ctttcagctc	420
tgaaacatca cacatctaag attcaagagt ttgccgacct aactcggggtt gaaacttttg	480
gctttcgggg gaaagctctg agctcacttt gtgcactgag tg atg tca cca ttt	534
Met Ser Pro Phe	
1	
cta cct gcc acg tat cgg cga agg ttg gga ctc gac tgg tgt ttg atc	582
Leu Pro Ala Thr Tyr Arg Arg Arg Leu Gly Leu Asp Trp Cys Leu Ile	
5 10 15 20	
acg atg gga aaa tca tcc aga aaa ccc cct acc ccc acc cca gag gga	630
Thr Met Gly Lys Ser Ser Arg Lys Pro Pro Thr Pro Thr Pro Glu Gly	
25 30 35	
ccc aca gtc agc gtg aag cag tta ttt tct acg cta cct gtg cgc cat	678
Pro Thr Val Ser Val Lys Gln Leu Phe Ser Thr Leu Pro Val Arg His	
40 45 50	
aag gaa ttt caa agg aat att aag aag aaa cgt gcc tgc ttc ccc ttc	726
Lys Glu Phe Gln Arg Asn Ile Lys Lys Lys Arg Ala Cys Phe Pro Phe	
55 60 65	
gcc ttc tgc cgt gat tgt cag ttt ctt gag ggc tcc cca gcc atg ctt	774
Ala Phe Cys Arg Asp Cys Gln Phe Leu Glu Gly Ser Pro Ala Met Leu	
70 75 80	
cct gta cag cct gca aaa ctt aca gaa cct gct aag gcc atc aaa cct	822
Pro Val Gln Pro Ala Lys Leu Thr Glu Pro Ala Lys Ala Ile Lys Pro	
85 90 95 100	
att gat cgg aag tca gtc cat cag att tgc tct ggg ccg gtg gta ctg	870
Ile Asp Arg Lys Ser Val His Gln Ile Cys Ser Gly Pro Val Val Leu	
105 110 115	
agt cta agc act gcg gtg aag aag ata gta gga aac agt ctg gat gct	918
Ser Leu Ser Thr Ala Val Lys Lys Ile Val Gly Asn Ser Leu Asp Ala	
120 125 130	
ggg gcc act aat att gat cta aag ctt aag gac tat gga atg gat ctc	966
Gly Ala Thr Asn Ile Asp Leu Lys Leu Lys Asp Tyr Gly Met Asp Leu	

<212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (460)..(3621)

<400> 7

```

gtctgctgta ataccctcta ctataggac cactttgtac aagaaagctg ggtacgcgta      60
agcttggggc cctcgaggga ttctctagag cgtccgcggg ggctgccagg gtatttcggg    120
aagggggcgt gaggaggcgg cggcggcagc ggcgggtagg gcaggcagca gagggaagga    180
gaaagaaagg aaggaagagg gcgggggagtc ctacagaggag gaggcgggac cggccggggca  240
cctcacctct cggcggcggc ggcggcggcg gcggcggcgg cgggcggccg gggaggcggc    300
gccgccccct gcccgcggtc ttctcagcgc agtcgggcgc ggaccgctg gtcccgggca    360
gcggccaagg ctactggggc gggagcagtg ggccggtcgg cggcggcagc ggcagcggcg    420
gaggaggagg aggctggagt gggcgcgagg gcgaccgcc      atg gcg ttc ctc aaa    474
                                         Met Ala Phe Leu Lys
                                         1           5

ctc cgt gac cag cca tca ctg gtg caa gct ata ttt aac gga gat cct    522
Leu Arg Asp Gln Pro Ser Leu Val Gln Ala Ile Phe Asn Gly Asp Pro
                        10                15                20

gat gaa gtt cga gca cta ata ttt aag aaa gaa gat gtt aac ttt cag    570
Asp Glu Val Arg Ala Leu Ile Phe Lys Lys Glu Asp Val Asn Phe Gln
                        25                30                35

gac aat gaa aag cga acc cca ttg cac gcc gca gct tac ctt gga gat    618
Asp Asn Glu Lys Arg Thr Pro Leu His Ala Ala Ala Tyr Leu Gly Asp
                        40                45                50

gca gaa atc att gaa ctt ctt att tta tct gga gct aga gtt aat gcc    666
Ala Glu Ile Ile Glu Leu Leu Ile Leu Ser Gly Ala Arg Val Asn Ala
                        55                60                65

aaa gac agc aaa tgg ttg aca cct tta cac aga gca gtt gca tct tgt    714
Lys Asp Ser Lys Trp Leu Thr Pro Leu His Arg Ala Val Ala Ser Cys
                        70                75                80                85

agt gag gaa gca gtt cag gta ctt ttg aag cat tct gca gat gtt aat    762
Ser Glu Glu Ala Val Gln Val Leu Leu Lys His Ser Ala Asp Val Asn
                        90                95                100

gct cga gac aaa aat tgg caa acc cct tta cat ata gct gct gct aat    810
Ala Arg Asp Lys Asn Trp Gln Thr Pro Leu His Ile Ala Ala Ala Asn
                        105                110                115

aaa gct gta aag tgt gct gaa gct ttg gta cct ctt ctg agt aat gta    858
Lys Ala Val Lys Cys Ala Glu Ala Leu Val Pro Leu Leu Ser Asn Val
                        120                125                130

```

aac gta tct gat cga gca ggg agg act gca tta cat cat gca gct ttc	906
Asn Val Ser Asp Arg Ala Gly Arg Thr Ala Leu His His Ala Ala Phe	
135 140 145	
agt gga cat ggt gag atg gtc aaa cta ctc ttg tct aga ggt gcc aat	954
Ser Gly His Gly Glu Met Val Lys Leu Leu Leu Ser Arg Gly Ala Asn	
150 155 160 165	
att aat gct ttt gac aag aaa gat agg cgt gct atc cat tgg gca gca	1002
Ile Asn Ala Phe Asp Lys Lys Asp Arg Arg Ala Ile His Trp Ala Ala	
170 175 180	
tat atg ggt cac att gaa gta gtg aaa ttg ctt gtg tcg cat gga gct	1050
Tyr Met Gly His Ile Glu Val Val Lys Leu Leu Val Ser His Gly Ala	
185 190 195	
gaa gtg aca tgc aag gat aaa aag tct tat aca cct ctt cat gca gca	1098
Glu Val Thr Cys Lys Asp Lys Lys Ser Tyr Thr Pro Leu His Ala Ala	
200 205 210	
gcc tct agt gga atg atc agc gta gtc aag tac ctt cta gat ctt gga	1146
Ala Ser Ser Gly Met Ile Ser Val Val Lys Tyr Leu Leu Asp Leu Gly	
215 220 225	
gtt gat atg aat gaa cca aat gcc tat gga aat aca cct ctt cat gta	1194
Val Asp Met Asn Glu Pro Asn Ala Tyr Gly Asn Thr Pro Leu His Val	
230 235 240 245	
gcc tgc tat aat gga caa gat gtt gta gtg aat gaa ctt ata gac tgt	1242
Ala Cys Tyr Asn Gly Gln Asp Val Val Val Asn Glu Leu Ile Asp Cys	
250 255 260	
ggg gct att gtg aat caa aag aat gaa aaa gga ttt act cct ttg cac	1290
Gly Ala Ile Val Asn Gln Lys Asn Glu Lys Gly Phe Thr Pro Leu His	
265 270 275	
ttt gct gct gca tca aca cat gga gca ttg tgt tta gag ctt cta gtt	1338
Phe Ala Ala Ala Ser Thr His Gly Ala Leu Cys Leu Glu Leu Leu Val	
280 285 290	
ggc aat ggg gcc gat gtc aat atg aag agt aaa gat ggg aaa acc cca	1386
Gly Asn Gly Ala Asp Val Asn Met Lys Ser Lys Asp Gly Lys Thr Pro	
295 300 305	
cta cac atg act gct ctc cac ggt aga ttc tcc cga tca caa acc att	1434
Leu His Met Thr Ala Leu His Gly Arg Phe Ser Arg Ser Gln Thr Ile	
310 315 320 325	
atc cag agt gga gct gta atc gac tgt gag gat aag aat gga aat acc	1482
Ile Gln Ser Gly Ala Val Ile Asp Cys Glu Asp Lys Asn Gly Asn Thr	
330 335 340	
cct ttg cac ata gca gca cgg tat ggc cat gag ctg ctg atc aac act	1530
Pro Leu His Ile Ala Ala Arg Tyr Gly His Glu Leu Leu Ile Asn Thr	
345 350 355	

ctt att aca agt ggt gct gac act gca aag cgt ggc ata cat gga atg Leu Ile Thr Ser Gly Ala Asp Thr Ala Lys Arg Gly Ile His Gly Met 360 365 370	1578
ttc ccc ctc cat ttg gca gcc tta agc ggc ttt tca gat tgc tgc aga Phe Pro Leu His Leu Ala Ala Leu Ser Gly Phe Ser Asp Cys Cys Arg 375 380 385	1626
aaa ctt ctt tct tca gga ttt gat ata gat acc cca gat gat ttt ggc Lys Leu Leu Ser Ser Gly Phe Asp Ile Asp Thr Pro Asp Asp Phe Gly 390 395 400 405	1674
agg act tgt cta cat gca gct gca gct gga ggg aat ttg gag tgc cta Arg Thr Cys Leu His Ala Ala Ala Ala Gly Gly Asn Leu Glu Cys Leu 410 415 420	1722
aac ctt ctg ctg aat act ggt gca gac ttt aat aaa aag gac aaa ttt Asn Leu Leu Leu Asn Thr Gly Ala Asp Phe Asn Lys Lys Asp Lys Phe 425 430 435	1770
ggg aga tct cca ctg cac tac gct gct gcc aac tgc aat tac cag tgc Gly Arg Ser Pro Leu His Tyr Ala Ala Ala Asn Cys Asn Tyr Gln Cys 440 445 450	1818
ctg ttt gct ctt gtg gga tca gga gca agt gtg aat gac ctt gat gaa Leu Phe Ala Leu Val Gly Ser Gly Ala Ser Val Asn Asp Leu Asp Glu 455 460 465	1866
aga ggc tgc aca ccc ctg cac tat gca gct aca tca gac aca gat ggc Arg Gly Cys Thr Pro Leu His Tyr Ala Ala Thr Ser Asp Thr Asp Gly 470 475 480 485	1914
aag tgc ctg gaa tac tta tta aga aac gat gca aat cca ggg atc cgt Lys Cys Leu Glu Tyr Leu Leu Arg Asn Asp Ala Asn Pro Gly Ile Arg 490 495 500	1962
gat aag caa gga tac aac gca gtt cat tat tca gct gct tat ggt cac Asp Lys Gln Gly Tyr Asn Ala Val His Tyr Ser Ala Ala Tyr Gly His 505 510 515	2010
cgt cta tgt ctt cag ctg att gca agt gaa act cct cta gat gtt tta Arg Leu Cys Leu Gln Leu Ile Ala Ser Glu Thr Pro Leu Asp Val Leu 520 525 530	2058
atg gaa acc tca gga aca gac atg ctg agt gat tca gat aat aga gca Met Glu Thr Ser Gly Thr Asp Met Leu Ser Asp Ser Asp Asn Arg Ala 535 540 545	2106
aca ata agc cct tta cac ttg gct gcc tat cat ggt cac cat caa gca Thr Ile Ser Pro Leu His Leu Ala Ala Tyr His Gly His His Gln Ala 550 555 560 565	2154
ctg gaa gtg ttg gta cag tct ttg tta gat ctt gat gtc aga aat agt Leu Glu Val Leu Val Gln Ser Leu Leu Asp Leu Asp Val Arg Asn Ser 570 575 580	2202
agt gga aga aca ccc cta gat ctt gca gct ttt aag ggc cat gtt gaa	2250

Ser Gly Arg Thr Pro Leu Asp Leu Ala Ala Phe Lys Gly His Val Glu	
585 590 595	
tgt gtg gat gta ctc att aat cag gga gcc tca atc tta gta aaa gat	2298
Cys Val Asp Val Leu Ile Asn Gln Gly Ala Ser Ile Leu Val Lys Asp	
600 605 610	
tac att ttg aag agg aca cct att cat gca gca gca aca aat ggt cat	2346
Tyr Ile Leu Lys Arg Thr Pro Ile His Ala Ala Ala Thr Asn Gly His	
615 620 625	
tca gaa tgc tta cgg cta tta ata gga aat gca gaa cca cag aat gca	2394
Ser Glu Cys Leu Arg Leu Leu Ile Gly Asn Ala Glu Pro Gln Asn Ala	
630 635 640 645	
gtg gat att caa gat gga aat gga cag acg cct ctg atg cta tct gtt	2442
Val Asp Ile Gln Asp Gly Asn Gly Gln Thr Pro Leu Met Leu Ser Val	
650 655 660	
ctc aac ggg cac aca gac tgt gtt tac tca ttg ctg aac aaa gga gca	2490
Leu Asn Gly His Thr Asp Cys Val Tyr Ser Leu Leu Asn Lys Gly Ala	
665 670 675	
aat gta gat gcc aaa gat aag tgg gga agg aca gcg ttg cat aga ggg	2538
Asn Val Asp Ala Lys Asp Lys Trp Gly Arg Thr Ala Leu His Arg Gly	
680 685 690	
gca gtt aca ggc cat gaa gaa tgt gta gat gca tta ctt caa cat ggt	2586
Ala Val Thr Gly His Glu Glu Cys Val Asp Ala Leu Leu Gln His Gly	
695 700 705	
gct aag tgc tta ctt cgg gat agc agg ggc cgg acg cct ata cac ctg	2634
Ala Lys Cys Leu Leu Arg Asp Ser Arg Gly Arg Thr Pro Ile His Leu	
710 715 720 725	
tct gct gcc tgt gga cac att ggt gtt ctt gga gcc ctt ttg cag tca	2682
Ser Ala Ala Cys Gly His Ile Gly Val Leu Gly Ala Leu Leu Gln Ser	
730 735 740	
gca gca tct atg gat gca aat cca gcc aca gca gac aat cat gga tat	2730
Ala Ala Ser Met Asp Ala Asn Pro Ala Thr Ala Asp Asn His Gly Tyr	
745 750 755	
acg gca ctt cac tgg gct tgc tac aat ggt cac gag aca tgt gta gaa	2778
Thr Ala Leu His Trp Ala Cys Tyr Asn Gly His Glu Thr Cys Val Glu	
760 765 770	
ctg ctt tta gaa cag gaa gtt ttc cag aaa acg gaa gga aat gct ttt	2826
Leu Leu Leu Glu Gln Glu Val Phe Gln Lys Thr Glu Gly Asn Ala Phe	
775 780 785	
agt cca ttg cat tgt gcc gtg ata aat gac aac gaa ggt gct gct gag	2874
Ser Pro Leu His Cys Ala Val Ile Asn Asp Asn Glu Gly Ala Ala Glu	
790 795 800 805	
atg tta att gat aca tta ggt gcc agc att gtg aac gcc aca gat tca	2922
Met Leu Ile Asp Thr Leu Gly Ala Ser Ile Val Asn Ala Thr Asp Ser	

810	815	820	
aaa gga aga act cct ctc cat gca gcc gcc ttc aca gac cat gta gag			2970
Lys Gly Arg Thr Pro Leu His Ala Ala Phe Thr Asp His Val Glu			
825	830	835	
tgt tta cag ctg ctg ctc agc cat aat gct caa gtc aat tct gtg gac			3018
Cys Leu Gln Leu Leu Leu Ser His Asn Ala Gln Val Asn Ser Val Asp			
840	845	850	
tct aca ggg aaa aca cct ctt atg atg gct gca gaa aat gga caa aca			3066
Ser Thr Gly Lys Thr Pro Leu Met Met Ala Ala Glu Asn Gly Gln Thr			
855	860	865	
aat aca gtt gag atg ctg gtt agc agt gct agt gca gaa ctg act tta			3114
Asn Thr Val Glu Met Leu Val Ser Ser Ala Ser Ala Glu Leu Thr Leu			
870	875	880	885
caa gat aac agt aaa aat act gcc ctc cat ttg gct tgt agc aag ggt			3162
Gln Asp Asn Ser Lys Asn Thr Ala Leu His Leu Ala Cys Ser Lys Gly			
890	895	900	
cat gaa act agt gcc ttg tta ata ctg gaa aag ata aca gat aga aac			3210
His Glu Thr Ser Ala Leu Leu Ile Leu Glu Lys Ile Thr Asp Arg Asn			
905	910	915	
ctc atc aat gca acc aac gca gcc ttg caa aca cct ctg cat gtt gct			3258
Leu Ile Asn Ala Thr Asn Ala Ala Leu Gln Thr Pro Leu His Val Ala			
920	925	930	
gcc cga aat ggg cta aca atg gtg gtt cag gaa ctt ttg gga aaa gga			3306
Ala Arg Asn Gly Leu Thr Met Val Val Gln Glu Leu Leu Gly Lys Gly			
935	940	945	
gca agt gtg ctt gca gta gat gaa aat ggc tat acc cca gct ttg gcc			3354
Ala Ser Val Leu Ala Val Asp Glu Asn Gly Tyr Thr Pro Ala Leu Ala			
950	955	960	965
tgt gct ccc aat aag gat gtg gct gat tgc ctg gct ctc att ttg gcc			3402
Cys Ala Pro Asn Lys Asp Val Ala Asp Cys Leu Ala Leu Ile Leu Ala			
970	975	980	
acc atg atg cct gtc tca tca agt agt cct tta tca tcc tta aca ttc			3450
Thr Met Met Pro Val Ser Ser Ser Ser Pro Leu Ser Ser Leu Thr Phe			
985	990	995	
aat gcc att aac cgt tat acc aac acc tca aaa aca gtc agc ttt gaa			3498
Asn Ala Ile Asn Arg Tyr Thr Asn Thr Ser Lys Thr Val Ser Phe Glu			
1000	1005	1010	
gct ttg ccc atc atg agg aat gaa cct agc tcc tat tgc agt ttc aat			3546
Ala Leu Pro Ile Met Arg Asn Glu Pro Ser Ser Tyr Cys Ser Phe Asn			
1015	1020	1025	
aac att gga ggg gaa cag gag tac tta tac act gac gtg gat gag ctc			3594
Asn Ile Gly Gly Glu Gln Glu Tyr Leu Tyr Thr Asp Val Asp Glu Leu			
1030	1035	1040	1045

aac gac tcc gat tct gag acc tac tga gaggc tgaggaggag ggagttctca 3646
 Asn Asp Ser Asp Ser Glu Thr Tyr *
 1050

cagtaaagct tcaaactgtg ctttttcagg aaacaggcac tttgatattc acgtagaaat 3706

tcaacctaag agggacagtt ccgttgaagc cgt 3739

<210> 8
 <211> 2162
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (387)..(1748)

<400> 8

cgtactgaag tgggtctgcat gagcgaccgg tccggaattc ccgggtcgac gatttcgttc 60

cggttgcact cttcctatag cccagagggc gagagggcct gtggcctggg ggaaggagga 120

cgaggttctg cctggatccc agcagtagga cgctgtgcc tttgggaaca aaggaatagt 180

ctgcctggaa tccctgcaga tcttggggcc ggaggccagt ccaacccttg gagcaggaag 240

aaacgcaaag ttgtcaagaa ccaagtcgag ctgcctcaga gccggcccg agtagctgca 300

gactccgccc gcgacgtgtg cgcgcttctc tgggccagag cgagcctgtt ttgtgctcgg 360

gttaagagat ttgtcccagc tatacc atg ggc cgc act cgg gaa gct ggc tgc 413
 Met Gly Arg Thr Arg Glu Ala Gly Cys
 1 5

gtg gcc gct ggt gtg gtt atc ggg gct ggt gcc tgc tac tgt gta tac 461
 Val Ala Ala Gly Val Val Ile Gly Ala Gly Ala Cys Tyr Cys Val Tyr
 10 15 20 25

aga ctg gct tgg gga aga gac gag aac gag aaa atc tgg gac gaa gac 509
 Arg Leu Ala Trp Gly Arg Asp Glu Asn Glu Lys Ile Trp Asp Glu Asp
 30 35 40

gag gag tct acg gac acc tca gag att ggg gtt gag act gtg aaa gga 557
 Glu Glu Ser Thr Asp Thr Ser Glu Ile Gly Val Glu Thr Val Lys Gly
 45 50 55

gct aaa act aac gct ggg gca ggg tct ggg gcc aaa ctt cag ggt gat 605
 Ala Lys Thr Asn Ala Gly Ala Gly Ser Gly Ala Lys Leu Gln Gly Asp
 60 65 70

tca gag gtc aag cct gag gtg agt ttg gga ctc gag gat tgt ccg ggt 653
 Ser Glu Val Lys Pro Glu Val Ser Leu Gly Leu Glu Asp Cys Pro Gly
 75 80 85

gta aaa gag aag gcc cat tca gga tcc cac agc gga ggt ggt cta gag	701
Val Lys Glu Lys Ala His Ser Gly Ser His Ser Gly Gly Gly Leu Glu	
90 95 100 105	
gcc aag gcc aag gcc ctt ttc aac acg ctg aag gaa cag gca agt gca	749
Ala Lys Ala Lys Ala Leu Phe Asn Thr Leu Lys Glu Gln Ala Ser Ala	
110 115 120	
aag gca ggc aaa ggg gct agg gtg ggt acc atc tct ggg aac agg acc	797
Lys Ala Gly Lys Gly Ala Arg Val Gly Thr Ile Ser Gly Asn Arg Thr	
125 130 135	
ctt gca ccg agt tta ccc tgc cca gga ggc agg ggt gga ggc tgc cac	845
Leu Ala Pro Ser Leu Pro Cys Pro Gly Gly Arg Gly Gly Gly Cys His	
140 145 150	
ccc acc agg agt gga tct agg gcc ggg ggc agg gca agt gga aaa tcc	893
Pro Thr Arg Ser Gly Ser Arg Ala Gly Gly Arg Ala Ser Gly Lys Ser	
155 160 165	
aag gga aag gcc cga agt aag agc acc agg gct cca gct aca aca tgg	941
Lys Gly Lys Ala Arg Ser Lys Ser Thr Arg Ala Pro Ala Thr Thr Trp	
170 175 180 185	
cct gtc cgg aga ggc aag ttc aac ttt cct tat aaa att gat gat att	989
Pro Val Arg Arg Gly Lys Phe Asn Phe Pro Tyr Lys Ile Asp Asp Ile	
190 195 200	
ctg agt gct ccc gac ctc caa aag gtc ctc aac atc ctg gag cga aca	1037
Leu Ser Ala Pro Asp Leu Gln Lys Val Leu Asn Ile Leu Glu Arg Thr	
205 210 215	
aat gat cct ttt att caa gaa gta gcc ttg gtc act ctg ggt aac aat	1085
Asn Asp Pro Phe Ile Gln Glu Val Ala Leu Val Thr Leu Gly Asn Asn	
220 225 230	
gca gca tat tca ttt aac cag aat gcc ata cgt gaa ttg ggt ggt gtc	1133
Ala Ala Tyr Ser Phe Asn Gln Asn Ala Ile Arg Glu Leu Gly Gly Val	
235 240 245	
cca att att gca aaa ctg ata aaa aca aaa gac ccc ata att agg gaa	1181
Pro Ile Ile Ala Lys Leu Ile Lys Thr Lys Asp Pro Ile Ile Arg Glu	
250 255 260 265	
aag act tac aat gcc ctt aat aac ttg agt gtg aac gca gaa aat cag	1229
Lys Thr Tyr Asn Ala Leu Asn Asn Leu Ser Val Asn Ala Glu Asn Gln	
270 275 280	
ggc aag att aag acg tac atc agt caa gtg tgt gat gac acc atg gtc	1277
Gly Lys Ile Lys Thr Tyr Ile Ser Gln Val Cys Asp Asp Thr Met Val	
285 290 295	
tgt cgc ttg gac tca gct gtg cag atg gct ggg cta aga ctg tta acc	1325
Cys Arg Leu Asp Ser Ala Val Gln Met Ala Gly Leu Arg Leu Leu Thr	
300 305 310	

aac atg act gtg act aat cat tac caa cat ttg ctt tcc tat tct ttt	1373
Asn Met Thr Val Thr Asn His Tyr Gln His Leu Leu Ser Tyr Ser Phe	
315 320 325	
cca gac ttt ttt gct ttg tta ttc ctg gga aat cac ttc acc aag ata	1421
Pro Asp Phe Phe Ala Leu Leu Phe Leu Gly Asn His Phe Thr Lys Ile	
330 335 340 345	
cag att atg aaa cta att ata aac ttt act gaa aat cca gcc atg aca	1469
Gln Ile Met Lys Leu Ile Ile Asn Phe Thr Glu Asn Pro Ala Met Thr	
350 355 360	
aga gag ctg gtc agt tgt aaa gta cca tca gaa ttg att tcc ctc ttt	1517
Arg Glu Leu Val Ser Cys Lys Val Pro Ser Glu Leu Ile Ser Leu Phe	
365 370 375	
aat aaa gaa tgg gat aga gag att ctt ctt aat atc ctt acc cta ttt	1565
Asn Lys Glu Trp Asp Arg Glu Ile Leu Leu Asn Ile Leu Thr Leu Phe	
380 385 390	
gag aat ata aat gac aac ata aaa aat gaa ggg ctc gca tca tcc agg	1613
Glu Asn Ile Asn Asp Asn Ile Lys Asn Glu Gly Leu Ala Ser Ser Arg	
395 400 405	
aaa gaa ttc agc aga agt tca ctt ttt ttc tta ttc aaa gag tct gga	1661
Lys Glu Phe Ser Arg Ser Ser Leu Phe Phe Leu Phe Lys Glu Ser Gly	
410 415 420 425	
gtt tgt gtt aag aaa atc aaa gca cta gca aat cac aat gat ctg gtg	1709
Val Cys Val Lys Lys Ile Lys Ala Leu Ala Asn His Asn Asp Leu Val	
430 435 440	
gtg aaa gta aaa gtc ctg aaa gta tta acc aaa ctc taa tttggagtct	1758
Val Lys Val Lys Val Leu Lys Val Leu Thr Lys Leu *	
445 450	
gtcccaaaca atattgagat atttgcagtt ggtacgatgt gatttgtaaa ttctttgttt	1818
ttcattgtgc gtatatggta aagagatctt ttcagctgct attttggaaat aatgactatc	1878
atatatcata acagtgactg atgttggttg taatggttgg gtttaggatg aaccatttta	1938
aggatgccaa atgaaatatt agtattttgta cacagaaaga atttattgat ttgatcttat	1998
tacctagatt gagatttttt aatctttcct ctacctaaac tgacaatgaa ttggttatac	2058
atcatgcata agctacactt ttatattagt ttatatttgt tattctaaga cttgtgtttc	2118
atcaataaag ttgtgtttta agcagcagaa aaaaaaaaaa aaaa	2162

<210> 9

<211> 1614

<212> DNA

<213> Homo sapiens

<220>
 <221> CDS
 <222> (202) .. (969)

<400> 9

aagctggtac gcctgcaggt accggtccgg aattcccggg tcgacgattt cgtcaaagag	60
acaaactcca ttttcttatg aatggaaagt gaaaaccctt gttccgctta aattgggttc	120
cttcctgtcc tgagaaacat agagaccccc aaaagggaag cagaggagag aaagtccac	180
accagacccc cgcgagaaga g atg acc atg acc acc atg cca gaa agt ctc	231
Met Thr Met Thr Thr Met Pro Glu Ser Leu	
1 5 10	
aac agc ccc gtg tcg ggc aag gcg gtg ttt atg gag ttt ggg ccg ccc	279
Asn Ser Pro Val Ser Gly Lys Ala Val Phe Met Glu Phe Gly Pro Pro	
15 20 25	
aac cag caa atg tct cct tcc ccc atg tcc cac ggg cac tac tcc atg	327
Asn Gln Gln Met Ser Pro Ser Pro Met Ser His Gly His Tyr Ser Met	
30 35 40	
cac tgt tta cac tcg gcg ggc cat tcg cag ccc gac ggc gcc tac agc	375
His Cys Leu His Ser Ala Gly His Ser Gln Pro Asp Gly Ala Tyr Ser	
45 50 55	
tca gcc tcg tcc ttc tcc cga ccg ctg ggc tac ccc tac gtc aac tcg	423
Ser Ala Ser Ser Phe Ser Arg Pro Leu Gly Tyr Pro Tyr Val Asn Ser	
60 65 70	
gtc agc agc cac gca tcc agc ccc tac atc agt tcg gtg cag tcc tac	471
Val Ser Ser His Ala Ser Ser Pro Tyr Ile Ser Ser Val Gln Ser Tyr	
75 80 85 90	
ccg ggc agc gcc agc ctc gcc cag agc cgc ctg gag gac cca ggg gcg	519
Pro Gly Ser Ala Ser Leu Ala Gln Ser Arg Leu Glu Asp Pro Gly Ala	
95 100 105	
gac tcg gag aag agc acg gtg gtg gaa ggc ggt gaa gtg cgc ttc aat	567
Asp Ser Glu Lys Ser Thr Val Val Glu Gly Gly Glu Val Arg Phe Asn	
110 115 120	
ggc aag gga aaa aag atc cgt aaa ccc agg acg att tat tcc agt ttg	615
Gly Lys Gly Lys Lys Ile Arg Lys Pro Arg Thr Ile Tyr Ser Ser Leu	
125 130 135	
cag ttg cag gct ttg aac cgg agg ttc cag caa act cag tac cta gct	663
Gln Leu Gln Ala Leu Asn Arg Arg Phe Gln Gln Thr Gln Tyr Leu Ala	
140 145 150	
ctg ccg gag agg gcg gag ctc gcg gcc tct ttg gga ctc aca cag act	711
Leu Pro Glu Arg Ala Glu Leu Ala Ala Ser Leu Gly Leu Thr Gln Thr	
155 160 165 170	
cag gtc aag atc tgg ttc caa aac aag cga tcc aag ttc aag aag ctg	759

Gln	Val	Lys	Ile	Trp	Phe	Gln	Asn	Lys	Arg	Ser	Lys	Phe	Lys	Lys	Leu	
				175					180					185		
atg	aag	cag	ggg	ggg	gcg	gct	ctg	gag	ggg	agt	gcg	ttg	gcc	aac	ggg	807
Met	Lys	Gln	Gly	Gly	Ala	Ala	Leu	Glu	Gly	Ser	Ala	Leu	Ala	Asn	Gly	
			190					195					200			
cgg	gcc	ctg	tct	gct	ggc	tcc	cca	ccc	gtg	ccg	ccc	ggc	tgg	aac	cct	855
Arg	Ala	Leu	Ser	Ala	Gly	Ser	Pro	Pro	Val	Pro	Pro	Gly	Trp	Asn	Pro	
		205					210					215				
aac	tct	tca	tcc	ggg	aag	ggc	tca	gga	gga	aac	gcg	ggc	tcc	tat	atc	903
Asn	Ser	Ser	Ser	Gly	Lys	Gly	Ser	Gly	Gly	Asn	Ala	Gly	Ser	Tyr	Ile	
	220					225					230					
ccc	agc	tac	aca	tcg	tgg	tac	cct	tca	gcg	cac	caa	gaa	gct	atg	cag	951
Pro	Ser	Tyr	Thr	Ser	Trp	Tyr	Pro	Ser	Ala	His	Gln	Glu	Ala	Met	Gln	
235					240					245					250	
caa	ccc	caa	ctt	atg	tga	ggttgc	ccgcccgtct	ccttcttctg	tccccggccc							1005
Gln	Pro	Gln	Leu	Met	*											
			255													
aggtccctcc	cgctccagg	tccatccatc	ccgtccggaa	aagaaggacc	cagaggggaag											1065
aaggaacagt	ggaggcggga	cgccctccat	ctcctcggag	ccccgcgagg	tccggccccag											1125
caacttccccg	gcatccgcgc	tctagcctga	acctggcct	gggccgagca	gtggcagcag											1185
agagtggcct	cggaggggaag	ccactgccac	ctgagacagc	ccaagcagca	agataaaccc											1245
gtccaccccg	acccgcccgc	cttcagcttt	gtgggactat	caggaaaaaa	caaaacaaaa											1305
acaaaatgta	gaaaaagcaa	aagctctttt	ctgtcctgtc	agtctcctgt	ctcctttttgc											1365
tctgtctgtg	cgctggtaaa	gtccagggtc	tcacccgtcc	gctgtcctca	ttctgcggcc											1425
tcagcaaaaa	gccacaaggt	ctgagcggcc	cgggtcctgc	cgggctgacc	atctccggat											1485
cctgggacac	tctgcctgac	catctgtgta	gctgggtgtg	gaatctgggg	gcattggagg											1545
gaggggggttt	acgaaatcgt	cgacagcgaa	gtcgaccggg	gaattccggg	ccggtacgtg											1605
caggcgtac																1614

<210> 10
 <211> 4351
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (154)..(3240)

<400> 10
ccctctaate tgatcactat agggaaatttg gccctcgagc cgaagaattc ggcacgaggt 60
ctggctggga gcagaaggca gcctcggtct ctgggcggcg gcggcggccc actctgccct 120
ggccgcgctg tgtggtgacc gcaggccccga gac atg agg gcg gcc cgt gct ctg 174
Met Arg Ala Ala Arg Ala Leu
1 5
ctg ccc ctg ctg ctg cag gcc tgc tgg aca gcc gcg cag gat gag ccg 222
Leu Pro Leu Leu Leu Gln Ala Cys Trp Thr Ala Ala Gln Asp Glu Pro
10 15 20
gag acc ccg agg gcc gtg gcc ttc cag gac tgc ccc gtg gac ctg ttc 270
Glu Thr Pro Arg Ala Val Ala Phe Gln Asp Cys Pro Val Asp Leu Phe
25 30 35
ttt gtg ctg gac acc tct gag agc gtg gcc ctg agg ctg aag ccc tac 318
Phe Val Leu Asp Thr Ser Glu Ser Val Ala Leu Arg Leu Lys Pro Tyr
40 45 50 55
ggg gcc ctc gtg gac aaa gtc aag tcc ttc acc aag cgc ttc atc gac 366
Gly Ala Leu Val Asp Lys Val Lys Ser Phe Thr Lys Arg Phe Ile Asp
60 65 70
aac ctg agg gac agg tac tac cgc tgt gac cga aac ctg gtg tgg aac 414
Asn Leu Arg Asp Arg Tyr Tyr Arg Cys Asp Arg Asn Leu Val Trp Asn
75 80 85
gca ggc gcg ctg cac tac agt gac gag gtg gag atc atc caa ggc ctc 462
Ala Gly Ala Leu His Tyr Ser Asp Glu Val Glu Ile Ile Gln Gly Leu
90 95 100
acg cgc atg cct ggc ggc cgc gac gca ctc aaa agc agc gtg gac gcg 510
Thr Arg Met Pro Gly Gly Arg Asp Ala Leu Lys Ser Ser Val Asp Ala
105 110 115
gtc aag tac ttt ggg aag ggc acc tac acc gac tgc gct atc aag aag 558
Val Lys Tyr Phe Gly Lys Gly Thr Tyr Thr Asp Cys Ala Ile Lys Lys
120 125 130 135
ggg ctg gag cag ctc ctc gtg ggg ggc tcc cac ctg aag gag aat aag 606
Gly Leu Glu Gln Leu Leu Val Gly Gly Ser His Leu Lys Glu Asn Lys
140 145 150
tac ctg att gtg gtg acc gac ggg cac ccc ctg gag ggc tac aag gaa 654
Tyr Leu Ile Val Val Thr Asp Gly His Pro Leu Glu Gly Tyr Lys Glu
155 160 165
ccc tgt ggg ggg ctg gag gat gct gtg aac gag gcc aag cac ctg ggc 702
Pro Cys Gly Gly Leu Glu Asp Ala Val Asn Glu Ala Lys His Leu Gly
170 175 180
gtc aaa gtc ttc tgc gtg gcc atc aca ccc gac cac ctg gag ccg cgt 750
Val Lys Val Phe Ser Val Ala Ile Thr Pro Asp His Leu Glu Pro Arg
185 190 195

ctg agc atc atc gcc acg gac cac acg tac cgg cgc aac ttc acg gcg	798
Leu Ser Ile Ile Ala Thr Asp His Thr Tyr Arg Arg Asn Phe Thr Ala	
200 205 210 215	
gct gac tgg ggc cag agc cgc gac gca gag gag gcc atc agc cag acc	846
Ala Asp Trp Gly Gln Ser Arg Asp Ala Glu Glu Ala Ile Ser Gln Thr	
220 225 230	
atc gac acc atc gtg gac atg atc aaa aat aac gtg gag caa gtg tgc	894
Ile Asp Thr Ile Val Asp Met Ile Lys Asn Asn Val Glu Gln Val Cys	
235 240 245	
tgc tcc ttc gaa tgc cag cct gca aga gga cct ccg ggg ctc cgg ggc	942
Cys Ser Phe Glu Cys Gln Pro Ala Arg Gly Pro Pro Gly Leu Arg Gly	
250 255 260	
gac ccc ggc ttt gag gga gaa cga ggc aag ccg ggg ctc cca gga gag	990
Asp Pro Gly Phe Glu Gly Glu Arg Gly Lys Pro Gly Leu Pro Gly Glu	
265 270 275	
aag gga gaa gcc gga gat cct gga aga ccc ggg gac ctc gga cct gtt	1038
Lys Gly Glu Ala Gly Asp Pro Gly Arg Pro Gly Asp Leu Gly Pro Val	
280 285 290 295	
ggg tac cag gga atg aag gga gaa aaa ggg agc cgt ggg gag aag ggc	1086
Gly Tyr Gln Gly Met Lys Gly Glu Lys Gly Ser Arg Gly Glu Lys Gly	
300 305 310	
tcc agg gga ccc aag ggc tac aag gga gag aag ggc aag cgt ggc atc	1134
Ser Arg Gly Pro Lys Gly Tyr Lys Gly Glu Lys Gly Lys Arg Gly Ile	
315 320 325	
gac ggg gtg gac ggc gtg aag ggg gag atg ggg tac cca ggc ctg cca	1182
Asp Gly Val Asp Gly Val Lys Gly Glu Met Gly Tyr Pro Gly Leu Pro	
330 335 340	
ggc tgc aag ggc tcg ccc ggg ttt gac ggc att caa gga ccc cct ggc	1230
Gly Cys Lys Gly Ser Pro Gly Phe Asp Gly Ile Gln Gly Pro Pro Gly	
345 350 355	
ccc aag gga gac ccc ggc gcc ttt gga ctg aaa gga gaa aag ggc gag	1278
Pro Lys Gly Asp Pro Gly Ala Phe Gly Leu Lys Gly Glu Lys Gly Glu	
360 365 370 375	
cct gga gct gac ggg gag gcg ggg aga cca ggg agc tcg gga cca tct	1326
Pro Gly Ala Asp Gly Glu Ala Gly Arg Pro Gly Ser Ser Gly Pro Ser	
380 385 390	
gga gac gag ggc cag ccg gga gag cct ggg ccc ccc gga gag aaa gga	1374
Gly Asp Glu Gly Gln Pro Gly Glu Pro Gly Pro Pro Gly Glu Lys Gly	
395 400 405	
gag gcg ggc gac gag ggg aac cca gga cct gac ggt gcc ccc ggg gag	1422
Glu Ala Gly Asp Glu Gly Asn Pro Gly Pro Asp Gly Ala Pro Gly Glu	
410 415 420	
cgg ggt ggc cct gga gag aga gga cca cgg ggg acc cca ggc acg cgg	1470

Arg Gly Gly Pro Gly Glu Arg Gly Pro Arg Gly Thr Pro Gly Thr Arg	
425 430 435	
gga cca aga gga gac cct ggt gaa gct ggc ccg cag ggt gat cag gga	1518
Gly Pro Arg Gly Asp Pro Gly Glu Ala Gly Pro Gln Gly Asp Gln Gly	
440 445 450 455	
aga gaa ggc ccc gtt ggt gtc cct gga gac ccg ggc gag gct ggc cct	1566
Arg Glu Gly Pro Val Gly Val Pro Gly Asp Pro Gly Glu Ala Gly Pro	
460 465 470	
atc gga cct aaa ggc tac cga ggc gat gag ggt ccc cca ggg tcc gag	1614
Ile Gly Pro Lys Gly Tyr Arg Gly Asp Glu Gly Pro Pro Gly Ser Glu	
475 480 485	
ggt gcc aga gga gcc cca gga cct gcc gga ccc cct gga gac ccg ggg	1662
Gly Ala Arg Gly Ala Pro Gly Pro Ala Gly Pro Pro Gly Asp Pro Gly	
490 495 500	
ctg atg ggt gaa agg gga gaa gac ggc ccc gct gga aat ggc acc gag	1710
Leu Met Gly Glu Arg Gly Glu Asp Gly Pro Ala Gly Asn Gly Thr Glu	
505 510 515	
ggc ttc ccc ggc ttc ccc ggg tat ccg ggc aac agg ggc gct ccc ggg	1758
Gly Phe Pro Gly Phe Pro Gly Tyr Pro Gly Asn Arg Gly Ala Pro Gly	
520 525 530 535	
ata aac ggc acg aag ggc tac ccc ggc ctc aag ggg gac gag gga gaa	1806
Ile Asn Gly Thr Lys Gly Tyr Pro Gly Leu Lys Gly Asp Glu Gly Glu	
540 545 550	
gcc ggg gac ccc gga gac gat aac aac gac att gca ccc cga gga gtc	1854
Ala Gly Asp Pro Gly Asp Asp Asn Asn Asp Ile Ala Pro Arg Gly Val	
555 560 565	
aaa gga gca aag ggg tac cgg ggt ccc gag ggc ccc cag gga ccc cca	1902
Lys Gly Ala Lys Gly Tyr Arg Gly Pro Glu Gly Pro Gln Gly Pro Pro	
570 575 580	
gga cac caa gga ccg cct ggg ccg gac gaa tgc gag att ttg gac atc	1950
Gly His Gln Gly Pro Pro Gly Pro Asp Glu Cys Glu Ile Leu Asp Ile	
585 590 595	
atc atg aaa atg tgc tct tgc tgt gaa tgc aag tgc ggc ccc atc gac	1998
Ile Met Lys Met Cys Ser Cys Cys Glu Cys Lys Cys Gly Pro Ile Asp	
600 605 610 615	
ctc ctg ttc gtg ctg gac agc tca gag agc att ggc ctg cag aac ttc	2046
Leu Leu Phe Val Leu Asp Ser Ser Glu Ser Ile Gly Leu Gln Asn Phe	
620 625 630	
gag att gcc aag gac ttc gtc gtc aag gtc atc gac ccg ctg agc cgg	2094
Glu Ile Ala Lys Asp Phe Val Val Lys Val Ile Asp Arg Leu Ser Arg	
635 640 645	
gac gag ctg gtc aag ttc gag cca ggg cag tcg tac gcg ggt gtg gtg	2142
Asp Glu Leu Val Lys Phe Glu Pro Gly Gln Ser Tyr Ala Gly Val Val	

cgg gcg tcg ctg cag ttc ctg cag aac tac acg gcc ctg gcc agt gcc	2862
Arg Ala Ser Leu Gln Phe Leu Gln Asn Tyr Thr Ala Leu Ala Ser Ala	
890 895 900	
gtc gat gcc atg gac ttt atc aac gac gcc acc gac gtc aac gat gcc	2910
Val Asp Ala Met Asp Phe Ile Asn Asp Ala Thr Asp Val Asn Asp Ala	
905 910 915	
ctg ggc tat gtg acc cgc ttc tac cgc gag gcc tcg tcc ggc gct gcc	2958
Leu Gly Tyr Val Thr Arg Phe Tyr Arg Glu Ala Ser Ser Gly Ala Ala	
920 925 930 935	
aag aag agg ctg ctg ctc ttc tca gat ggc aac tcg cag ggc gcc acg	3006
Lys Lys Arg Leu Leu Leu Phe Ser Asp Gly Asn Ser Gln Gly Ala Thr	
940 945 950	
ccc gct gcc atc gag aag gcc gtg cag gaa gcc cag cgg gca ggc atc	3054
Pro Ala Ala Ile Glu Lys Ala Val Gln Glu Ala Gln Arg Ala Gly Ile	
955 960 965	
gag atc ttc gtg gtg gtc gtg ggc cgc cag gtg aat gag ccc cac atc	3102
Glu Ile Phe Val Val Val Val Gly Arg Gln Val Asn Glu Pro His Ile	
970 975 980	
cgc gtc ctg gtc acc ggc aag acg gcc gag tac gac gtg gcc tac ggc	3150
Arg Val Leu Val Thr Gly Lys Thr Ala Glu Tyr Asp Val Ala Tyr Gly	
985 990 995	
gag agc cac ctg ttc cgt gtc ccc agc tac cag gcc ctg ctc cgc ggt	3198
Glu Ser His Leu Phe Arg Val Pro Ser Tyr Gln Ala Leu Leu Arg Gly	
1000 1005 1010 1015	
gtc ttc cac cag aca gtc tcc agg aag gtg gcg ctg ggc tag cccaccc	3247
Val Phe His Gln Thr Val Ser Arg Lys Val Ala Leu Gly *	
1020 1025	
tgcacgccgg caccaaacc tgtcctccca cccctcccca ctcatcacta aacagagaaa	3307
agcttggaag gccaggacac aacgctgctg cctgctttgt gcagggtcct ccggggctca	3367
gccctgagtt ggcacacact gcgcagggcc ctctggggct cagccctgag ctagtgtcac	3427
ctgcacaggg ccctctgggg ctcagccctg agctggcgtc acctgtgcag ggccctctgg	3487
ggctcagccc tgagctggcc tcacctgggt tccccacccc gggtctctct gccctgccct	3547
cctgcccgcc ctccctctctg cctgcgcagc tccttcctta ggcacctctg tgctgcgtcc	3607
caccagcctg agcaagacgc cctctcgggg cctgtgccgc actagcctcc ctctcctctg	3667
tcccatagc tggtttttcc caccaatcct cacctaacag ttactttaca attaaactca	3727
aagcaagctc ttctcctcag cttggggcag ccattggcct ctgtctcggt ttgggaaacc	3787
aaggtcagga ggccgttgca gacataaatc tcggcgactc ggccccgtct cctgaggggtc	3847

ctgctggtga ccggcctgga ccttggccct acagccctgg aggccgctgc tgaccagcac 3907
 tgaccccgac ctcagagagt actcgcaggg gcgctggctg cactcaagac cctcgagatt 3967
 aacggtgcta acccgtctg ctccctccctc ccgcagagac tggggcctgg actggacatg 4027
 agagcccctt ggtgccacag agggctgtgt cttactagaa acaacgcaa cctctccttc 4087
 ctcagaatag tgatgtgttc gacgttttat caaaggcccc ctttctatgt tcatgttagt 4147
 tttgtcctt ctgtgttttt ttctgaacca tatccatgtt gctgactttt ccaaataaag 4207
 gttttcactc ctcaaaaaaa aaaaaaaaag ggcggccgct ctagagtatc cctcgagggg 4267
 cccaagctta cgcgtacca gctttcttgt acaaagtggg ccctatagtg agtcgtatat 4327
 aagctaggga cgtttcagcg acag 4351

<210> 11
 <211> 1622
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (864) .. (1424)

<400> 11
 taagcttgcg gccgctgaac tacagctctg cgccctgcca ggcggccgca cgctcagggg 60
 cgtggcatgg gtgggtcgtg agttgggcgg ggcccacagg gcgtgcgcga cgcagcggcg 120
 cggcgcgagg cgtaaggggc gtggcgccag tgggcgtggc gtggcgagcgt gcgaagggac 180
 gcggtgcgca tgcgcgtgag ggcttccacg ggtgggtggg atcgaggcct gtcgggtcag 240
 ggcgggttcgc ggggtgctgtc agagctgggc cggggcccct aggcaggccc agacatgtcc 300
 gtccttgtaa gttaaaagct tccatgggag ctttccttcc taatcaagat gcaaatagta 360
 cgggtattccg aacagacact aaaaatagct gtcatctcaa agaatccagt gcttgtgtca 420
 cagtatgaga aagtagatgc tggggaacag cgtttaatga atgaggcatg caagccagcc 480
 agtgatctct ttggaccttg cattctccat cagattggat cacctcccac cctgaggccc 540
 cccaagactt tgaacagttc ttcagtcatc cttacagaaa gataccctct ccagacaaac 600
 gcagtattta tatacgggtcc attggatctc tatgaagcac cagaattatc agtgaagaat 660
 atattaaatg gtcacgggc tactgtaaag catatttcta tcgcttgaga gtaaaactcc 720
 tagaaccagt tcctgtttct gtaacaagat gttcctttag agtcaatgag aacacacaca 780

acctacaaat tcatgcaggg gacatcctga agttcttgaa aaagaagaaa cctgaagatg 840

ccttctgtgt tgtgggaata aca atg att gat ctt tac cca aga gac tcg 890
Met Ile Asp Leu Tyr Pro Arg Asp Ser
1 5

tgg aat ttt gtc ttt gga cag gcc tct ttg aca gat ggt gtg ggg ata 938
Trp Asn Phe Val Phe Gly Gln Ala Ser Leu Thr Asp Gly Val Gly Ile
10 15 20 25

ttc agc ttt gcc agg tat ggc agt gat ttt tat agc atg cac tat aaa 986
Phe Ser Phe Ala Arg Tyr Gly Ser Asp Phe Tyr Ser Met His Tyr Lys
30 35 40

ggc aaa gtg aag aag ctc aag aaa aca tct tca agt gac tat tca att 1034
Gly Lys Val Lys Lys Leu Lys Lys Thr Ser Ser Ser Asp Tyr Ser Ile
45 50 55

ttc gac aac tat tat att cca gaa ata act agt gtt tta cta ctt cga 1082
Phe Asp Asn Tyr Tyr Ile Pro Glu Ile Thr Ser Val Leu Leu Leu Arg
60 65 70

tcc tgt aag act tta acc cat gag atc gga cac ata ttt gga ctg cga 1130
Ser Cys Lys Thr Leu Thr His Glu Ile Gly His Ile Phe Gly Leu Arg
75 80 85

cac tgc cag tgg ctt gca tgc ctc atg caa ggc tcc aac cac ttg gaa 1178
His Cys Gln Trp Leu Ala Cys Leu Met Gln Gly Ser Asn His Leu Glu
90 95 100 105

gaa gct gac cgg cgc cct cta aac ctt tgc cct atc tgt ttg cac aag 1226
Glu Ala Asp Arg Arg Pro Leu Asn Leu Cys Pro Ile Cys Leu His Lys
110 115 120

ttg cag tgt gct gtt ggc ttc agc att gta gaa aga tac aaa gca ctg 1274
Leu Gln Cys Ala Val Gly Phe Ser Ile Val Glu Arg Tyr Lys Ala Leu
125 130 135

gtg agg tgg att gat gat gaa tct tct gac aca cct gga gca act cca 1322
Val Arg Trp Ile Asp Asp Glu Ser Ser Asp Thr Pro Gly Ala Thr Pro
140 145 150

gaa cac agt cac gag gat aat ggg aat tta ccg aaa ccc gtg gaa gcc 1370
Glu His Ser His Glu Asp Asn Gly Asn Leu Pro Lys Pro Val Glu Ala
155 160 165

ttt aag gaa tgg aaa gag tgg ata ata aaa tgc ctg gct gtt ctc caa 1418
Phe Lys Glu Trp Lys Glu Trp Ile Ile Lys Cys Leu Ala Val Leu Gln
170 175 180 185

aaa tga ggaccttcaa ataggagtga ttgaaataaa taactacttg catgttatgc 1474
Lys *

tttcatttgg gtggaatact tcattggaat aaactactga tcttgtgctg tgtcaaagta 1534

acagactaga accttctttc aagtacctga attgaaatga aactcatttt gaataataaa 1594

aactctagaa actcttaaaaa aaaaaaaaa

1622

<210> 12
<211> 882
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (147) .. (284)

<400> 12
gagccgagat cagccactg cactccagcc tgggcaacaa gagcaaaact ctgtctcaat 60
aaaaaaaaag gaaaaccac agccaacatt atcactagt gtaacaggga aaatgtcccc 120
tttgaccccc accccaagta cagaac atg caa gga cgc ctg ttc tca ctg cat 173
Met Gln Gly Arg Leu Phe Ser Leu His
1 5
gtc ttc tgt gtt gta ctg aag ctc cta gct ggt gca gtg agg caa gac 221
Val Phe Cys Val Val Leu Lys Leu Leu Ala Gly Ala Val Arg Gln Asp
10 15 20 25
aaa gaa agg aaa tat agg tgc act ggg aag gaa gaa gaa aca ctg cct 269
Lys Glu Arg Lys Tyr Arg Cys Thr Gly Lys Glu Glu Glu Thr Leu Pro
30 35 40
tta ttc ttt agg tga catgattgtg tgcttttaaa ataataaagg aatcaacaga 324
Leu Phe Phe Arg *
45
aaagttgctt aacctaataga atgagtttat caaagtcaca agatacaagg tcagtataca 384
aaaatcagtt ggatttctac atggtagaaa caactgtaca tggaaaaatg tttaatagt 444
taagatatgt acattggaaa ctatgaaaga gtgtaaaaaa taaagaagtg aaataaatgg 504
agaagatacc accttgatgg atggaagcct taaggtaaag atgctcattc tccccacact 564
gacctgtaca ttccccacag gcctaataca aacccccaca ggcttctgtg tagaaattga 624
catgctgata ctgaaattta tatgaaaatg caaagagtct ggaataacca aaataatttt 684
gtaaaagaac aaagaagact tctactacct gggtataaga cttctctgaa gcacagaagt 744
caaggcagtg tgggtggtggc ataagtaatg taaatcatcc aggtgtggtg gctcaagcca 804
gtcatccagc actttgggag gctgaggcag gagaatcgct agagcccagg agttggaaac 864
cagcctgggc agcatagc 882

<210> 13
 <211> 442
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (58) .. (336)

<400> 13

```

gctccggaat tcccgggaac atttacactg acagcaagta cgccttagct actgtgc      57
atg tac ata aag cca tct acc agg aaa tca ggc tac tca cct cag cag      105
Met Tyr Ile Lys Pro Ser Thr Arg Lys Ser Gly Tyr Ser Pro Gln Gln
   1             5             10             15

gta gct gtg atc cac tgc aaa gga cat caa aaa gaa aac acg gcc gtt      153
Val Ala Val Ile His Cys Lys Gly His Gln Lys Glu Asn Thr Ala Val
           20             25             30

gcc cat agt aac cag aaa gct gat tca gca gct cag gtc act gcc aga      201
Ala His Ser Asn Gln Lys Ala Asp Ser Ala Ala Gln Val Thr Ala Arg
           35             40             45

ctt tca gtc acg cct cca aac ttg ctg ccc aca gtc tcc ttt cca cag      249
Leu Ser Val Thr Pro Pro Asn Leu Leu Pro Thr Val Ser Phe Pro Gln
           50             55             60

cca gat ctg cct gac aat ccc gta tac tca aca aca aca gaa aaa ctg      297
Pro Asp Leu Pro Asp Asn Pro Val Tyr Ser Thr Thr Thr Glu Lys Leu
           65             70             75             80

gct tca gat ctc aga gcc aat aaa aat cag gaa agt tag tagattcttc      346
Ala Ser Asp Leu Arg Ala Asn Lys Asn Gln Glu Ser *
           85             90

ctgactctgg aatcttcata ccttgaactt aaaccagtta cctacagtct accacccatt      406

taagaagagc aaagttacct cagctcctcc ggaggg      442

```

<210> 14
 <211> 2058
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (173) .. (1909)

<400> 14

```

taagcttgcg gccgcccggc ggtgtagatg gtacgcttgc ttgtacgcag caggggaggc      60

tctgacttgc actggctggt ggtccgcttc cggggagagc gccagtgcc cacaggtggc      120

```

aggagcctgt ccagggactc tgcattgctg ctgcagagag ggatgtactc gg	atg Met 1	175
tgg ccc aga gct gcc cac ctc cca ggt acc cgc tgg cat ggc cgg ccc Trp Pro Arg Ala Ala His Leu Pro Gly Thr Arg Trp His Gly Arg Pro	5 10 15	223
aca ggg cat ccc ccc cca ccc cca atg agc ctg ccc gcc tca gcc atg Thr Gly His Pro Pro Pro Pro Pro Met Ser Leu Pro Ala Ser Ala Met	20 25 30	271
cct gtt gag ggg gtg ggg ggg gat gcc ctg tgg gcc ggc cat gcc agc Pro Val Glu Gly Val Gly Gly Asp Ala Leu Trp Ala Gly His Ala Ser	35 40 45	319
ggg tac ctg gga ggt ggc cag ctc tgg gcc aca tcc gag tac atc cct Gly Tyr Leu Gly Gly Gly Gln Leu Trp Ala Thr Ser Glu Tyr Ile Pro	50 55 60 65	367
ctc tgc agc agc aat gca gag tcc ctg gac agg ctc ctg cca cct gtg Leu Cys Ser Ser Asn Ala Glu Ser Leu Asp Arg Leu Leu Pro Pro Val	70 75 80	415
ggc act ggg cgc tct ccc cgg aag cgg acc acc agc cag tgc aag tca Gly Thr Gly Arg Ser Pro Arg Lys Arg Thr Thr Ser Gln Cys Lys Ser	85 90 95	463
gag cct ccc ctg ctg cgt aca agc aag cgt acc atc tac acc gcc ggg Glu Pro Pro Leu Leu Arg Thr Ser Lys Arg Thr Ile Tyr Thr Ala Gly	100 105 110	511
cgg ccg ccc tgg tac aat gaa cac ggc acg caa tcc aaa gag gcc ttc Arg Pro Pro Trp Tyr Asn Glu His Gly Thr Gln Ser Lys Glu Ala Phe	115 120 125	559
gcc atc ggc ttg gga ggc ggc agt gcc tct ggg aag acc act gtg gcc Ala Ile Gly Leu Gly Gly Gly Ser Ala Ser Gly Lys Thr Thr Val Ala	130 135 140 145	607
aga atg atc atc gag gcc ctg gat gtg ccc tgg gtg gtc ttg ctg tcc Arg Met Ile Ile Glu Ala Leu Asp Val Pro Trp Val Val Leu Leu Ser	150 155 160	655
atg gac tcc ttc tac aag gtg ctg act gag cag cag cag gaa cag gcc Met Asp Ser Phe Tyr Lys Val Leu Thr Glu Gln Gln Gln Glu Gln Ala	165 170 175	703
gca cac aac aac ttc aac ttc gac cac cca gat gcc ttt gac ttc gac Ala His Asn Asn Phe Asn Phe Asp His Pro Asp Ala Phe Asp Phe Asp	180 185 190	751
ctc atc att tcc acc ctc aag aag ctg aag cag ggg aag agt gtc aag Leu Ile Ile Ser Thr Leu Lys Lys Leu Lys Gln Gly Lys Ser Val Lys	195 200 205	799

gtg ccc att tat gac ttc acc acg cac agc cgg aag aag gac tgg aaa	847
Val Pro Ile Tyr Asp Phe Thr Thr His Ser Arg Lys Lys Asp Trp Lys	
210 215 220 225	
aca ctg tat ggt gca aac gtc atc atc ttt gag ggc atc atg gcc ttt	895
Thr Leu Tyr Gly Ala Asn Val Ile Ile Phe Glu Gly Ile Met Ala Phe	
230 235 240	
gct gac aag aca ctg ttg gag ctc ctg gac atg aag atc ttt gtg gac	943
Ala Asp Lys Thr Leu Leu Glu Leu Leu Asp Met Lys Ile Phe Val Asp	
245 250 255	
aca gac tcc gac atc cgc ctg gta cgg cgg ctg cgc cgg gac atc agt	991
Thr Asp Ser Asp Ile Arg Leu Val Arg Arg Leu Arg Arg Asp Ile Ser	
260 265 270	
gag cgc ggc cgg gac atc gag ggt gtc atc aag cag tac aac aag ttt	1039
Glu Arg Gly Arg Asp Ile Glu Gly Val Ile Lys Gln Tyr Asn Lys Phe	
275 280 285	
gtc aag ccc tcc ttc gac cag tac atc cag ccc acc atg cgc ctg gca	1087
Val Lys Pro Ser Phe Asp Gln Tyr Ile Gln Pro Thr Met Arg Leu Ala	
290 295 300 305	
gac atc gtg gtc ccc aga ggg agc ggc aac acg gtg gcc atc gac ctg	1135
Asp Ile Val Val Pro Arg Gly Ser Gly Asn Thr Val Ala Ile Asp Leu	
310 315 320	
att gtg cag cac gtg cac agc cag ctg gag gag cgt gaa ctc agc gtc	1183
Ile Val Gln His Val His Ser Gln Leu Glu Glu Arg Glu Leu Ser Val	
325 330 335	
agg gct gcg ctg gcc tcg gca cac cag tgc cac ccg ctg ccc cgg acg	1231
Arg Ala Ala Leu Ala Ser Ala His Gln Cys His Pro Leu Pro Arg Thr	
340 345 350	
ctg agc gtc ctg aag agc acg ccg cag gta cgg ggc atg cac acc atc	1279
Leu Ser Val Leu Lys Ser Thr Pro Gln Val Arg Gly Met His Thr Ile	
355 360 365	
atc agg gac aag gag acc agt cgc gac gag ttc atc ttc tac tcc aag	1327
Ile Arg Asp Lys Glu Thr Ser Arg Asp Glu Phe Ile Phe Tyr Ser Lys	
370 375 380 385	
aga ctg atg cgg ctg ctc atc gag cac gcg ctc tcc ttc ctg ccc ttt	1375
Arg Leu Met Arg Leu Leu Ile Glu His Ala Leu Ser Phe Leu Pro Phe	
390 395 400	
cag gac tgc gtc gta cag acc ccg cag ggg cag gac tat gcg ggc aag	1423
Gln Asp Cys Val Val Gln Thr Pro Gln Gly Gln Asp Tyr Ala Gly Lys	
405 410 415	
tgc tat gcg ggg aag cag atc acc ggt gtg tcc att ctg cgc gcc ggt	1471
Cys Tyr Ala Gly Lys Gln Ile Thr Gly Val Ser Ile Leu Arg Ala Gly	
420 425 430	
gaa acc atg gag ccc gcg ctg cgc gct gtg tgc aaa gac gtg cgc atc	1519

Glu Thr Met Glu Pro Ala Leu Arg Ala Val Cys Lys Asp Val Arg Ile	
435 440 445	
ggc acc atc ctc atc cag acc aac cag ctt acc ggg gag ccc gag ctc	1567
Gly Thr Ile Leu Ile Gln Thr Asn Gln Leu Thr Gly Glu Pro Glu Leu	
450 455 460 465	
cac tac ctg agg ctg ccc aag gac atc agc gat gac cac gtg atc ctc	1615
His Tyr Leu Arg Leu Pro Lys Asp Ile Ser Asp Asp His Val Ile Leu	
470 475 480	
atg gac tgc acc gtg tcc acg ggc gcg gcg gcc atg atg gca gtg cgc	1663
Met Asp Cys Thr Val Ser Thr Gly Ala Ala Ala Met Met Ala Val Arg	
485 490 495	
gtg ctc ctg gac cac gac gtg cct gag gac aag atc ttt ttg ctg tcg	1711
Val Leu Leu Asp His Asp Val Pro Glu Asp Lys Ile Phe Leu Leu Ser	
500 505 510	
ctg ctc atg gca gag atg ggc gtg cac tca gtg gcc tat gca ttt ccg	1759
Leu Leu Met Ala Glu Met Gly Val His Ser Val Ala Tyr Ala Phe Pro	
515 520 525	
cga gtg aga atc atc acc acg gcg gtg gac aag cgg gtc aat gac ctt	1807
Arg Val Arg Ile Ile Thr Thr Ala Val Asp Lys Arg Val Asn Asp Leu	
530 535 540 545	
ttc cgc atc atc cca ggc att ggg aac ttt ggc gac cgc tac ttt ggg	1855
Phe Arg Ile Ile Pro Gly Ile Gly Asn Phe Gly Asp Arg Tyr Phe Gly	
550 555 560	
aca gac gcg gtc ccc gat ggc agt gac gag gag gaa gtg gcc tac acg	1903
Thr Asp Ala Val Pro Asp Gly Ser Asp Glu Glu Glu Val Ala Tyr Thr	
565 570 575	
ggg tag ctgcccagtg agccatcccg tccccaccac cctcctcctg cctcctgacc	1959
Gly *	
caggactgct gaatacaaag atgttaattt ttaaaatggt actagtataa tttattctat	2019
gcattttata aaataaataa agcttttagaa aaaaaaaaaa	2058

<210> 15
 <211> 1705
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (545) .. (1672)

aaggatcctt aattttatta atcccccccc ccccggaaggc cccagccgc ctgtgccccg	60
-------------------------------------------------------------------	----

cgggtgcgcgc cccgctccgg gacctgccgc caccgccgcc ccgccctcgg cggcaccac 120
 acccaggcgc gcccgcgcg gcgcccggcc ccgtccctgc ctggaagcac agctgaagat 180
 ggcgagcccc gcgcctccgg agcacgccga ggagggatgc ccggctcctg ccgccgagga 240
 gcaggcgccg ccgtcgccgc caccgccccca ggcatcccc gcagagcggc agcagcagga 300
 ggaggaagcg caggaagctg gggcggcgga gggcgcgggg ttgcaggtgg aggaggccgc 360
 gggccggggc gcggcccgcg taacctggct gctcggggag ccggtgctgt ggctgggctg 420
 ccgcgccgac gagctgctga gctggaagag gccgctgcgg agcctgctcg gcttcgtcgc 480
 tgccaacctg ctgttctggt tccttgcat t gactccatgg agagtatatc acctgatttc 540
 cgtc atg ata ctt ggg cgt gtt att atg caa ata ata aag gat atg gtt 589
 Met Ile Leu Gly Arg Val Ile Met Gln Ile Ile Lys Asp Met Val
 1 5 10 15
 ttg tct aga aca aga ggt gca cag ttg tgg aga agc ctc agt gaa agc 637
 Leu Ser Arg Thr Arg Gly Ala Gln Leu Trp Arg Ser Leu Ser Glu Ser
 20 25 30
 tgg gaa gtt atc aat tcc aaa cca gat gaa aga ccc agg ctc agc cac 685
 Trp Glu Val Ile Asn Ser Lys Pro Asp Glu Arg Pro Arg Leu Ser His
 35 40 45
 tgt att gca gaa tca tgg atg aat ttc agc ata ttt ctt caa gaa atg 733
 Cys Ile Ala Glu Ser Trp Met Asn Phe Ser Ile Phe Leu Gln Glu Met
 50 55 60
 tct ctt ttt aaa cag cag agc cct ggc aag ttt tgt ctc ctg gtc tgt 781
 Ser Leu Phe Lys Gln Gln Ser Pro Gly Lys Phe Cys Leu Leu Val Cys
 65 70 75
 agt gtg tgc aca ttt ttt acg atc ttg gga agt tac att cct ggg gtt 829
 Ser Val Cys Thr Phe Thr Ile Leu Gly Ser Tyr Ile Pro Gly Val
 80 85 90 95
 ata ctc agc tat cta ctg tta ctg tgt gca ttt ttg tgt cca ttg ttt 877
 Ile Leu Ser Tyr Leu Leu Leu Leu Cys Ala Phe Leu Cys Pro Leu Phe
 100 105 110
 aaa tgt aat gat att gga caa aaa att tac agc aaa att aag tca gtt 925
 Lys Cys Asn Asp Ile Gly Gln Lys Ile Tyr Ser Lys Ile Lys Ser Val
 115 120 125
 ctg ctg aaa ctg gat ttt gga att gga gaa tat att aat cag aag aaa 973
 Leu Leu Lys Leu Asp Phe Gly Ile Gly Glu Tyr Ile Asn Gln Lys Lys
 130 135 140
 cgt gag aga tct gaa gca gat aaa gaa aaa agt cac aaa gat gac agt 1021
 Arg Glu Arg Ser Glu Ala Asp Lys Glu Lys Ser His Lys Asp Asp Ser
 145 150 155

gaa tta gac ttt tca gct ctt tgt cct aag att agc ctc acg gtt gct	1069
Glu Leu Asp Phe Ser Ala Leu Cys Pro Lys Ile Ser Leu Thr Val Ala	
160 165 170 175	
gcc aaa gag tta tct gtg tct gac aca gac gtc tca gag gta tcc tgg	1117
Ala Lys Glu Leu Ser Val Ser Asp Thr Asp Val Ser Glu Val Ser Trp	
180 185 190	
act gat aat ggg acc ttc aac ctt tca gaa gga tac act cca cag aca	1165
Thr Asp Asn Gly Thr Phe Asn Leu Ser Glu Gly Tyr Thr Pro Gln Thr	
195 200 205	
gac act tct gat gat ctt gac cga ccc agt gag gaa gtt ttc tct aga	1213
Asp Thr Ser Asp Asp Leu Asp Arg Pro Ser Glu Glu Val Phe Ser Arg	
210 215 220	
gat ctt tca gat ttt cca tct cta gaa aat ggc atg gga aca aat gat	1261
Asp Leu Ser Asp Phe Pro Ser Leu Glu Asn Gly Met Gly Thr Asn Asp	
225 230 235	
gaa gat gaa tta agc ctt ggt ttg ccc act gag ctc aag aga aag aag	1309
Glu Asp Glu Leu Ser Leu Gly Leu Pro Thr Glu Leu Lys Arg Lys Lys	
240 245 250 255	
gaa cag ttg gac agt ggt cac aga cca agc aaa gag acg caa tca gca	1357
Glu Gln Leu Asp Ser Gly His Arg Pro Ser Lys Glu Thr Gln Ser Ala	
260 265 270	
gct ggt ctc acc ctt cct ctg aac agt gac caa acc ttt cac ctg atg	1405
Ala Gly Leu Thr Leu Pro Leu Asn Ser Asp Gln Thr Phe His Leu Met	
275 280 285	
agc aac ctg gct ggg gat gtt atc aca gct gca gtg act gca gct atc	1453
Ser Asn Leu Ala Gly Asp Val Ile Thr Ala Ala Val Thr Ala Ala Ile	
290 295 300	
aaa gac cag tta gag ggt gtg cag caa gca ctt tct cag gct gcc ccc	1501
Lys Asp Gln Leu Glu Gly Val Gln Gln Ala Leu Ser Gln Ala Ala Pro	
305 310 315	
atc cca gaa gag gac aca gac act gaa gaa ggt gat gac ttt gaa cta	1549
Ile Pro Glu Glu Asp Thr Asp Thr Glu Glu Gly Asp Asp Phe Glu Leu	
320 325 330 335	
ctt gac cag tca gag ctg gat caa att gag agt gaa ttg gga ctt aca	1597
Leu Asp Gln Ser Glu Leu Asp Gln Ile Glu Ser Glu Leu Gly Leu Thr	
340 345 350	
caa gac cag gaa gca gaa gca cag caa aat aag aag tct tca ggt ttc	1645
Gln Asp Gln Glu Ala Glu Ala Gln Gln Asn Lys Lys Ser Ser Gly Phe	
355 360 365	
ctt tca aat ctg ctg gga ggc cat taa tctag gaatcagctt gcaacagagc	1697
Leu Ser Asn Leu Leu Gly Gly His *	
370 375	
acaaaaaa	1705

ctg aca gtg aca cgg ctg tac ttc agc gcc gag gag ggg ggt gag cgc Leu Thr Val Thr Arg Leu Tyr Phe Ser Ala Glu Glu Gly Gly Glu Arg 155 160 165	654
tct gtc tgc ctc acc ttt gcc ttc ctc ttc ctg ctg ctg gcc atg ctg Ser Val Cys Leu Thr Phe Ala Phe Leu Phe Leu Leu Leu Ala Met Leu 170 175 180	702
gtg caa gtg gtg cgg gag gag acc ctc gag ctg ggc ctg gag cct ggt Val Gln Val Val Arg Glu Glu Thr Leu Glu Leu Gly Leu Glu Pro Gly 185 190 195	750
ctg gcc agc atg acc cag aac tta gag cca ctt ctg aag aag cag ggc Leu Ala Ser Met Thr Gln Asn Leu Glu Pro Leu Leu Lys Lys Gln Gly 200 205 210 215	798
tgg gac tgg gcg ctt cct gtg gcc aag ctg gct atc cgc gtg gga ctg Trp Asp Trp Ala Leu Pro Val Ala Lys Leu Ala Ile Arg Val Gly Leu 220 225 230	846
gca gtg gtg ggc tct gtg ctg ggt gcc ttc ctc acc ttc cca ggc ctg Ala Val Val Gly Ser Val Leu Gly Ala Phe Leu Thr Phe Pro Gly Leu 235 240 245	894
cgg ctg gcc cag acc cac cgg gac gca ctg acc atg tcg gag gac aga Arg Leu Ala Gln Thr His Arg Asp Ala Leu Thr Met Ser Glu Asp Arg 250 255 260	942
ccc atg ctg cag ttc ctc ctg cac acc agc ttc ctg tct ccc ctg ttc Pro Met Leu Gln Phe Leu Leu His Thr Ser Phe Leu Ser Pro Leu Phe 265 270 275	990
atc ctg tgg ctc tgg aca aag ccc att gca cgg gac ttc ctg cac cag Ile Leu Trp Leu Trp Thr Lys Pro Ile Ala Arg Asp Phe Leu His Gln 280 285 290 295	1038
ccg ccg ttt ggg gag acg cgt ttc tcc ctg ctg tcc gat tct gcc ttc Pro Pro Phe Gly Glu Thr Arg Phe Ser Leu Leu Ser Asp Ser Ala Phe 300 305 310	1086
gac tct ggg cgc ctc tgg ttg ctg gtg gtg ctg tgc ctg ctg cgg ctg Asp Ser Gly Arg Leu Trp Leu Leu Val Val Leu Cys Leu Leu Arg Leu 315 320 325	1134
gcg gtg acc cgg ccc cac ctg cag gcc tac ctg tgc ctg gcc aag gcc Ala Val Thr Arg Pro His Leu Gln Ala Tyr Leu Cys Leu Ala Lys Ala 330 335 340	1182
cgg gtg gag cag ctg cga agg gag gct ggc cgc atc gaa gcc cgt gaa Arg Val Glu Gln Leu Arg Arg Glu Ala Gly Arg Ile Glu Ala Arg Glu 345 350 355	1230
atc cag cag agg gtg gtc cga gtc tac tgc tat gtg acc gtg gtg agc Ile Gln Gln Arg Val Val Arg Val Tyr Cys Tyr Val Thr Val Val Ser 360 365 370 375	1278

ttg cag tac ctg acg ccg ctc atc ctc acc ctc aac tgc aca ctt ctg	1326
Leu Gln Tyr Leu Thr Pro Leu Ile Leu Thr Leu Asn Cys Thr Leu Leu	
380 385 390	
ctc aag acg ctg gga ggc tat tcc tgg ggc ctg ggc cca gct cct cta	1374
Leu Lys Thr Leu Gly Gly Tyr Ser Trp Gly Leu Gly Pro Ala Pro Leu	
395 400 405	
cta tcc ccc gac cca tcc tca gcc agc gct gcc ccc atc ggc tct ggg	1422
Leu Ser Pro Asp Pro Ser Ser Ala Ser Ala Ala Pro Ile Gly Ser Gly	
410 415 420	
gag gac gaa gtc cag cag act gca gcg cgg att gcc ggg gct ctg ggt	1470
Glu Asp Glu Val Gln Gln Thr Ala Ala Arg Ile Ala Gly Ala Leu Gly	
425 430 435	
ggc ctg ctt act ccc ctc ttc ctc cgt ggc gtc ctg gcc tac ctc atc	1518
Gly Leu Leu Thr Pro Leu Phe Leu Arg Gly Val Leu Ala Tyr Leu Ile	
440 445 450 455	
tgg tgg acg gct gcc tgc cag ctg ctc gcc agc ctt ttc ggc ctc tac	1566
Trp Trp Thr Ala Ala Cys Gln Leu Leu Ala Ser Leu Phe Gly Leu Tyr	
460 465 470	
ttc cac cag cac ttg gca ggc tcc tag ctgcc tgcagaccct cctggggccc	1618
Phe His Gln His Leu Ala Gly Ser *	
475 480	
tgaggtctgt tccctggggca gcgggacact agcctgcccc ctctgtttgc gccccctgt	1678
ccccagctgc aaggtggggc cggactcccc ggcgttcctc tcaccacagt gcctgacccg	1738
cggccccccct tggacgccga gtttctgcct cagaactgtc tctcctgggc ccagcagcat	1798
gaggggtcccg aggccattgt ctccgaagcg tatgtgccag gtttgagtgg cgaggggtgat	1858
gctggctgct cttctgaaca aataaaggag catgccgatt tttaaaaaa aaaaaa	1914

<210> 17

<211> 859

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (61) .. (795)

<400> 17

atttggccct cgaggccaag aattcggcac gaggaaggag gtccccacg gcccttcagg	60
------------------------------------------------------------------	----

atg aaa gct gcg gtg ctg acc ttg gcc gtg ctc ttc ctg acg ggg agc	108
Met Lys Ala Ala Val Leu Thr Leu Ala Val Leu Phe Leu Thr Gly Ser	
1 5 10 15	

cag gct cgg cat ttc tgg cag caa gat gaa ccc ccc cag agc ccc tgg	156
Gln Ala Arg His Phe Trp Gln Gln Asp Glu Pro Pro Gln Ser Pro Trp	
20 25 30	
gat cga gtg aag gac ctg gcc act gtg tac gtg gat gtg ctc aaa gac	204
Asp Arg Val Lys Asp Leu Ala Thr Val Tyr Val Asp Val Leu Lys Asp	
35 40 45	
agc ggc aaa gac agc gtg acc tcc acc ttc agc aag ctg cgc gaa cag	252
Ser Gly Lys Asp Ser Val Thr Ser Thr Phe Ser Lys Leu Arg Glu Gln	
50 55 60	
ctc ggc cct gtg acc cag gag ttc tgg gat aac ctg gaa aag gag aca	300
Leu Gly Pro Val Thr Gln Glu Phe Trp Asp Asn Leu Glu Lys Glu Thr	
65 70 75 80	
gag ggc ctg agg cag gag atg agc aag gat ctg gag gag gtg aag gcc	348
Glu Gly Leu Arg Gln Glu Met Ser Lys Asp Leu Glu Glu Val Lys Ala	
85 90 95	
aag gtg cag ccc tac ctg gac gac ttc cag aag aag tgg cag gag gag	396
Lys Val Gln Pro Tyr Leu Asp Asp Phe Gln Lys Lys Trp Gln Glu Glu	
100 105 110	
atg gag ctc tac cgc cag aag gtg gag ccg ctg cgc gca gag ctc caa	444
Met Glu Leu Tyr Arg Gln Lys Val Glu Pro Leu Arg Ala Glu Leu Gln	
115 120 125	
gag ggc gcg cgc cag aag ctg cac gag ctg caa gag aag ctg agc cca	492
Glu Gly Ala Arg Gln Lys Leu His Glu Leu Gln Glu Lys Leu Ser Pro	
130 135 140	
ctg ggc gag gag atg cgc gac cgc gcg cgc gcc cat gtg gac gcg ctg	540
Leu Gly Glu Glu Met Arg Asp Arg Ala Arg Ala His Val Asp Ala Leu	
145 150 155 160	
cgc acg cat ctg gcc ccc tac agc gac gag ctg cgc cag cgc ttg gcc	588
Arg Thr His Leu Ala Pro Tyr Ser Asp Glu Leu Arg Gln Arg Leu Ala	
165 170 175	
gcg cgc ctt gag gct ctc aag gag aac ggc ggc gcc aga ctg gcc gag	636
Ala Arg Leu Glu Ala Leu Lys Glu Asn Gly Gly Ala Arg Leu Ala Glu	
180 185 190	
tac cac gcc aag gcc acc gag cat ctg agc acg ctc agc gag aag gcc	684
Tyr His Ala Lys Ala Thr Glu His Leu Ser Thr Leu Ser Glu Lys Ala	
195 200 205	
aag ccc gcg ctc gag gac ctc cgc caa ggc ctg ctg ccc gtg ctg gag	732
Lys Pro Ala Leu Glu Asp Leu Arg Gln Gly Leu Leu Pro Val Leu Glu	
210 215 220	
agc ttc aag gtc agc ttc ctg agc gct ctc gag gag tac act aag aag	780
Ser Phe Lys Val Ser Phe Leu Ser Ala Leu Glu Tyr Thr Lys Lys	
225 230 235 240	
ctc aac acc cag tga ggcgccccgcc gccgcccccc ttccccggtgc tcagaataaa	835

Leu Asn Thr Gln *
245

cgtttccaaa gtgggaaaaa aaaa

859

<210> 18
<211> 2454
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (154)..(1710)

<400> 18

gtgaaaggag ggaacgcagg tgagaaagcg agacaggcag gtagggaaat cgtgaggtga 60

gcgtgatacct agctccttgt gccagagcct agagagaagg cgaggacgct gaagaaccag 120

gcggacagct gccagagaga gaagttggct agc atg gaa tca cca gag gag cct 174
Met Glu Ser Pro Glu Glu Pro
1 5

gga gca tcc atg gat gag aac tac ttt gtg aac tac act ttc aaa gat 222
Gly Ala Ser Met Asp Glu Asn Tyr Phe Val Asn Tyr Thr Phe Lys Asp
10 15 20

cgg tca cat tca ggc cgt gtg gct caa ggc atc atg aaa ctg tgt cta 270
Arg Ser His Ser Gly Arg Val Ala Gln Gly Ile Met Lys Leu Cys Leu
25 30 35

gag gag gag ctc ttt gct gat gtc acc att tcg gtg gaa ggc cgg gag 318
Glu Glu Glu Leu Phe Ala Asp Val Thr Ile Ser Val Glu Gly Arg Glu
40 45 50 55

ttt cag ctc cat cgg ctg gtc ctc tca gct cag agc tgc ttc ttc cga 366
Phe Gln Leu His Arg Leu Val Leu Ser Ala Gln Ser Cys Phe Phe Arg
60 65 70

tcc atg ttc act tcc aac ctg aag gag gcc cac aac cgg gtg att gtg 414
Ser Met Phe Thr Ser Asn Leu Lys Glu Ala His Asn Arg Val Ile Val
75 80 85

ctg cag gat gtc agc gag tct gtt ttc cag ctc ctg gtt gat tat atc 462
Leu Gln Asp Val Ser Glu Ser Val Phe Gln Leu Leu Val Asp Tyr Ile
90 95 100

tac cat ggg act gtg aaa ctt cga gct gag gag ttg cag gaa att tat 510
Tyr His Gly Thr Val Lys Leu Arg Ala Glu Glu Leu Gln Glu Ile Tyr
105 110 115

gag gtg tca gac atg tat cag ctg aca tct ctc ttt gag gaa tgc tct 558
Glu Val Ser Asp Met Tyr Gln Leu Thr Ser Leu Phe Glu Glu Cys Ser
120 125 130 135

cgg ttt ttg gcc cgc aca gtg caa gtg gga aac tgc ctt cag gtg atg	606
Arg Phe Leu Ala Arg Thr Val Gln Val Gly Asn Cys Leu Gln Val Met	
140 145 150	
tggttg gca gat cgg cac agt gat cct gag ctc tat acg gct gcc aag	654
Trp Leu Ala Asp Arg His Ser Asp Pro Glu Leu Tyr Thr Ala Ala Lys	
155 160 165	
cac tgt gcc aag acc cac ctg gcc cag ctg cag aat aca gag gaa ttt	702
His Cys Ala Lys Thr His Leu Ala Gln Leu Gln Asn Thr Glu Glu Phe	
170 175 180	
ctc cac ttg ccc cac cgc tta ctc aca gat atc atc tcg gat gga gtt	750
Leu His Leu Pro His Arg Leu Leu Thr Asp Ile Ile Ser Asp Gly Val	
185 190 195	
ccg tgt tct cag aac cca aca gag gca ata gaa gcc tgg atc aac ttt	798
Pro Cys Ser Gln Asn Pro Thr Glu Ala Ile Glu Ala Trp Ile Asn Phe	
200 205 210 215	
aat aaa gag gaa aga gag gct ttt gca gag tca ctc agg aca agc ttg	846
Asn Lys Glu Glu Arg Glu Ala Phe Ala Glu Ser Leu Arg Thr Ser Leu	
220 225 230	
aag gaa att ggg gag aat gtg cac att tac ctg att ggg aaa gag tca	894
Lys Glu Ile Gly Glu Asn Val His Ile Tyr Leu Ile Gly Lys Glu Ser	
235 240 245	
tct cgt acc cac tcg ttg gct gtg tcc ttg cac tgt gca gaa gat gac	942
Ser Arg Thr His Ser Leu Ala Val Ser Leu His Cys Ala Glu Asp Asp	
250 255 260	
tcc atc agt gta agt ggc caa aac agt ttg tgc cac cag atc act gcg	990
Ser Ile Ser Val Ser Gly Gln Asn Ser Leu Cys His Gln Ile Thr Ala	
265 270 275	
gcc tgc aag cat ggt gga gac ttg tat gtg gtg gga ggg tcc atc cca	1038
Ala Cys Lys His Gly Gly Asp Leu Tyr Val Val Gly Gly Ser Ile Pro	
280 285 290 295	
cgg cgc atg tgg aag tgc aac aat gcc acc gtt gac tgg gag tgg tgt	1086
Arg Arg Met Trp Lys Cys Asn Asn Ala Thr Val Asp Trp Glu Trp Cys	
300 305 310	
gct cct ttg cct cgg gac cgg ctc cag cac acc ctg gtg tct gtg ccc	1134
Ala Pro Leu Pro Arg Asp Arg Leu Gln His Thr Leu Val Ser Val Pro	
315 320 325	
ggg aaa gat gcc ata tat tca ctg ggt ggc aag aca ctg caa gat acc	1182
Gly Lys Asp Ala Ile Tyr Ser Leu Gly Gly Lys Thr Leu Gln Asp Thr	
330 335 340	
ctc tcc aac gca gtc att tat tat cgc gta ggt gat aat gtg tgg aca	1230
Leu Ser Asn Ala Val Ile Tyr Tyr Arg Val Gly Asp Asn Val Trp Thr	
345 350 355	

gag aca act cag cta gag gtg gct gtg tca ggg gct gct ggt gcc aac	1278
Glu Thr Thr Gln Leu Glu Val Ala Val Ser Gly Ala Ala Gly Ala Asn	
360 365 370 375	
ctc aac ggg atc atc tac tta cta ggg ggg gag gag aat gat ctg gac	1326
Leu Asn Gly Ile Ile Tyr Leu Leu Gly Gly Glu Glu Asn Asp Leu Asp	
380 385 390	
ttc ttt acc aaa cct tcc cga ctc atc cag tgc ttt gac aca gag aca	1374
Phe Phe Thr Lys Pro Ser Arg Leu Ile Gln Cys Phe Asp Thr Glu Thr	
395 400 405	
gac aaa tgc cat gtg aag ccc tat gtg ctg ccc ttt gca ggc cgc atg	1422
Asp Lys Cys His Val Lys Pro Tyr Val Leu Pro Phe Ala Gly Arg Met	
410 415 420	
cac gca gct gtg cat aaa gat ctg gtg ttc atc gtg gct gaa ggg gac	1470
His Ala Ala Val His Lys Asp Leu Val Phe Ile Val Ala Glu Gly Asp	
425 430 435	
tcc ctg gtg tgc tac aat ccc ttg cta gac agc ttc acc cgg ctt tgc	1518
Ser Leu Val Cys Tyr Asn Pro Leu Leu Asp Ser Phe Thr Arg Leu Cys	
440 445 450 455	
ctt cct gag gcc tgg agc tct gcc cca tcc ctc tgg aag att gcc agc	1566
Leu Pro Glu Ala Trp Ser Ser Ala Pro Ser Leu Trp Lys Ile Ala Ser	
460 465 470	
tgt aac ggg agc atc tat gtc ttc cgg gac cga tat aaa aag ggg gat	1614
Cys Asn Gly Ser Ile Tyr Val Phe Arg Asp Arg Tyr Lys Lys Gly Asp	
475 480 485	
gcc aac acc tac aag ctt gac cct gcc act tca gcc gta act gtc aca	1662
Ala Asn Thr Tyr Lys Leu Asp Pro Ala Thr Ser Ala Val Thr Val Thr	
490 495 500	
aga ggt att aag gtg ctg ctt acc aat ttg cag ttt gtg ttg gcc taa	1710
Arg Gly Ile Lys Val Leu Thr Asn Leu Gln Phe Val Leu Ala *	
505 510 515	
ggctgtgggg aggggaggag aactgctcac tccttttccc tccccataca aactcaaagt	1770
cccctggggcc ccaattcaga gttatgtttt ttttggcaca tactagaaag gcagtgcctc	1830
agcccttccc tgaatccatg gaggtgttct gtttggggct ttttagactg ctgctgctca	1890
gctggttgct tgaactgaca gtaggccagc ctgttctctg ccattcccta gtcctcctgt	1950
gcctcaccac agcttgctta gagcaagcct tttctcagac cttaggcaca gcctctcctc	2010
tttacctgat caatgttaaa tgtaagcacc cctgatccca ggacataagg aaagatgccc	2070
aattgtactt ttgttctata gcctgtgaaa tggctagttg atcatttttc cacaaagaat	2130
taggtgttaa gagttttcct tcaggcttta cttaggagaa tggactaagc tgaaggtgta	2190
cttcaccagc aagagtcaac tctagaattc aggatgttcc ttctattggt ttcttagcca	2250

```

tctgtcagga aatgtaactt tggtttttatt tttggcttat tccaaggggt aagccagaaa 2310
atagaaatga ttattttctga ttaatagcag aaactttttc aatctcaa atataaggtg 2370
tctgctcttt taaaagctct aagctaagtc aagagctagg aactgttgat acaaataaaa 2430
gtttttgaag ggaaaaaaaa aaaa 2454

```

```

<210> 19
<211> 3586
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (106)..(1860)

```

```

<400> 19
gtaccggtcc ggaattcccg ggtcgacgat ttcgtggagc aggcagacgt cgaagccgtc 60

ctgcagccga ccctcctgca tctgggcacg ttcagggttc gacac atg aag gct 114
Met Lys Ala
1

ttg atg aac gag aag gcc cag gcc gcc ctg gtg gag ttt gtg gag gac 162
Leu Met Asn Glu Lys Ala Gln Ala Ala Leu Val Glu Phe Val Glu Asp
5 10 15

gtc aat cac gct gcc att ccc agg gag atc cca cgc aag gat ggg gtc 210
Val Asn His Ala Ala Ile Pro Arg Glu Ile Pro Arg Lys Asp Gly Val
20 25 30 35

tgg agg gtt ctg tgg aag gac cgt gcg cag gac acg agg gtc ctg agg 258
Trp Arg Val Leu Trp Lys Asp Arg Ala Gln Asp Thr Arg Val Leu Arg
40 45 50

cag atg aca cgc ctg ctg ctg gat gac ggg ccc acg cag gcc gcg gag 306
Gln Met Thr Arg Leu Leu Leu Asp Asp Gly Pro Thr Gln Ala Ala Glu
55 60 65

gct ggg acc ccc ggt gag gca ccc acc cct ccc gct tcg gag acg cag 354
Ala Gly Thr Pro Gly Glu Ala Pro Thr Pro Pro Ala Ser Glu Thr Gln
70 75 80

gcc cag gat tct ggt gag gta aca ggg cat gct ggc tcg ctt ctt ggg 402
Ala Gln Asp Ser Gly Glu Val Thr Gly His Ala Gly Ser Leu Leu Gly
85 90 95

gca ccc agg aac cca agg agg ggc cgt cgg ggt cgc aga aac aga acc 450
Ala Pro Arg Asn Pro Arg Arg Gly Arg Arg Arg Arg Asn Arg Thr
100 105 110 115

aga cgc aac agg ttg acc cag aag ggc aag aag aga agc cga gga gga 498

```

Arg	Arg	Asn	Arg	Leu	Thr	Gln	Lys	Gly	Lys	Lys	Arg	Ser	Arg	Gly	Gly	
				120					125					130		
cgg	ccg	tct	gct	ccc	gcg	agg	agt	gag	gcc	gag	gac	tct	tcc	gac	gag	546
Arg	Pro	Ser	Ala	Pro	Ala	Arg	Ser	Glu	Ala	Glu	Asp	Ser	Ser	Asp	Glu	
			135					140					145			
agc	ctg	ggc	atc	gtg	atc	gag	gag	atc	gac	cag	ggc	gac	ctg	agc	gga	594
Ser	Leu	Gly	Ile	Val	Ile	Glu	Glu	Ile	Asp	Gln	Gly	Asp	Leu	Ser	Gly	
		150					155					160				
gaa	gag	gac	cag	agc	gcg	ctg	tac	gcc	acg	ctg	cag	gcc	gct	gcc	agg	642
Glu	Glu	Asp	Gln	Ser	Ala	Leu	Tyr	Ala	Thr	Leu	Gln	Ala	Ala	Ala	Arg	
		165				170					175					
gag	ctg	gtt	agg	cag	tgg	gcg	ccc	tgc	aac	tcc	gag	ggg	gaa	gaa	gac	690
Glu	Leu	Val	Arg	Gln	Trp	Ala	Pro	Cys	Asn	Ser	Glu	Gly	Glu	Glu	Asp	
180					185					190					195	
ggc	ccc	cg	gag	ttc	ttg	gct	ctg	gtc	acc	gtc	acc	gac	aaa	tcg	aag	738
Gly	Pro	Arg	Glu	Phe	Leu	Ala	Leu	Val	Thr	Val	Thr	Asp	Lys	Ser	Lys	
				200					205					210		
aaa	gaa	gag	gca	gag	aag	gag	cca	gct	ggg	gcc	gaa	tcc	atc	cgc	ttg	786
Lys	Glu	Glu	Ala	Glu	Lys	Glu	Pro	Ala	Gly	Ala	Glu	Ser	Ile	Arg	Leu	
			215					220					225			
aac	acc	aaa	gaa	gac	aaa	aat	ggc	gtc	ccc	gac	tta	gtg	gcc	ctg	ctg	834
Asn	Thr	Lys	Glu	Asp	Lys	Asn	Gly	Val	Pro	Asp	Leu	Val	Ala	Leu	Leu	
		230					235					240				
gct	gtg	aga	gac	acc	ccg	gac	gag	gag	ccg	gtg	gac	agc	gac	act	tcg	882
Ala	Val	Arg	Asp	Thr	Pro	Asp	Glu	Glu	Pro	Val	Asp	Ser	Asp	Thr	Ser	
		245				250					255					
gag	agc	gac	tcg	cag	gaa	agt	ggg	gac	caa	gaa	aca	gag	gag	ttg	gat	930
Glu	Ser	Asp	Ser	Gln	Glu	Ser	Gly	Asp	Gln	Glu	Thr	Glu	Glu	Leu	Asp	
260					265				270					275		
aat	cct	gag	ttc	gtg	gcc	att	gtg	gcc	tat	acc	gac	ccg	tcg	gac	ccc	978
Asn	Pro	Glu	Phe	Val	Ala	Ile	Val	Ala	Tyr	Thr	Asp	Pro	Ser	Asp	Pro	
			280						285					290		
tgg	gcc	cg	gag	gag	atg	ttg	aaa	atc	gct	tct	gtt	atc	gag	tcg	ctg	1026
Trp	Ala	Arg	Glu	Glu	Met	Leu	Lys	Ile	Ala	Ser	Val	Ile	Glu	Ser	Leu	
			295					300					305			
ggc	tgg	agc	gac	gag	aaa	gac	aag	cga	gac	ccc	ctc	cga	cag	gtc	ttg	1074
Gly	Trp	Ser	Asp	Glu	Lys	Asp	Lys	Arg	Asp	Pro	Leu	Arg	Gln	Val	Leu	
		310					315					320				
tcc	gtc	atg	tcc	aag	gac	act	aac	ggg	acc	cgc	gtg	aag	gtg	gaa	gag	1122
Ser	Val	Met	Ser	Lys	Asp	Thr	Asn	Gly	Thr	Arg	Val	Lys	Val	Glu	Glu	
		325				330					335					
gcg	ggc	cgc	gag	gtg	gac	gcc	gtg	gtc	ctg	cgc	aag	gcc	ggg	gat	gac	1170
Ala	Gly	Arg	Glu	Val	Asp	Ala	Val	Val	Leu	Arg	Lys	Ala	Gly	Asp	Asp	

340	345	350	355	
ggg gac ctc cgg gag tgc att tcc acc ttg gcg cag ccg gat ctc cct				1218
Gly Asp Leu Arg Glu Cys Ile Ser Thr Leu Ala Gln Pro Asp Leu Pro	360	365	370	
ccc cag gcg aag aag gct ggg cgt ggc ctc ttc ggg ggc tgg agc gag				1266
Pro Gln Ala Lys Lys Ala Gly Arg Gly Leu Phe Gly Gly Trp Ser Glu	375	380	385	
cac cgt gag gac gaa ggg ggt ctt ctg gag ctg gtg gcg ctc ctg gct				1314
His Arg Glu Asp Glu Gly Gly Leu Leu Glu Leu Val Ala Leu Leu Ala	390	395	400	
gcc cag gac atg gcg gag gtg atg aag gag gaa aaa gaa aac gcc tgg				1362
Ala Gln Asp Met Ala Glu Val Met Lys Glu Glu Lys Glu Asn Ala Trp	405	410	415	
gaa ggc ggg aag tac aaa tac ccc aaa ggc aaa ctg ggg gag gta ttg				1410
Glu Gly Gly Lys Tyr Lys Tyr Pro Lys Gly Lys Leu Gly Glu Val Leu	420	425	430	435
gcg ctc ctg gcc gcc cgg gag aac atg ggg tcc aac gag ggg tcg gag				1458
Ala Leu Leu Ala Ala Arg Glu Asn Met Gly Ser Asn Glu Gly Ser Glu	440	445	450	
gag gct tcg gac gaa cag tcc gag gag gag tcg gag gac acc gag agc				1506
Glu Ala Ser Asp Glu Gln Ser Glu Glu Glu Ser Glu Asp Thr Glu Ser	455	460	465	
gag gcg tcg gag ccg gag gac agg gca tcc agg aag ccc cgg gcc aag				1554
Glu Ala Ser Glu Pro Glu Asp Arg Ala Ser Arg Lys Pro Arg Ala Lys	470	475	480	
agg gcg cgc acg gcc ccc agg ggc ctg act ccg gcc ggc gcg cct ccc				1602
Arg Ala Arg Thr Ala Pro Arg Gly Leu Thr Pro Ala Gly Ala Pro Pro	485	490	495	
acc gct tcc ggg gcc cgc aaa acc cgc gcg ggc ggc cga ggc cga ggc				1650
Thr Ala Ser Gly Ala Arg Lys Thr Arg Ala Gly Gly Arg Gly Arg Gly	500	505	510	515
cga ggc cgg ggc gtc act ccc gag aag aaa gcc ggg agc cgg ggc tcg				1698
Arg Gly Arg Gly Val Thr Pro Glu Lys Lys Ala Gly Ser Arg Gly Ser	520	525	530	
gcc cag gac gac gcc gca gga agc agg aag aag aag ggg agc gcg ggc				1746
Ala Gln Asp Asp Ala Ala Gly Ser Arg Lys Lys Lys Gly Ser Ala Gly	535	540	545	
tcc ggg gcc cat gcc agg gca ggc gag gcc aag ggc cag gcg ccc act				1794
Ser Gly Ala His Ala Arg Ala Gly Glu Ala Lys Gly Gln Ala Pro Thr	550	555	560	
gga tcc aag gcc gcg cgc ggg aag aag gcc cgt cgg ggc cgg agg ctg				1842
Gly Ser Lys Ala Ala Arg Gly Lys Lys Ala Arg Arg Gly Arg Arg Leu	565	570	575	


```

gtggtgttgc ctgtgggtgg ggagggcccc aggtgtgctt ctcccgtagc agtcctaggc 3516
ttctctccct gtgccctgtg tcacctggat cctccagtaa agtgaaattc agcactgtaa 3576
aaaaaaaaaa 3586

```

```

<210> 20
<211> 3340
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (47)..(1894)

```

```

<400> 20
gatacagttc tgaccatcat tacgccaatc ttggcacgag gggaag atg gcg gag 55
Met Ala Glu
1

tcc ggc ggt agc agc ggt ggt gct ggt ggc ggc ggc gct ttc ggc gcg 103
Ser Gly Gly Ser Ser Gly Gly Ala Gly Gly Gly Ala Phe Gly Ala
5 10 15

ggc ccg ggc ccc gag cgc ccg aac agc acg gcc gac aag aac ggg gcc 151
Gly Pro Gly Pro Glu Arg Pro Asn Ser Thr Ala Asp Lys Asn Gly Ala
20 25 30 35

ctc aag tgc acc ttc tcg gca ccc agc cac agc acc agc ctc ctg cag 199
Leu Lys Cys Thr Phe Ser Ala Pro Ser His Ser Thr Ser Leu Leu Gln
40 45 50

ggc ctg gcc acc ctc cgc gct cag ggc cag ctc ctc gat gtt gtg ctg 247
Gly Leu Ala Thr Leu Arg Ala Gln Gly Gln Leu Leu Asp Val Val Leu
55 60 65

act att aac aga gag gcc ttt cct gca cac aag gtc gtc ctg gct gcc 295
Thr Ile Asn Arg Glu Ala Phe Pro Ala His Lys Val Val Leu Ala Ala
70 75 80

tgc agc gac tac ttc agg gcc atg ttc acc ggc ggc atg cgg gag gca 343
Cys Ser Asp Tyr Phe Arg Ala Met Phe Thr Gly Gly Met Arg Glu Ala
85 90 95

agc cag gac gtc atc gag ctg aag ggc gtg tcg gcc cgt ggc ctg cgg 391
Ser Gln Asp Val Ile Glu Leu Lys Gly Val Ser Ala Arg Gly Leu Arg
100 105 110 115

cac atc atc gac ttc gcc tac agc gcc gag gtg aca ctg gac ctg gac 439
His Ile Ile Asp Phe Ala Tyr Ser Ala Glu Val Thr Leu Asp Leu Asp
120 125 130

tgc gtg cag gac gtg ctg ggc gcg gcc gtg ttc ttg cag atg ctg ccc 487
Cys Val Gln Asp Val Leu Gly Ala Ala Val Phe Leu Gln Met Leu Pro

```


cag tac cgc agc ggc gag ggc gca gtg gac gcc tgc tac cgc tac gac Gln Tyr Arg Ser Gly Glu Gly Ala Val Asp Ala Cys Tyr Arg Tyr Asp 375 380 385	1207
ccc cac ctg aat cgc tgg ctg cgc ctg cag gcc atg cag gaa agc cgc Pro His Leu Asn Arg Trp Leu Arg Leu Gln Ala Met Gln Glu Ser Arg 390 395 400	1255
atc cag ttc cag ctg aac gtg ctg tgc ggc atg gtg tac gcc acg ggc Ile Gln Phe Gln Leu Asn Val Leu Cys Gly Met Val Tyr Ala Thr Gly 405 410 415	1303
ggc cgc aac cga gcc ggc agc ctg gcc tcc gtg gag cgg tac tgc ccc Gly Arg Asn Arg Ala Gly Ser Leu Ala Ser Val Glu Arg Tyr Cys Pro 420 425 430 435	1351
cgg cgc aat gag tgg ggc tac gcc tgc tgc ctg aag cgc cgt acc tgg Arg Arg Asn Glu Trp Gly Tyr Ala Cys Ser Leu Lys Arg Arg Thr Trp 440 445 450	1399
ggc cat gct ggg gcc gcc tca ggg ggc cgc ctc tac atc tgc ggt ggc Gly His Ala Gly Ala Ala Ser Gly Gly Arg Leu Tyr Ile Ser Gly Gly 455 460 465	1447
tac ggg atc tca gtg gag gac aag aag gcc ctg cac tgc tac gac ccc Tyr Gly Ile Ser Val Glu Asp Lys Lys Ala Leu His Cys Tyr Asp Pro 470 475 480	1495
gtg gcc gac cag tgg gag ttc aag gcg ccc atg agc gaa ccc cgc gtg Val Ala Asp Gln Trp Glu Phe Lys Ala Pro Met Ser Glu Pro Arg Val 485 490 495	1543
cta cac gcc atg gtg ggt gcc ggc ggc cgc atc tat gcc ctc ggg ggc Leu His Ala Met Val Gly Ala Gly Gly Arg Ile Tyr Ala Leu Gly Gly 500 505 510 515	1591
cgc atg gac cac gcg gac cgc tgc ttc gac gtg ctg gct gtg gag tac Arg Met Asp His Ala Asp Arg Cys Phe Asp Val Leu Ala Val Glu Tyr 520 525 530	1639
tat gtg ccg gag acg gac cag tgg acc agc gtg agc ccc atg cgg gcc Tyr Val Pro Glu Thr Asp Gln Trp Thr Ser Val Ser Pro Met Arg Ala 535 540 545	1687
ggc cag tca gag gcc ggc tgc tgc ctg ctg gag agg aag atc tac atc Gly Gln Ser Glu Ala Gly Cys Cys Leu Leu Glu Arg Lys Ile Tyr Ile 550 555 560	1735
gtc ggg ggc tac aac tgg cgt ctc aac aac gtc acg ggc atc gta cag Val Gly Gly Tyr Asn Trp Arg Leu Asn Asn Val Thr Gly Ile Val Gln 565 570 575	1783
gtg tac aac acg gac acc gac gag tgg gag cgg gac ctg cac ttc ccg Val Tyr Asn Thr Asp Thr Asp Glu Trp Glu Arg Asp Leu His Phe Pro 580 585 590 595	1831

gag tcc ttc gca ggc ata gcc tgc gcc ccc gtc ctg ctg ccc cgg gcc	1879
Glu Ser Phe Ala Gly Ile Ala Cys Ala Pro Val Leu Leu Pro Arg Ala	
600 605 610	
ggg acc agg agg tag cccccaagac ccccgaggacc ctggcctgac cgcattgtgt	1934
Gly Thr Arg Arg *	
615	
ctccaagtgg ggcttggcga atgcacgtct gcctgagaac cccagtgcc cccttcgccc	1994
gggctgccct tgaggggcct gctgcgttga taagcccccc tcccaggggt ccctccctcc	2054
ctccttccca aagcagatcc tggctgcgag tccatccgag ggagcctgcc ggcaaagcgt	2114
ctgacatgtg gtggcagcaa attcgtcccc ggggtggttt cctcgcttgg ccccgagtc	2174
cccacgggct ggcgggtgga atcccaggtc tccagggggt ccctgtgcag ctccatctca	2234
cttctctact gcctcccagc cccacgggtt caggcattca gatgtgagct catcaacatt	2294
gaacccaaag tcgggtggtat atgactcacc ctccttccaa gtctcctgcg cgcgtgtttt	2354
taaaataaac tcacccgaaa cgtccgtaac acacggacct ccaggagcag tgagaggtgg	2414
cttagaagcc cctggtttgg ggtgggtgga ggaggagggc acgtgtctgc ctcccctggg	2474
gtgccctcct tcccccatcc caagctgctg aggggaggcc ctggctcatgc ctcagttcct	2534
gccttcattc gctttccgga ggaaaaacca tatcaactcc tagaaacgct ccttaggggc	2594
ttgggacctt ccatttggca ctgagcatct tgtggggcct taactggctg agacatcccg	2654
gccctctaca tttgccctgt tggccaggca gtccccttcc cgcaattgga gggcgacgct	2714
aacttcagaa tcccatagtg gcgcttgccg caggctctggt ggggtgtctt ttttctctc	2774
cctcctttcc acccctccgc gccctgccac tcctggcct gccctgtttt tgggtcaaca	2834
ttgctacgga gccagcagtg aggcctttcc cttcaagggc tctgtggtat ctctggccac	2894
atttgttcta atgtctgaac ctctaaatct tttctttttg attggtttta ctgttttta	2954
gaagccagca ctgctgtctc atagatggga tttgtactct tggggcaact taaagtgtct	3014
ctctcgctgc taacagacga ttgatgtctt gtctctgtga cccactcacc atgtaaagaa	3074
ttaacctcct atcttagcag acatcgctc ctaatatattt cctttattta ataaaaatgt	3134
tatggtgaag agatggagcc ggcccagcac tgagcttggt cggttggtgt ctgattggtc	3194
acagattcct cgtgtgtcct ccgcgtgtct gggggctcct ctccccgcc tcagcctccc	3254
cccgatccc cccaagaaag gaaaattatt tttcgtattg taaactttta acatgaaaaa	3314
gctgttttta atttaaaaaa aaaaaa	3340

<210> 21
 <211> 2547
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (143)..(2275)

<400> 21

```

aaggatcctt aattaaatta atcccccccc cccccccgcc cccgcggtgg cggcggcggc      60

ggcgggtggct gccgtggcgg ctgagagtcc agagccggac gttccggccg cttegggctg      120

gcggctggag agcgctcggg tc atg tct gcc cag ggg gac tgc gag ttc ctg      172
                Met Ser Ala Gln Gly Asp Cys Glu Phe Leu
                  1             5             10

gtg cag cga gcc cgg gag ttg gtg ccg caa gac ctg tgg gca gcc aag      220
Val Gln Arg Ala Arg Glu Leu Val Pro Gln Asp Leu Trp Ala Ala Lys
                  15             20             25

gcg tgg ctg atc acg gcc cgc agc ctc tac ccg gca gac ttt aac atc      268
Ala Trp Leu Ile Thr Ala Arg Ser Leu Tyr Pro Ala Asp Phe Asn Ile
                  30             35             40

cag tat gag atg tac acc atc gag cgg aat gca gag cgg acc gcc acc      316
Gln Tyr Glu Met Tyr Thr Ile Glu Arg Asn Ala Glu Arg Thr Ala Thr
                  45             50             55

gcc ggg agg ctg ctg tac gac atg ttt gtg aat ttc cca gac cag ccg      364
Ala Gly Arg Leu Leu Tyr Asp Met Phe Val Asn Phe Pro Asp Gln Pro
                  60             65             70

gtg gtg tgg aga gaa atc agc att att aca tca gca tta agg aac gat      412
Val Val Trp Arg Glu Ile Ser Ile Ile Thr Ser Ala Leu Arg Asn Asp
                  75             80             85             90

tca cag gac aaa caa acc caa ttt tta aga agt tta ttt gaa act ctt      460
Ser Gln Asp Lys Gln Thr Gln Phe Leu Arg Ser Leu Phe Glu Thr Leu
                  95             100             105

cct ggt cgg gtc cag tgt gaa atg tta cta aag gtc acg gaa caa tgc      508
Pro Gly Arg Val Gln Cys Glu Met Leu Leu Lys Val Thr Glu Gln Cys
                  110             115             120

ttc aac acg tta gaa cga tca gaa atg ttg ctt cta ctt ttg agg cgc      556
Phe Asn Thr Leu Glu Arg Ser Glu Met Leu Leu Leu Leu Leu Arg Arg
                  125             130             135

ttc cct gaa acg gtg gtg cag cat ggg gtt ggc ctt ggg gag gca cta      604
Phe Pro Glu Thr Val Val Gln His Gly Val Gly Leu Gly Glu Ala Leu
                  140             145             150

tta gag gct gaa act att gaa gaa caa gaa tct cca gtg aac tgc ttt      652

```

Leu	Glu	Ala	Glu	Thr	Ile	Glu	Glu	Gln	Glu	Ser	Pro	Val	Asn	Cys	Phe	
155					160					165					170	
aga	aaa	tta	ttt	gtt	tgt	gat	gtc	ctt	cct	cta	ata	att	aac	aac	cat	700
Arg	Lys	Leu	Phe	Val	Cys	Asp	Val	Leu	Pro	Leu	Ile	Ile	Asn	Asn	His	
				175					180					185		
gat	gtt	cga	tta	cct	gcc	aat	tta	ttg	tat	aag	tac	ttg	aac	aaa	gca	748
Asp	Val	Arg	Leu	Pro	Ala	Asn	Leu	Leu	Tyr	Lys	Tyr	Leu	Asn	Lys	Ala	
			190					195					200			
gct	gaa	ttt	tat	atc	aat	tat	gtc	act	agg	tct	act	caa	ata	gaa	aat	796
Ala	Glu	Phe	Tyr	Ile	Asn	Tyr	Val	Thr	Arg	Ser	Thr	Gln	Ile	Glu	Asn	
		205					210					215				
cag	cat	caa	ggc	gcc	cag	gat	aca	tct	gat	tta	atg	tca	cct	agc	aaa	844
Gln	His	Gln	Gly	Ala	Gln	Asp	Thr	Ser	Asp	Leu	Met	Ser	Pro	Ser	Lys	
		220				225					230					
cgt	agc	tct	cag	aag	tac	ata	ata	gaa	ggg	ctg	acg	gaa	aaa	tca	tcc	892
Arg	Ser	Ser	Gln	Lys	Tyr	Ile	Ile	Glu	Gly	Leu	Thr	Glu	Lys	Ser	Ser	
					240					245					250	
cag	atc	gtg	gac	cct	tgg	gag	agg	ttg	ttt	aag	att	ttg	aat	gtt	gtt	940
Gln	Ile	Val	Asp	Pro	Trp	Glu	Arg	Leu	Phe	Lys	Ile	Leu	Asn	Val	Val	
				255					260					265		
gga	atg	aga	tgt	gaa	tgg	cag	atg	gat	aaa	gga	aga	cga	agc	tat	gga	988
Gly	Met	Arg	Cys	Glu	Trp	Gln	Met	Asp	Lys	Gly	Arg	Arg	Ser	Tyr	Gly	
			270					275					280			
gat	att	ttg	cat	aga	atg	aag	gat	ctc	tgc	aga	tac	atg	aac	aac	ttt	1036
Asp	Ile	Leu	His	Arg	Met	Lys	Asp	Leu	Cys	Arg	Tyr	Met	Asn	Asn	Phe	
		285					290					295				
gat	agt	gaa	gca	cat	gca	aaa	tat	aaa	aac	caa	gtg	gtg	tat	tcc	acc	1084
Asp	Ser	Glu	Ala	His	Ala	Lys	Tyr	Lys	Asn	Gln	Val	Val	Tyr	Ser	Thr	
		300				305					310					
atg	ctg	gtc	ttc	ttt	aag	aat	gca	ttc	cag	tat	gtc	aac	agc	ata	cag	1132
Met	Leu	Val	Phe	Phe	Lys	Asn	Ala	Phe	Gln	Tyr	Val	Asn	Ser	Ile	Gln	
		315				320				325					330	
cca	tct	ctc	ttc	caa	ggc	cct	aat	gcc	ccg	agc	caa	gtt	cca	ctg	gtt	1180
Pro	Ser	Leu	Phe	Gln	Gly	Pro	Asn	Ala	Pro	Ser	Gln	Val	Pro	Leu	Val	
				335					340					345		
ctt	ctt	gaa	gat	gta	tcg	aat	gtg	tat	ggc	gat	gta	gaa	att	gat	cgt	1228
Leu	Leu	Glu	Asp	Val	Ser	Asn	Val	Tyr	Gly	Asp	Val	Glu	Ile	Asp	Arg	
			350					355					360			
aat	aaa	cac	atc	cat	aaa	aag	agg	aaa	cta	gct	gaa	gga	aga	gaa	aaa	1276
Asn	Lys	His	Ile	His	Lys	Lys	Arg	Lys	Leu	Ala	Glu	Gly	Arg	Glu	Lys	
		365					370					375				
acc	atg	agt	tca	gac	gat	gaa	gac	tgt	tcg	gcg	aaa	gga	aga	aat	cgt	1324
Thr	Met	Ser	Ser	Asp	Asp	Glu	Asp	Cys	Ser	Ala	Lys	Gly	Arg	Asn	Arg	

380	385	390	
cac att gta gtc aat aaa gcc gaa ctt gct aac tcc act gaa gtg tta			1372
His Ile Val Val Asn Lys Ala Glu Leu Ala Asn Ser Thr Glu Val Leu			
395	400	405	410
gaa agc ttt aaa ttg gcc agg gag agc tgg gag ttg ctc tat tcc cta			1420
Glu Ser Phe Lys Leu Ala Arg Glu Ser Trp Glu Leu Leu Tyr Ser Leu			
	415	420	425
gaa ttc ctt gac aaa gaa ttt aca agg att tgc ttg gcc tgg aag acg			1468
Glu Phe Leu Asp Lys Glu Phe Thr Arg Ile Cys Leu Ala Trp Lys Thr			
	430	435	440
gat act tgg ctt tgg tta aga atc ttc ctc act gat atg atc atc tat			1516
Asp Thr Trp Leu Trp Leu Arg Ile Phe Leu Thr Asp Met Ile Ile Tyr			
	445	450	455
cag ggt caa tat aaa aag gcg ata gcc agc ctg cat cac tta gca gct			1564
Gln Gly Gln Tyr Lys Lys Ala Ile Ala Ser Leu His His Leu Ala Ala			
	460	465	470
ctc cag gga tcc att tct cag cca cag atc aca ggg cag ggg acc ctg			1612
Leu Gln Gly Ser Ile Ser Gln Pro Gln Ile Thr Gly Gln Gly Thr Leu			
	475	480	490
gag cat cag agg gcg ctc atc cag ctg gcg acg tgc cac ttt gcg cta			1660
Glu His Gln Arg Ala Leu Ile Gln Leu Ala Thr Cys His Phe Ala Leu			
	495	500	505
ggg gag tac aga atg aca tgt gaa aaa gtc ctt gat ttg atg tgc tac			1708
Gly Glu Tyr Arg Met Thr Cys Glu Lys Val Leu Asp Leu Met Cys Tyr			
	510	515	520
atg gta ctc ccc att caa gat gga ggc aaa tcc cag gag gaa ccc tcg			1756
Met Val Leu Pro Ile Gln Asp Gly Gly Lys Ser Gln Glu Glu Pro Ser			
	525	530	535
aaa gta aag ccc aaa ttt aga aaa ggt tcg gat ctg aag ctc ctg cct			1804
Lys Val Lys Pro Lys Phe Arg Lys Gly Ser Asp Leu Lys Leu Leu Pro			
	540	545	550
tgt acc agc aag gct atc atg cca tac tgc ctc cat tta atg tta gcc			1852
Cys Thr Ser Lys Ala Ile Met Pro Tyr Cys Leu His Leu Met Leu Ala			
	555	560	570
tgt ttt aag ctt aga gct ttc aca gac aac aga gac gac atg gca ttg			1900
Cys Phe Lys Leu Arg Ala Phe Thr Asp Asn Arg Asp Asp Met Ala Leu			
	575	580	585
ggg cat gtg att gtg ttg ctt cag caa gag tgg cca cgg ggc gag aat			1948
Gly His Val Ile Val Leu Leu Gln Gln Glu Trp Pro Arg Gly Glu Asn			
	590	595	600
ctt ttc ctg aaa gct gtc aat aaa att tgc caa caa gga aat ttc caa			1996
Leu Phe Leu Lys Ala Val Asn Lys Ile Cys Gln Gln Gly Asn Phe Gln			
	605	610	615

tat gag aat ttt ttc aat tac gtt aca aat att gat atg ctg gag gaa	2044
Tyr Glu Asn Phe Phe Asn Tyr Val Thr Asn Ile Asp Met Leu Glu Glu	
620 625 630	
ttt gcc tac ttg aga act cag gaa ggt ggg aaa att cat ctg gaa tta	2092
Phe Ala Tyr Leu Arg Thr Gln Glu Gly Gly Lys Ile His Leu Glu Leu	
635 640 645 650	
cta ccc aat caa gga atg ctg atc aag cac cac act gta act cga ggc	2140
Leu Pro Asn Gln Gly Met Leu Ile Lys His His Thr Val Thr Arg Gly	
655 660 665	
atc acc aaa ggc gtg aag gag gac ttt cgc ctg gcc atg gag cgc cag	2188
Ile Thr Lys Gly Val Lys Glu Asp Phe Arg Leu Ala Met Glu Arg Gln	
670 675 680	
gtc tcc cgc tgt gga gag aat ctg atg gtg gtt ctg cac agg ttc tgc	2236
Val Ser Arg Cys Gly Glu Asn Leu Met Val Val Leu His Arg Phe Cys	
685 690 695	
att aat gag aag atc ttg ctc ctt cag act ctg acc tga gtggagacct	2285
Ile Asn Glu Lys Ile Leu Leu Leu Gln Thr Leu Thr *	
700 705 710	
ttccaccaga cacagctcgg gcctgtgtaa ttgtaggaga agacactcag cagtgattgc	2345
catggcacag agccgtggtc attgttgctg ttacaaagaa gaaaaccatc tgagttctaa	2405
ctccttgggtt gcttaaaagt agttcccaag agtctgagaa gctattttcta tttttaagag	2465
tcatttttttg taattttttgt aaaacaaaag taccaatctg ttttgtaaataaaaaatcatc	2525
ctaaaatttg aaaaaaaaaa aa	2547

<210> 22
 <211> 2625
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (143)..(2353)

aaggatcctt aattaaatta atcccccccc cccccccgcc cccgcggtgg cggcggcgcc	60
ggcgggtgget gccgtggcgg ctgagagtcc agagccggac gttccggccg cttcgggctg	120
gcggctggag agcgcctcggg tc atg tct gcc cag ggg gac tgc gag ttc ctg	172
Met Ser Ala Gln Gly Asp Cys Glu Phe Leu	
1 5 10	
gtg cag cga gcc cgg gag ttg gtg ccg caa gac ctg tgg gca gcc aag	220

Val	Gln	Arg	Ala	Arg	Glu	Leu	Val	Pro	Gln	Asp	Leu	Trp	Ala	Ala	Lys		
				15					20					25			
gcg	tgg	ctg	atc	acg	gcc	cgc	agc	ctc	tac	ccg	gca	gac	ttt	aac	atc	268	
Ala	Trp	Leu	Ile	Thr	Ala	Arg	Ser	Leu	Tyr	Pro	Ala	Asp	Phe	Asn	Ile		
		30						35					40				
cag	tat	gag	atg	tac	acc	atc	gag	cgg	aat	gca	gag	cgg	acc	gcc	acc	316	
Gln	Tyr	Glu	Met	Tyr	Thr	Ile	Glu	Arg	Asn	Ala	Glu	Arg	Thr	Ala	Thr		
		45					50					55					
gcc	ggg	agg	ctg	ctg	tac	gac	atg	ttt	gtg	aat	ttc	cca	gac	cag	ccg	364	
Ala	Gly	Arg	Leu	Leu	Tyr	Asp	Met	Phe	Val	Asn	Phe	Pro	Asp	Gln	Pro		
	60					65					70						
gtg	gtg	tgg	aga	gaa	atc	agc	att	att	aca	tca	gca	tta	agg	aac	gat	412	
Val	Val	Trp	Arg	Glu	Ile	Ser	Ile	Ile	Thr	Ser	Ala	Leu	Arg	Asn	Asp		
	75				80					85					90		
tca	cag	gac	aaa	caa	acc	caa	ttt	tta	aga	agt	tta	ttt	gaa	act	ctt	460	
Ser	Gln	Asp	Lys	Gln	Thr	Gln	Phe	Leu	Arg	Ser	Leu	Phe	Glu	Thr	Leu		
				95				100						105			
cct	ggt	cgg	gtc	cag	tgt	gaa	atg	tta	cta	aag	gtc	acg	gaa	caa	tgc	508	
Pro	Gly	Arg	Val	Gln	Cys	Glu	Met	Leu	Leu	Lys	Val	Thr	Glu	Gln	Cys		
			110					115					120				
ttc	aac	acg	tta	gaa	cga	tca	gaa	atg	ttg	ctt	cta	ctt	ttg	agg	cgc	556	
Phe	Asn	Thr	Leu	Glu	Arg	Ser	Glu	Met	Leu	Leu	Leu	Leu	Leu	Arg	Arg		
		125					130					135					
ttc	cct	gaa	acg	gtg	gtg	cag	cat	ggg	gtt	ggc	ctt	ggg	gag	gca	cta	604	
Phe	Pro	Glu	Thr	Val	Val	Gln	His	Gly	Val	Gly	Leu	Gly	Glu	Ala	Leu		
	140					145					150						
tta	gag	gct	gaa	act	att	gaa	gaa	caa	gaa	tct	cca	gtg	aac	tgc	ttt	652	
Leu	Glu	Ala	Glu	Thr	Ile	Glu	Glu	Gln	Glu	Ser	Pro	Val	Asn	Cys	Phe		
	155				160					165					170		
aga	aaa	tta	ttt	gtt	tgt	gat	gtc	ctt	cct	cta	ata	att	aac	aac	cat	700	
Arg	Lys	Leu	Phe	Val	Cys	Asp	Val	Leu	Pro	Leu	Ile	Ile	Asn	Asn	His		
				175					180					185			
gat	gtt	cga	tta	cct	gcc	aat	tta	ttg	tat	aag	tac	ttg	aac	aaa	gca	748	
Asp	Val	Arg	Leu	Pro	Ala	Asn	Leu	Leu	Tyr	Lys	Tyr	Leu	Asn	Lys	Ala		
			190					195					200				
gct	gaa	ttt	tat	atc	aat	tat	gtc	act	agg	tct	act	caa	ata	gaa	aat	796	
Ala	Glu	Phe	Tyr	Ile	Asn	Tyr	Val	Thr	Arg	Ser	Thr	Gln	Ile	Glu	Asn		
		205					210					215					
cag	cat	caa	ggc	gcc	cag	gat	aca	tct	gat	tta	atg	tca	cct	agc	aaa	844	
Gln	His	Gln	Gly	Ala	Gln	Asp	Thr	Ser	Asp	Leu	Met	Ser	Pro	Ser	Lys		
	220					225					230						
cgt	agc	tct	cag	aag	tac	ata	ata	gaa	ggg	ctg	acg	gaa	aaa	tca	tcc	892	
Arg	Ser	Ser	Gln	Lys	Tyr	Ile	Ile	Glu	Gly	Leu	Thr	Glu	Lys	Ser	Ser		

235	240	245	250	
cag atc gtg gac cct tgg gag agg ttg ttt aag att ttg aat gtt gtt				940
Gln Ile Val Asp 255	Pro Trp Glu Arg Leu Phe Lys Ile Leu Asn Val Val 265			
gga atg aga tgt gaa tgg cag atg gat aaa gga aga cga agc tat gga				988
Gly Met Arg Cys 270	Glu Trp Gln Met Asp Lys Gly Arg Arg Ser Tyr Gly 280			
gat att ttg cat aga atg aag gat ctc tgc aga tac atg aac aac ttt				1036
Asp Ile Leu His Arg Met Lys Asp Leu Cys Arg Tyr Met Asn Asn Phe 285				
gat agt gaa gca cat gca aaa tat aaa aac caa gtg gtg tat tcc acc				1084
Asp Ser Glu Ala His Ala Lys Tyr Lys Asn Gln Val Val Tyr Ser Thr 300				
atg ctg gtc ttc ttt aag aat gca ttc cag tat gtc aac agc ata cag				1132
Met Leu Val Phe Phe Lys Asn Ala Phe Gln Tyr Val Asn Ser Ile Gln 315				
cca tct ctc ttc caa ggt cct aat gcc ccg agc caa gtt cca ctg gtt				1180
Pro Ser Leu Phe Gln Gly Pro Asn Ala Pro Ser Gln Val Pro Leu Val 335				
ctt ctt gaa gat gta tgc aat gtg tat ggt gat gta gaa att gat cgt				1228
Leu Leu Glu Asp Val Ser Asn Val Tyr Gly Asp Val Glu Ile Asp Arg 350				
aat aaa cac atc cat aaa aag agg aaa cta gct gaa gga aga gaa aaa				1276
Asn Lys His Ile His Lys Lys Arg Lys Leu Ala Glu Gly Arg Glu Lys 365				
acc atg agt tca gac gat gaa gac tgt tgc gcg aaa gga aga aat cgt				1324
Thr Met Ser Ser Asp Asp Glu Asp Cys Ser Ala Lys Gly Arg Asn Arg 380				
cac att gta gtc aat aaa gcc gaa ctt gct aac tcc act gaa gtg tta				1372
His Ile Val Val Asn Lys Ala Glu Leu Ala Asn Ser Thr Glu Val Leu 395				
gaa agc ttt aaa ttg gcc agg gag agc tgg gag ttg ctc tat tcc cta				1420
Glu Ser Phe Lys Leu Ala Arg Glu Ser Trp Glu Leu Leu Tyr Ser Leu 415				
gaa ttc ctt gac aaa gaa ttt aca agg att tgc ttg gcc tgg aag acg				1468
Glu Phe Leu Asp Lys Glu Phe Thr Arg Ile Cys Leu Ala Trp Lys Thr 430				
gat act tgg ctt tgg tta aga atc ttc ctc act gat atg atc atc tat				1516
Asp Thr Trp Leu Trp Leu Arg Ile Phe Leu Thr Asp Met Ile Ile Tyr 445				
cag ggt caa tat aaa aag gcg ata gcc agc ctg cat cac tta gca gct				1564
Gln Gly Gln Tyr Lys Lys Ala Ile Ala Ser Leu His His Leu Ala Ala 460				

ctc	cag	gga	tcc	att	tct	cag	cca	cag	atc	aca	ggg	cag	ggg	acc	ctg	1612
Leu	Gln	Gly	Ser	Ile	Ser	Gln	Pro	Gln	Ile	Thr	Gly	Gln	Gly	Thr	Leu	
475					480					485					490	
gag	cat	cag	agg	gcg	ctc	atc	cag	ctg	gcg	acg	tgc	cac	ttt	gcg	cta	1660
Glu	His	Gln	Arg	Ala	Leu	Ile	Gln	Leu	Ala	Thr	Cys	His	Phe	Ala	Leu	
			495						500					505		
ggg	gag	tac	aga	atg	aca	tgt	gaa	aaa	gtc	ctt	gat	ttg	atg	tgc	tac	1708
Gly	Glu	Tyr	Arg	Met	Thr	Cys	Glu	Lys	Val	Leu	Asp	Leu	Met	Cys	Tyr	
			510					515					520			
atg	gta	ctc	ccc	att	caa	gat	gga	ggc	aaa	tcc	cag	gag	gaa	ccc	tcg	1756
Met	Val	Leu	Pro	Ile	Gln	Asp	Gly	Gly	Lys	Ser	Gln	Glu	Glu	Pro	Ser	
			525				530						535			
aaa	gta	aag	ccc	aaa	ttt	aga	aaa	ggg	tcg	gat	ctg	aag	ctc	ctg	cct	1804
Lys	Val	Lys	Pro	Lys	Phe	Arg	Lys	Gly	Ser	Asp	Leu	Lys	Leu	Leu	Pro	
			540			545					550					
tgt	acc	agc	aag	gct	atc	atg	cca	tac	tgc	ctc	cat	tta	atg	tta	gcc	1852
Cys	Thr	Ser	Lys	Ala	Ile	Met	Pro	Tyr	Cys	Leu	His	Leu	Met	Leu	Ala	
555					560				565						570	
tgt	ttt	aag	ctt	aga	gct	ttc	aca	gac	aac	aga	gac	gac	atg	gca	ttg	1900
Cys	Phe	Lys	Leu	Arg	Ala	Phe	Thr	Asp	Asn	Arg	Asp	Asp	Met	Ala	Leu	
				575					580					585		
ggg	cat	gtg	att	gtg	ttg	ctt	cag	caa	gag	tgg	cca	cgg	ggc	gag	aat	1948
Gly	His	Val	Ile	Val	Leu	Leu	Gln	Gln	Glu	Trp	Pro	Arg	Gly	Glu	Asn	
				590				595					600			
ctt	ttc	ctg	aaa	gct	gtc	aat	aaa	att	tgc	caa	caa	gga	aat	ttc	caa	1996
Leu	Phe	Leu	Lys	Ala	Val	Asn	Lys	Ile	Cys	Gln	Gln	Gly	Asn	Phe	Gln	
			605				610					615				
tat	gag	aat	ttt	ttc	aat	tac	gtt	aca	aat	att	gat	atg	ctg	gag	gaa	2044
Tyr	Glu	Asn	Phe	Phe	Asn	Tyr	Val	Thr	Asn	Ile	Asp	Met	Leu	Glu	Glu	
			620			625					630					
ttt	gcc	tac	ttg	aga	act	cag	gaa	ggg	ggg	aaa	att	cat	ctg	gaa	tta	2092
Phe	Ala	Tyr	Leu	Arg	Thr	Gln	Glu	Gly	Gly	Lys	Ile	His	Leu	Glu	Leu	
635					640					645					650	
cta	ccc	aat	caa	gga	atg	ctg	atc	aag	cct	tct	agc	cct	ccc	atg	ggg	2140
Leu	Pro	Asn	Gln	Gly	Met	Leu	Ile	Lys	Pro	Ser	Ser	Pro	Pro	Met	Gly	
				655					660					665		
tta	ctg	cag	cag	gaa	ttc	tta	cct	gtg	ctt	cag	ccc	agc	ata	cag	act	2188
Leu	Leu	Gln	Gln	Glu	Phe	Leu	Pro	Val	Leu	Gln	Pro	Ser	Ile	Gln	Thr	
				670				675					680			
gct	gac	agg	cac	cac	act	gta	act	cga	ggc	atc	acc	aaa	ggc	gtg	aag	2236
Ala	Asp	Arg	His	His	Thr	Val	Thr	Arg	Gly	Ile	Thr	Lys	Gly	Val	Lys	
			685				690					695				

gag gac ttt cgc ctg gcc atg gag cgc cag gtc tcc cgc tgt gga gag 2284
 Glu Asp Phe Arg Leu Ala Met Glu Arg Gln Val Ser Arg Cys Gly Glu
 700 705 710

aat ctg atg gtg gtt ctg cac agg ttc tgc att aat gag aag atc ttg 2332
 Asn Leu Met Val Val Leu His Arg Phe Cys Ile Asn Glu Lys Ile Leu
 715 720 725 730

ctc ctt cag act ctg acc tga gt ggagaccttt ccaccagaca cagctcgggc 2385
 Leu Leu Gln Thr Leu Thr *
 735

ctgtgtaatt gtaggagaag acactcagca gtgattgccca tggcacagag ccgtgggtcat 2445

tgttgctgtt acaaagaaga aaaccatctg agttctaact ccttggttgc ttaaaagtag 2505

ttcccaagag tctgagaagc tatttctatt tttaagagtc attttttgta atttttgtaa 2565

aacaaaagta ccaatctgtt ttgtaaataa aaatcatcct aaaatttgaa aaaaaaaaaa 2625

<210> 23

<211> 6288

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (115) .. (5253)

<400> 23

aaggagagccc cgctcagcgc ggggagcgcc cggccccctc cccgccccat gcgcccgcgg 60

ctctgaagcc tgagcggggc cgggggctgg gcggggccgg ggcccgcctg aggc atg 117
 Met
 1

gcg tcc ggg agc cgg tgg cgg ccg aca ccg ccg ccg ctg ctg ttg ctg 165
 Ala Ser Gly Ser Arg Trp Arg Pro Thr Pro Pro Pro Leu Leu Leu Leu
 5 10 15

ctg ctg ctg gcg ctg gcg gcg cgc gcg gac ggc ctg gag ttc ggc ggc 213
 Leu Leu Leu Ala Leu Ala Ala Arg Ala Asp Gly Leu Glu Phe Gly Gly
 20 25 30

ggc ccc ggg cag tgg gct cgc tac gcg cgc tgg gcg ggc gcg gcg agc 261
 Gly Pro Gly Gln Trp Ala Arg Tyr Ala Arg Trp Ala Gly Ala Ala Ser
 35 40 45

agc ggc gag ctc agc ttc agc ctg cgc acc aac gcc acg cgc gcg ctg 309
 Ser Gly Glu Leu Ser Phe Ser Leu Arg Thr Asn Ala Thr Arg Ala Leu
 50 55 60 65

ctg ctc tac ctg gac gac ggc ggc gac tgc gac ttc ctg gag ctg ctg 357
 Leu Leu Tyr Leu Asp Asp Gly Gly Asp Cys Asp Phe Leu Glu Leu Leu

	70	75	80	
ctg gtg gac ggc cgc ctg cgg ctg cgc ttc acg ctt tcg tgc gcc gag				405
Leu Val Asp Gly Arg Leu Arg Leu Arg Phe Thr Leu Ser Cys Ala Glu				
	85	90	95	
ccg gcc acg ctg cag ctg gac acg ccg gtg gcc gac gac cgc tgg cac				453
Pro Ala Thr Leu Gln Leu Asp Thr Pro Val Ala Asp Asp Arg Trp His				
	100	105	110	
atg gtg ctg ctg acc cgc gac gcg cgc cgc acg gcg ctg gcg gtg gac				501
Met Val Leu Leu Thr Arg Asp Ala Arg Arg Thr Ala Leu Ala Val Asp				
	115	120	125	
ggc gag gcc cgc gcc gcc gag gtg cgc tcc aag cgg cgc gag atg cag				549
Gly Glu Ala Arg Ala Ala Glu Val Arg Ser Lys Arg Arg Glu Met Gln				
	130	135	140	145
gtg gcc agc gac ctg ttc gtg ggc ggc atc ccg ccc gac gtg cgc ctc				597
Val Ala Ser Asp Leu Phe Val Gly Gly Ile Pro Pro Asp Val Arg Leu				
	150	155	160	
tcg gcg ctt acg ctg agc acc gtc aag tac gag ccg ccc ttc cgc ggt				645
Ser Ala Leu Thr Leu Ser Thr Val Lys Tyr Glu Pro Pro Phe Arg Gly				
	165	170	175	
ctc ttg gcc aac ctg aag ctg ggc gag cgg ccc ccc gcg ctg ctg ggc				693
Leu Leu Ala Asn Leu Lys Leu Gly Glu Arg Pro Pro Ala Leu Leu Gly				
	180	185	190	
agc cag ggc ctg cgc ggc gcc acc gcc gac ccg ctg tgc gcg ccc gcg				741
Ser Gln Gly Leu Arg Gly Ala Thr Ala Asp Pro Leu Cys Ala Pro Ala				
	195	200	205	
cgc aac ccc tgc gcc aac ggc ggc ctc tgc acc gtg ctg gcc ccc ggc				789
Arg Asn Pro Cys Ala Asn Gly Gly Leu Cys Thr Val Leu Ala Pro Gly				
	210	215	220	225
gag gtg ggc tgc gac tgc agc cac acg ggc ttc ggc ggc aag ttc tgc				837
Glu Val Gly Cys Asp Cys Ser His Thr Gly Phe Gly Gly Lys Phe Cys				
	230	235	240	
agc gaa gag gag cac ccc atg gaa ggt ccg gct cac ctg acg tta aac				885
Ser Glu Glu Glu His Pro Met Glu Gly Pro Ala His Leu Thr Leu Asn				
	245	250	255	
agc gaa gta ggg tcc tta ctg ttc tcc gag ggg ggg gcc ggg aga gga				933
Ser Glu Val Gly Ser Leu Leu Phe Ser Glu Gly Gly Ala Gly Arg Gly				
	260	265	270	
gga gcc ggc gat gtg cac cag cca aca aaa ggc aag gag gag ttt gtg				981
Gly Ala Gly Asp Val His Gln Pro Thr Lys Gly Lys Glu Glu Phe Val				
	275	280	285	
gcg acc ttc aaa ggc aat gag ttc ttc tgc tac gac ctg tca cac aac				1029
Ala Thr Phe Lys Gly Asn Glu Phe Phe Cys Tyr Asp Leu Ser His Asn				
	290	295	300	305

ccc atc cag agc agc act gat gag atc aca ctg gcc ttc cgc acc ctg	1077
Pro Ile Gln Ser Ser Thr Asp Glu Ile Thr Leu Ala Phe Arg Thr Leu	
310 315 320	
caa cgc aac ggc ctg atg ctg cat aca ggc aag tcg gcc gac tac gtc	1125
Gln Arg Asn Gly Leu Met Leu His Thr Gly Lys Ser Ala Asp Tyr Val	
325 330 335	
aac ctg tcc ctc aag tct ggg gct gtc tgg ctg gtc atc aac cta ggc	1173
Asn Leu Ser Leu Lys Ser Gly Ala Val Trp Leu Val Ile Asn Leu Gly	
340 345 350	
tca ggt gcc ttc gag gcc ctt gtg gaa ccc gtc aat ggc aag ttc aac	1221
Ser Gly Ala Phe Glu Ala Leu Val Glu Pro Val Asn Gly Lys Phe Asn	
355 360 365	
gac aac gcc tgg cac gac gtc cgg gtc acc cga aac ctg cgc cag cac	1269
Asp Asn Ala Trp His Asp Val Arg Val Thr Arg Asn Leu Arg Gln His	
370 375 380 385	
gca ggg att gga cac gct atg gta aac aaa ctg cat tat ctg gtg acc	1317
Ala Gly Ile Gly His Ala Met Val Asn Lys Leu His Tyr Leu Val Thr	
390 395 400	
atc tcg gtg gac ggg atc ctg acc acc aca ggc tac acg cag gag gat	1365
Ile Ser Val Asp Gly Ile Leu Thr Thr Thr Gly Tyr Thr Gln Glu Asp	
405 410 415	
tac acc atg ctg ggc tct gat gac ttc ttc tac att ggg ggc agc ccc	1413
Tyr Thr Met Leu Gly Ser Asp Asp Phe Phe Tyr Ile Gly Gly Ser Pro	
420 425 430	
aac aca gct gac ctg ccg ggc tcg ccc gtc agc aac aac ttc atg ggc	1461
Asn Thr Ala Asp Leu Pro Gly Ser Pro Val Ser Asn Asn Phe Met Gly	
435 440 445	
tgc ctc aag gac gtg gtc tat aag aac aat gac ttc aaa ttg gaa cta	1509
Cys Leu Lys Asp Val Val Tyr Lys Asn Asn Asp Phe Lys Leu Glu Leu	
450 455 460 465	
tcc cgc ctg gca aag gaa ggg gac ccc aag atg aag ctg cag ggg gac	1557
Ser Arg Leu Ala Lys Glu Gly Asp Pro Lys Met Lys Leu Gln Gly Asp	
470 475 480	
ttg tca ttc cgc tgt gag gat gtg gct gcc ctg gac cct gtg acc ttt	1605
Leu Ser Phe Arg Cys Glu Asp Val Ala Ala Leu Asp Pro Val Thr Phe	
485 490 495	
gag agt ccc gag gcc ttt gtg gcg ctg ccc cgc tgg agc gct aag cgc	1653
Glu Ser Pro Glu Ala Phe Val Ala Leu Pro Arg Trp Ser Ala Lys Arg	
500 505 510	
act ggc tcc atc tcc cta gac ttc cgc acc acc gag ccc aat ggg ctg	1701
Thr Gly Ser Ile Ser Leu Asp Phe Arg Thr Thr Glu Pro Asn Gly Leu	
515 520 525	

ctg ctc ttc agc cag ggc cgg cgg gct ggg ggt gga gct ggc agc cac	1749
Leu Leu Phe Ser Gln Gly Arg Arg Ala Gly Gly Gly Ala Gly Ser His	
530 535 540 545	
agc tct gct cag cgg gcc gac tac ttt gcc atg gag cta ttg gac ggc	1797
Ser Ser Ala Gln Arg Ala Asp Tyr Phe Ala Met Glu Leu Leu Asp Gly	
550 555 560	
cac ctc tat ctt ctg ctg gac atg gga tct ggg ggc atc aag ctg cgg	1845
His Leu Tyr Leu Leu Leu Asp Met Gly Ser Gly Gly Ile Lys Leu Arg	
565 570 575	
gca tcc agc cgc aag gtc aat gat ggc gag tgg tgt cac gtg gac ttc	1893
Ala Ser Ser Arg Lys Val Asn Asp Gly Glu Trp Cys His Val Asp Phe	
580 585 590	
cag agg gat ggg cga aaa ggc tcc atc tca gtg aat agt cgc agc acg	1941
Gln Arg Asp Gly Arg Lys Gly Ser Ile Ser Val Asn Ser Arg Ser Thr	
595 600 605	
ccg ttc ttg gcc act gga gac agc gag att ctg gac ctg gag agt gag	1989
Pro Phe Leu Ala Thr Gly Asp Ser Glu Ile Leu Asp Leu Glu Ser Glu	
610 615 620 625	
ctg tac ctg ggc ggt ctc cct gag ggg ggc cgg gtg gac ctg ccc ctg	2037
Leu Tyr Leu Gly Gly Leu Pro Glu Gly Gly Arg Val Asp Leu Pro Leu	
630 635 640	
ccc cca gag gtg tgg aca gca gca ctc cgg gca ggc tac gtg ggc tgt	2085
Pro Pro Glu Val Trp Thr Ala Ala Leu Arg Ala Gly Tyr Val Gly Cys	
645 650 655	
gtg cgg gac ctc ttc ata gat ggg cgt agc cga gac ctc cgg ggc ctg	2133
Val Arg Asp Leu Phe Ile Asp Gly Arg Ser Arg Asp Leu Arg Gly Leu	
660 665 670	
gct gag gct cag ggg gct gtg ggc gtt gcc ccc ttt tgc tcc cgg gag	2181
Ala Glu Ala Gln Gly Ala Val Gly Val Ala Pro Phe Cys Ser Arg Glu	
675 680 685	
acg ctg aag cag tgt gca tct gcc ccc tgt cgc aat ggg ggc gtc tgt	2229
Thr Leu Lys Gln Cys Ala Ser Ala Pro Cys Arg Asn Gly Gly Val Cys	
690 695 700 705	
cga gaa ggc tgg aac cgc ttc atc tgt gac tgc atc ggg acc ggc ttt	2277
Arg Glu Gly Trp Asn Arg Phe Ile Cys Asp Cys Ile Gly Thr Gly Phe	
710 715 720	
ctt ggg cgg gtc tgt gag aga gag gcc acg gtc ctg agc tac gat ggc	2325
Leu Gly Arg Val Cys Glu Arg Glu Ala Thr Val Leu Ser Tyr Asp Gly	
725 730 735	
tcc atg tac atg aag atc atg ctg cct aac gcc atg cac acg gag gca	2373
Ser Met Tyr Met Lys Ile Met Leu Pro Asn Ala Met His Thr Glu Ala	
740 745 750	
gag gat gtg tcc ctg cgt ttc atg tcc cag cgg gcc tac gga ctc atg	2421

Glu Asp Val Ser Leu Arg Phe Met Ser Gln Arg Ala Tyr Gly Leu Met	
755 760 765	
atg gcc acc act tcc agg gag tct gcc gac acc cta cgc ctg gag ctg	2469
Met Ala Thr Thr Ser Arg Glu Ser Ala Asp Thr Leu Arg Leu Glu Leu	
770 775 780 785	
gat ggg ggg cag atg aag ctc act gtc aac ctc gac tgc ctg cgc gtc	2517
Asp Gly Gly Gln Met Lys Leu Thr Val Asn Leu Asp Cys Leu Arg Val	
790 795 800	
ggc tgc gca ccc agt aaa ggc ccc gaa acg ctg ttt gcg ggg cac aag	2565
Gly Cys Ala Pro Ser Lys Gly Pro Glu Thr Leu Phe Ala Gly His Lys	
805 810 815	
ctc aat gac aat gag tgg cac acg gtg agg gtg gtc cgg cgt ggc aag	2613
Leu Asn Asp Asn Glu Trp His Thr Val Arg Val Val Arg Arg Gly Lys	
820 825 830	
agc ctg cag ctg tct gtg gac aac gtg act gtg gag gga cag atg gca	2661
Ser Leu Gln Leu Ser Val Asp Asn Val Thr Val Glu Gly Gln Met Ala	
835 840 845	
gga gcc cat atg cgg ctg gag ttc cac aac att gag acg ggc atc atg	2709
Gly Ala His Met Arg Leu Glu Phe His Asn Ile Glu Thr Gly Ile Met	
850 855 860 865	
acg gag cgg cgg ttt atc tcc gtg gtg ccc tcc aac ttc atc ggg cat	2757
Thr Glu Arg Arg Phe Ile Ser Val Val Pro Ser Asn Phe Ile Gly His	
870 875 880	
ctg agt ggg ctc gtg ttc aat ggc cag ccc tac atg gac cag tgc aag	2805
Leu Ser Gly Leu Val Phe Asn Gly Gln Pro Tyr Met Asp Gln Cys Lys	
885 890 895	
gat ggt gac atc acc tac tgt gag ctc aat gct cgc ttt ggc ctg cgt	2853
Asp Gly Asp Ile Thr Tyr Cys Glu Leu Asn Ala Arg Phe Gly Leu Arg	
900 905 910	
gcc att gtg gcc gat ccc gtc acc ttc aag agt cgc agc agc tac ctg	2901
Ala Ile Val Ala Asp Pro Val Thr Phe Lys Ser Arg Ser Ser Tyr Leu	
915 920 925	
gca ctc gcc acg ctc caa gcc tat gct tcc atg cac ctc ttc ttc cag	2949
Ala Leu Ala Thr Leu Gln Ala Tyr Ala Ser Met His Leu Phe Phe Gln	
930 935 940 945	
ttc aag acc acg gcc cct gat ggg ctt ctt ctg ttc aac tcg ggc aac	2997
Phe Lys Thr Thr Ala Pro Asp Gly Leu Leu Leu Phe Asn Ser Gly Asn	
950 955 960	
ggc aat gac ttc att gtc atc gag ctg gtc aag ggg tac atc cac tac	3045
Gly Asn Asp Phe Ile Val Ile Glu Leu Val Lys Gly Tyr Ile His Tyr	
965 970 975	
gtg ttt gac ctg ggg aat ggc ccg tcc ttg atg aag ggg aac tca gac	3093
Val Phe Asp Leu Gly Asn Gly Pro Ser Leu Met Lys Gly Asn Ser Asp	

980	985	990	
aaa cca gtc aat gac aac cag tgg cac aac gtg gtg gtg tcc agg gac			3141
Lys Pro Val Asn Asp Asn Gln Trp His Asn Val Val Ser Arg Asp			
995	1000	1005	
cca ggc aac gtg cac acg ctc aag att gac tcc cgc act gtc acg cag			3189
Pro Gly Asn Val His Thr Leu Lys Ile Asp Ser Arg Thr Val Thr Gln			
1010	1015	1020	1025
cac tcc aat ggc gcc cga aac ctc gat ctc aaa ggg gag ttg tac att			3237
His Ser Asn Gly Ala Arg Asn Leu Asp Leu Lys Gly Glu Leu Tyr Ile			
	1030	1035	1040
ggc ggt ctg agc aag aat atg ttc agc aac ctg ccc aag ctg gtg gcc			3285
Gly Gly Leu Ser Lys Asn Met Phe Ser Asn Leu Pro Lys Leu Val Ala			
	1045	1050	1055
tcc cgg gat ggc ttt cag ggc tgc ctg gcc tca gtg gac ctc aac gga			3333
Ser Arg Asp Gly Phe Gln Gly Cys Leu Ala Ser Val Asp Leu Asn Gly			
	1060	1065	1070
cgt ctc cca gac ctc atc gcc gac gcc ctg cac cgc att ggg cag gtg			3381
Arg Leu Pro Asp Leu Ile Ala Asp Ala Leu His Arg Ile Gly Gln Val			
	1075	1080	1085
gag agg ggc tgt gat ggc ccc agc acc acc tgc act gaa gag tcc tgt			3429
Glu Arg Gly Cys Asp Gly Pro Ser Thr Thr Cys Thr Glu Glu Ser Cys			
1090	1095	1100	1105
gcc aac cag ggc gtc tgc ttg cag cag tgg gat ggc ttc acc tgc gac			3477
Ala Asn Gln Gly Val Cys Leu Gln Gln Trp Asp Gly Phe Thr Cys Asp			
	1110	1115	1120
tgc acc atg act tcc tat gga ggc cct gtc tgc aat gat ccc ggg acc			3525
Cys Thr Met Thr Ser Tyr Gly Gly Pro Val Cys Asn Asp Pro Gly Thr			
	1125	1130	1135
aca tac atc ttt ggg aag ggg gga gcg ctc atc acc tac acg tgg ccc			3573
Thr Tyr Ile Phe Gly Lys Gly Gly Ala Leu Ile Thr Tyr Thr Trp Pro			
	1140	1145	1150
ccc aat gac agg ccc agc acg agg atg gat cgc ctg gcc gtg ggc ttc			3621
Pro Asn Asp Arg Pro Ser Thr Arg Met Asp Arg Leu Ala Val Gly Phe			
	1155	1160	1165
agc acc cac cag cgg agc gct gtg ctg gtg cgg gtg gac agc gcc tcc			3669
Ser Thr His Gln Arg Ser Ala Val Leu Val Arg Val Asp Ser Ala Ser			
1170	1175	1180	1185
ggc ctt gga gac tac ctg cag ctg cac atc gac cag ggc acc gtg ggg			3717
Gly Leu Gly Asp Tyr Leu Gln Leu His Ile Asp Gln Gly Thr Val Gly			
	1190	1195	1200
gtg atc ttt aac gtg ggc acg gac gac att acc atc gac gag ccc aac			3765
Val Ile Phe Asn Val Gly Thr Asp Asp Ile Thr Ile Asp Glu Pro Asn			
	1205	1210	1215

gcc ata gta agc gac ggc aaa tac cac gtg gtg cgc ttc act cga agc	3813
Ala Ile Val Ser Asp Gly Lys Tyr His Val Val Arg Phe Thr Arg Ser	
1220 1225 1230	
ggc ggc aac gcc acc ctg cag gtg gac agc tgg ccg gtc aac gag cgg	3861
Gly Gly Asn Ala Thr Leu Gln Val Asp Ser Trp Pro Val Asn Glu Arg	
1235 1240 1245	
tac ccg gca gga aac ttt gat aac gag cgc ctg gcg att gct aga cag	3909
Tyr Pro Ala Gly Asn Phe Asp Asn Glu Arg Leu Ala Ile Ala Arg Gln	
1250 1255 1260 1265	
aga atc ccc tac ccg ctt ggt cga gta gta gat gag tgg ctg ctc gac	3957
Arg Ile Pro Tyr Arg Leu Gly Arg Val Val Asp Glu Trp Leu Leu Asp	
1270 1275 1280	
aaa ggc cgc cag ctg acc atc ttc aac agc cag gct gcc atc aag atc	4005
Lys Gly Arg Gln Leu Thr Ile Phe Asn Ser Gln Ala Ala Ile Lys Ile	
1285 1290 1295	
ggg ggc ccg gat cag ggc cgc ccc ttc cag ggc cag gtg tcc ggc ctc	4053
Gly Gly Arg Asp Gln Gly Arg Pro Phe Gln Gly Gln Val Ser Gly Leu	
1300 1305 1310	
tac tac aat ggg ctc aag gtg ctg gcg ctg gcc gcc gag agc gac ccc	4101
Tyr Tyr Asn Gly Leu Lys Val Leu Ala Leu Ala Ala Glu Ser Asp Pro	
1315 1320 1325	
aat gtg ccg act gag ggt cac ctg cgc ctg gtg ggg gag ggg ccg tcc	4149
Asn Val Arg Thr Glu Gly His Leu Arg Leu Val Gly Glu Gly Pro Ser	
1330 1335 1340 1345	
gtg ctg ctc agt gcg gag acc acg gcc acc acc ctg ctg gct gac atg	4197
Val Leu Leu Ser Ala Glu Thr Thr Ala Thr Thr Leu Leu Ala Asp Met	
1350 1355 1360	
gcc acc acc atc atg gag act acc acc acc atg gcc act acc acc acg	4245
Ala Thr Thr Ile Met Glu Thr Thr Thr Met Ala Thr Thr Thr Thr	
1365 1370 1375	
cgc ccg ggc cgc tcc ccc aca ctg agg gac agc acc acc cag aac aca	4293
Arg Arg Gly Arg Ser Pro Thr Leu Arg Asp Ser Thr Thr Gln Asn Thr	
1380 1385 1390	
gat gac ctg ctg gtg gcc tct gct gag tgt cca agc gat gat gag gac	4341
Asp Asp Leu Leu Val Ala Ser Ala Glu Cys Pro Ser Asp Asp Glu Asp	
1395 1400 1405	
ctg gag gag tgt gag ccc agt act gga gga gag tta ata ttg ccc att	4389
Leu Glu Glu Cys Glu Pro Ser Thr Gly Gly Glu Leu Ile Leu Pro Ile	
1410 1415 1420 1425	
atc acg gag gac tcc tta gac ccc cct ccc gtg gcc acc cga tcc ccc	4437
Ile Thr Glu Asp Ser Leu Asp Pro Pro Val Ala Thr Arg Ser Pro	
1430 1435 1440	

ttc gtg ccc ccg ccc cct acc ttc tac ccc ttc ctc acg gga gtg ggc	4485
Phe Val Pro Pro Pro Pro Thr Phe Tyr Pro Phe Leu Thr Gly Val Gly	
1445 1450 1455	
gcc acc caa gac acg ctg ccc ccg ccc gcc gcg cgc cgc ccg ccc tct	4533
Ala Thr Gln Asp Thr Leu Pro Pro Pro Ala Ala Arg Arg Pro Pro Ser	
1460 1465 1470	
ggg gcc ccg tgc cag gcc gag cgg gac gac agc gac tgc gag gag ccc	4581
Gly Gly Pro Cys Gln Ala Glu Arg Asp Asp Ser Asp Cys Glu Glu Pro	
1475 1480 1485	
atc gag gcc tcg ggc ttc gcc tcc ggg gag gtc ttt gac tcc agc ctc	4629
Ile Glu Ala Ser Gly Phe Ala Ser Gly Glu Val Phe Asp Ser Ser Leu	
1490 1495 1500 1505	
ccc ccc acg gac gac gag gac ttt tac acc acc ttt ccc ctg gtc acg	4677
Pro Pro Thr Asp Asp Glu Asp Phe Tyr Thr Thr Phe Pro Leu Val Thr	
1510 1515 1520	
gac cgc acc acc ctc ctg tca ccc cgc aaa ccc gct ccc cgg ccc aac	4725
Asp Arg Thr Thr Leu Leu Ser Pro Arg Lys Pro Ala Pro Arg Pro Asn	
1525 1530 1535	
ctc agg aca gat ggg gcc acg ggc gcc cct ggg gtg ctg ttt gcc ccc	4773
Leu Arg Thr Asp Gly Ala Thr Gly Ala Pro Gly Val Leu Phe Ala Pro	
1540 1545 1550	
tcc gcc ccg gcc ccc aac ctg ccg gcg gga aaa atg aac cac cga gac	4821
Ser Ala Pro Ala Pro Asn Leu Pro Ala Gly Lys Met Asn His Arg Asp	
1555 1560 1565	
ccg ctt cag ccc ttg ctg gag aac ccg ccc ttg ggg ccc ggg gcc ccc	4869
Pro Leu Gln Pro Leu Leu Glu Asn Pro Pro Leu Gly Pro Gly Ala Pro	
1570 1575 1580 1585	
acg tcc ttt gag ccg ccg agg ccc cct ccc ctg cgc ccc ggc gtg acc	4917
Thr Ser Phe Glu Pro Arg Arg Pro Pro Pro Leu Arg Pro Gly Val Thr	
1590 1595 1600	
tca gcc ccc ggc ttc ccc cat ctg ccc aca gcc aac ccc aca ggg cct	4965
Ser Ala Pro Gly Phe Pro His Leu Pro Thr Ala Asn Pro Thr Gly Pro	
1605 1610 1615	
ggg gag cgg gcc ccg ccg ggc gca gtg gag gtg atc cgg gag tcc agc	5013
Gly Glu Arg Gly Pro Pro Gly Ala Val Glu Val Ile Arg Glu Ser Ser	
1620 1625 1630	
agc acc acg ggc atg gtg gtg ggc att gtg gcg gcg gcg gcg ctc tgc	5061
Ser Thr Thr Gly Met Val Val Gly Ile Val Ala Ala Ala Ala Leu Cys	
1635 1640 1645	
atc ctc atc ctc ctc tac gcc atg tat aag tac cgc aat cgt gat gag	5109
Ile Leu Ile Leu Leu Tyr Ala Met Tyr Lys Tyr Arg Asn Arg Asp Glu	
1650 1655 1660 1665	
ggc tcc tac cag gtg gac cag agc cga aac tac atc agt aac tcg gcc	5157

Gly Ser Tyr Gln Val Asp Gln Ser Arg Asn Tyr Ile Ser Asn Ser Ala	
1670 1675 1680	
cag agc aat ggg gcg gtg gtg aaa gag aag gcc ccg gct gcc ccc aag	5205
Gln Ser Asn Gly Ala Val Val Lys Glu Lys Ala Pro Ala Ala Pro Lys	
1685 1690 1695	
acg ccc agc aag gcc aag aag aac aaa gac aag gag tat tat gtc tga	5253
Thr Pro Ser Lys Ala Lys Lys Asn Lys Asp Lys Glu Tyr Tyr Val *	
1700 1705 1710	
gcccccgcca ctgcgccccca ctgccagctg cccctcctgg gagggcccg gagggagggtg	5313
ccaccctctc cctgccaggg gcctggggac cctctccctg gctgcctcag gcttctctta	5373
cgaagaggaa acgcaaaaaa agaaaaggaa aaaccccggtg ctgcgcccct tctcctgcc	5433
gtccactgcg cggcctcgtc agtcccgggg ctgactgtcc ctctcagctc tgcgcctgcc	5493
aggcagggca cgtgctcaca gccctgggtt gatttatattt tttaaggggg tagttttatt	5553
ttggtgggggt tgggtgggaa ggaaggctgg gggttttgta aagtgtccac tgctcgtcct	5613
gttaattttc ctcaattttt cttcttcttc cttctgtccc tctgccttc cttctcttcc	5673
caagccctcc aatccccatc ccaggcttgc tgtgtctcac tgtccccacc ctccttccct	5733
acttcttttt ttgtgtgtct gggtttctccc ttcttttctt ccttttgggt ttccagagtc	5793
gggtgggagaa gggcgggagg gtggggccga gtggcccagt ggggtgggtgg ggtgggggtgg	5853
ggcaagtgcc ccaactcccc tcaccaggag aggcacctgc ttggtgccgc ccagggaagg	5913
ggctcaggcc tgacggaagg cctgttctgt gtgtgccgcc gggcgacgtg cattgatggg	5973
gaagctgctg gaggagcagg ggtgggggggt gggagggagg ggaaaggcaa atgcagatat	6033
atattacaga caaatactct agattccacg agcagcagcc tgtggcacc cgtgggcgcg	6093
ggcagcaggg aagagggagc aaggcattgt ccacagactg ctggggtcac ttctttgccc	6153
acgggctccc tgctccccca gttttttttc tctctttgtt aacaaatgtg tctgagtctt	6213
ggaaaacacc ccaaccccg aaatgtgtgg gaaaaagaaa acaaaaaactt tccaaattcc	6273
aaaaaaaaa aaaaa	6288

<210> 24
 <211> 1016
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS

<222> (254) .. (724)

<400> 24

ttaatac	gat	gactataggg	aatttg	gccc	tcgagcaaga	attcggcacg	aggctggctc	60
tgactcttca	cccatcttca	cccaggctgg	ccccctttggt	gaaactacaa	ctcccagggg	120		
tctgtgcgcg	agaaggcagg	cggtgttttc	taccggaagt	ccgctctagc	tctgggacct	180		
acaactgcac	cctgagccgg	agctgcccag	tcgccgcggg	accggggccg	ctgggggtctg	240		
gacggggggtc	gcc	atg atc cgc ttt atc ctc atc cag aac cgg gca ggc	289					
		Met Ile Arg Phe Ile Leu Ile Gln Asn Arg Ala Gly						
		1 5 10						
aag acg cgc ctg gcc aag tgg tac atg cag ttt gat gat gat gag aaa	337							
Lys Thr Arg Leu Ala Lys Trp Tyr Met Gln Phe Asp Asp Asp Glu Lys								
	15 20 25							
cag aag ctg atc gag gag gtg cat gcc gtg gtc acc gtc cga gac gcc	385							
Gln Lys Leu Ile Glu Glu Val His Ala Val Val Thr Val Arg Asp Ala								
	30 35 40							
aaa cac acc aac ttt gtg gag gtc ctg gca agc tcc gtt gct gac agc	433							
Lys His Thr Asn Phe Val Glu Val Leu Ala Ser Ser Val Ala Asp Ser								
	45 50 55 60							
ctc tct gtt ctg cag ttc cgg aac ttt aag atc att tac cgc cgc tat	481							
Leu Ser Val Leu Gln Phe Arg Asn Phe Lys Ile Ile Tyr Arg Arg Tyr								
	65 70 75							
gct ggc ctc tac ttc tgc atc tgt gtg gat gtc aat gac aac aac ctg	529							
Ala Gly Leu Tyr Phe Cys Ile Cys Val Asp Val Asn Asp Asn Asn Leu								
	80 85 90							
gct tac ctg gag gcc att cac aac ttc gtg gag gtc tta aac gaa tat	577							
Ala Tyr Leu Glu Ala Ile His Asn Phe Val Glu Val Leu Asn Glu Tyr								
	95 100 105							
ttc cac aat gtc tgt gaa ctg gac ctg gtg ttc aac ttc tac aag gtt	625							
Phe His Asn Val Cys Glu Leu Asp Leu Val Phe Asn Phe Tyr Lys Val								
	110 115 120							
tac acg gtc gtg gac gag atg ttc ctg gct ggc gaa atc cga gag acc	673							
Tyr Thr Val Val Asp Glu Met Phe Leu Ala Gly Glu Ile Arg Glu Thr								
	125 130 135 140							
agc cag acg aag gtg ctg aaa cag ctg ctg atg cta cag tcc ctg gag	721							
Ser Gln Thr Lys Val Leu Lys Gln Leu Leu Met Leu Gln Ser Leu Glu								
	145 150 155							
tga gggc aggcgagccc caccgcggcc ccggcccttc ctggactcgc ctgctcgctt	778							
*								
ccccctccca ggcccgtggc caaccagca gtccttcctt cagctgccta ggaggaaggg	838							

ctg aac gag aac gtc ctc aag cct gcg cag gag aag gtg aag gag gga	822
Leu Asn Glu Asn Val Leu Lys Pro Ala Gln Glu Lys Val Lys Glu Gly	
250 255 260	
aag att ttt gat gat gtc tcc agt ggg gtc tct cag ttg gcg tcc aag	870
Lys Ile Phe Asp Asp Val Ser Ser Gly Val Ser Gln Leu Ala Ser Lys	
265 270 275	
gtc cag gga gtc ggt agt aag gga tgg cgg gac gtc acc acc ttt ttt	918
Val Gln Gly Val Gly Ser Lys Gly Trp Arg Asp Val Thr Thr Phe Phe	
280 285 290	
tcg ggg aaa gca gag ggc ccc ttg gac agc ccc tcg gag ggc cac agt	966
Ser Gly Lys Ala Glu Gly Pro Leu Asp Ser Pro Ser Glu Gly His Ser	
295 300 305 310	
tat cag aac agc ggt ctg gac cac ttc caa aac agc aac ata gac cag	1014
Tyr Gln Asn Ser Gly Leu Asp His Phe Gln Asn Ser Asn Ile Asp Gln	
315 320 325	
agc ttc tgg gag acc ttt gga agt gct gag ccc acc aag acc cgc aag	1062
Ser Phe Trp Glu Thr Phe Gly Ser Ala Glu Pro Thr Lys Thr Arg Lys	
330 335 340	
tcc ccg agc agc gac agc tgg acg tgc gcg gac acc tcc acc gag agg	1110
Ser Pro Ser Ser Asp Ser Trp Thr Cys Ala Asp Thr Ser Thr Glu Arg	
345 350 355	
agg agc tcg gac agc tgg gag gtg tgg ggc tcg gcc tcc acc aac agg	1158
Arg Ser Ser Asp Ser Trp Glu Val Trp Gly Ser Ala Ser Thr Asn Arg	
360 365 370	
aac agc aac agc gac ggc ggg gag ggc ggg gag ggc acc aag aag gca	1206
Asn Ser Asn Ser Asp Gly Gly Glu Gly Gly Glu Gly Thr Lys Lys Ala	
375 380 385 390	
gtg ccg ccg gcc gtg ccc act gat gat ggc tgg gac aac cag aac tgg	1254
Val Pro Pro Ala Val Pro Thr Asp Asp Gly Trp Asp Asn Gln Asn Trp	
395 400 405	
tag ggcc cactgcgccc ccgtccccag cgccccggg cgacttcgtg tttgcactct	1311
*	
gccctcgtcg ttctctctcc ttccatttga cccaagaatc agcaactgca gtgtgaggac	1371
agcgtctcgg gaggcaggac cctagggaga cccgggtgtg cgccgcctgc gcgtggggag	1431
tcttcggtgc gtgggggagg cttgctgtcc agcctgtgtg ggggccgtcc cgtcccacac	1491
tcccctgggc attcttgagc tcaaggccgg ggctctgcgt ggcttgctgg gaggtgggct	1551
gcagcacaga ggctgtgac tgcgttccag cggccagttc actacgcagt atctctgggg	1611
cctgggacca gccacgtgcc gagctgtcag cgacgtgagg tgtcccttct cgttgagata	1671

<212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (596)..(1840)

<400> 27

tttttgtatg caccacgggc ggcggtggtc ggtgcgggag gagggagggg agcttgcggg	60
cccgagaggg ggcgacggcg gcggcggtgg cctgaggagg cccgagcggc ggcggtggcg	120
gcgaaggccg aggcgtctag gtgttttttg aagagctgca gccctcttct cacagatgag	180
ctacgaggag atgatgacac tgactgagca gcacctggag tctcagaacg tcaccaaagg	240
tgcccgccac aagatagccc tgagcatcca gaagctgcgt gagagacaga gcgtcctcaa	300
gtccctagag aaggatgtgc tggaaggcgg gaacctacga aacgctctgc aggagctgca	360
gcagatcatt atcactccca tcaaggcta cagtgtcctc caggccaccg tggtgcccgc	420
caccaccacc cctactgcca aggatggggc cccgggggaa ccaccgctgc caggtgctga	480
gcctccccta gcccaccccg gcacagacaa aggcaccgag gccagagccg ggaccatgtg	540
acggcgctgg ccctcgccac cgccgtcccc cgacctggc cccaggcccg gcacc atg	598
	Met
	1
atg ttc cga gac cag gtg ggc atc ctc gct ggc tgg ttc aaa ggc tgg	646
Met Phe Arg Asp Gln Val Gly Ile Leu Ala Gly Trp Phe Lys Gly Trp	
	5 10 15
aat gag tgt gag cag aca gtg gcc ctc ctg tca ctt ccg aaa cgg gtc	694
Asn Glu Cys Glu Gln Thr Val Ala Leu Leu Ser Leu Pro Lys Arg Val	
	20 25 30
acc cgt acc cag gcc cgc ttc ctg cag ctc tgc ctg gag cac tca ctg	742
Thr Arg Thr Gln Ala Arg Phe Leu Gln Leu Cys Leu Glu His Ser Leu	
	35 40 45
gcg gac tgc aat gac atc cac ctg ctg gag tcg gag gcc aac agt gct	790
Ala Asp Cys Asn Asp Ile His Leu Leu Glu Ser Glu Ala Asn Ser Ala	
	50 55 60 65
gcc atc gtc agc cag tgg cag cag gag tcc aaa gag aag gtg gtg tcc	838
Ala Ile Val Ser Gln Trp Gln Gln Glu Ser Lys Glu Lys Val Val Ser	
	70 75 80
ctc ctg ctg tcc cac ctt ccc ctg ctt cag cca ggc aac aca gag gcc	886
Leu Leu Leu Ser His Leu Pro Leu Leu Gln Pro Gly Asn Thr Glu Ala	
	85 90 95
aag tcg gag tac atg agg cta ctg cag aaa gtg ctg gcc tac tca atc	934
Lys Ser Glu Tyr Met Arg Leu Leu Gln Lys Val Leu Ala Tyr Ser Ile	
	100 105 110

gag agc aat gct ttc atc gag gag agt cgc cag ctg ctt tcc tat gcc	982
Glu Ser Asn Ala Phe Ile Glu Glu Ser Arg Gln Leu Leu Ser Tyr Ala	
115 120 125	
ctc atc cac cca gcc acc aca ctg gag gac cgc aac gca ctg gcc ctc	1030
Leu Ile His Pro Ala Thr Thr Leu Glu Asp Arg Asn Ala Leu Ala Leu	
130 135 140 145	
tgg ctg agc cac ctg gaa gag cgg ttg gct agt ggc ttc cgc tcc cgg	1078
Trp Leu Ser His Leu Glu Glu Arg Leu Ala Ser Gly Phe Arg Ser Arg	
150 155 160	
cca gag ccc tcc tac cat tca cgt caa ggc tca gat gag tgg ggg ggc	1126
Pro Glu Pro Ser Tyr His Ser Arg Gln Gly Ser Asp Glu Trp Gly Gly	
165 170 175	
cct gca gag cta ggc cct ggg gag gca ggg cca ggc tgg cag gac aag	1174
Pro Ala Glu Leu Gly Pro Gly Glu Ala Gly Pro Gly Trp Gln Asp Lys	
180 185 190	
cca ccc cgg gaa aat gga cac gtg ccc ttc cac cca tcc agc tca gtg	1222
Pro Pro Arg Glu Asn Gly His Val Pro Phe His Pro Ser Ser Ser Val	
195 200 205	
ccg cca gcc atc aac agt att ggg agc aat gca aac aca ggt ctc ccc	1270
Pro Pro Ala Ile Asn Ser Ile Gly Ser Asn Ala Asn Thr Gly Leu Pro	
210 215 220 225	
tgc caa atc cac cct agc cca ctg aag cgc tcc atg tca ctc atc cct	1318
Cys Gln Ile His Pro Ser Pro Leu Lys Arg Ser Met Ser Leu Ile Pro	
230 235 240	
aca agc ccc cag gtc cct ggt gag tgg ccg agt cca gag gag ctt ggg	1366
Thr Ser Pro Gln Val Pro Gly Glu Trp Pro Ser Pro Glu Glu Leu Gly	
245 250 255	
gcc cgg gct gct ttt acc acg ccc gat cac gca cct ctc tcg ccc cag	1414
Ala Arg Ala Ala Phe Thr Thr Pro Asp His Ala Pro Leu Ser Pro Gln	
260 265 270	
agc agc gtg gcc tcc tct ggc agt gag cag aca gag gag cag ggc tcc	1462
Ser Ser Val Ala Ser Ser Gly Ser Glu Gln Thr Glu Glu Gln Gly Ser	
275 280 285	
agc cgg aac acc ttc cag gag gat ggc agt ggc atg aaa gat gtg ccc	1510
Ser Arg Asn Thr Phe Gln Glu Asp Gly Ser Gly Met Lys Asp Val Pro	
290 295 300 305	
tca tgg ctc aag agc ctc cgt ttg cac aag tat gca gcc ctc ttc tca	1558
Ser Trp Leu Lys Ser Leu Arg Leu His Lys Tyr Ala Ala Leu Phe Ser	
310 315 320	
cag atg agc tac gag gag atg atg aca ctg act gag cag cac ctg gag	1606
Gln Met Ser Tyr Glu Glu Met Met Thr Leu Thr Glu Gln His Leu Glu	
325 330 335	

tct cag aac gtc acc aaa ggt gcc cgc cac aag ata gcc ctg agc atc	1654
Ser Gln Asn Val Thr Lys Gly Ala Arg His Lys Ile Ala Leu Ser Ile	
340 345 350	
cag aag ctg cgt gag aga cag agc gtc ctc aag tcc cta gag aag gat	1702
Gln Lys Leu Arg Glu Arg Gln Ser Val Leu Lys Ser Leu Glu Lys Asp	
355 360 365	
gtg ctg gaa ggc ggg aac cta cga aac gct ctg cag gag ctg cag cag	1750
Val Leu Glu Gly Gly Asn Leu Arg Asn Ala Leu Gln Glu Leu Gln Gln	
370 375 380 385	
atc atc atc act ccc atc aag gcc tac agt gtc ctc cag gcc acc gtg	1798
Ile Ile Ile Thr Pro Ile Lys Ala Tyr Ser Val Leu Gln Ala Thr Val	
390 395 400	
gct gcc gcc acc ctg tat tgc ggc cgc tct aga gga tcc tag cttcgta	1847
Ala Ala Ala Thr Leu Tyr Cys Gly Arg Ser Arg Gly Ser *	
405 410 415	
cgc	1850

<210> 28
 <211> 4781
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> CDS
 <222> (813) .. (3635)

<400> 28	
tttgtctgtgg aattccccggg tcgacccacg cgtccgcgga cgcgtggggtt tagaggtaag	60
tttgccctact ttgtcgtcta gtgggtaaaa ttttgcgag agcgttgat ctgggaagcg	120
ggatagggat ggatgggttc atttgagagc cacggcttaa agcggttgcg atcaggatgg	180
gacacagggtt tgtttgggga caacaaagat ggcatttgtg agtgttttga agcaaccogt	240
actgattaca tctttctccc ttgtgttcc tttatcccag gtttgaattt tctcggagaa	300
agacaggccg gccacgagga aacagaaac aagccgcagc aacatctaag cccttgaaag	360
gaccttgaga gaggggggaa agggaaaaca gcagccacca gcccaaccac ttgtgtcttc	420
tgccccctcc cacctatctt gccacccca ccagcccacg ctgcttggga cttgaaatct	480
gtggccgaag gaccgtcact acataacttc aaaaataatc aaccaccctc ccttcccaaa	540
ccacccaaat tcatcatcc agcgtttact tttttgaatc cactcagaac tttttctgc	600
gacccccctc cctaaatgga gttgggtggg ggggaaatga atactgagtt ggcctttatt	660

ttttaaaaga ctttttgatc caatgaggcc ccctaaataa ttgagttttg ggtcctggtt	720
ggttggtttta ttttttttcc tccaaaattt taccctctcc cccctgagcc cgaggtgctg	780
acgtcgcaaa aaaattggat aaaaccacca tc atg ggt tcg ggt ccc ata gac	833
Met Gly Ser Gly Pro Ile Asp	
1 5	
ccc aaa gaa ctt ctc aag ggc ctg gac agc ttc ctt aac cga gat ggg	881
Pro Lys Glu Leu Leu Lys Gly Leu Asp Ser Phe Leu Asn Arg Asp Gly	
10 15 20	
gaa gtc aaa agt gtg gat ggg att tcc aag atc ttc agt ttg atg aag	929
Glu Val Lys Ser Val Asp Gly Ile Ser Lys Ile Phe Ser Leu Met Lys	
25 30 35	
gaa gca cga aag atg gtg agt cga tgc act tac ttg aac att ctc ctg	977
Glu Ala Arg Lys Met Val Ser Arg Cys Thr Tyr Leu Asn Ile Leu Leu	
40 45 50 55	
cag acc cgt tca cca gaa ata ttg gtc aaa ttt att gac gtt ggc ggc	1025
Gln Thr Arg Ser Pro Glu Ile Leu Val Lys Phe Ile Asp Val Gly Gly	
60 65 70	
tac aaa ctt ctt aac aat tgg ctg acg tat tca aag aca acc aac aac	1073
Tyr Lys Leu Leu Asn Asn Trp Leu Thr Tyr Ser Lys Thr Thr Asn Asn	
75 80 85	
att ccc ctc ctc cag caa att cta ctg acc ctg cag cat cta ccg ctc	1121
Ile Pro Leu Leu Gln Gln Ile Leu Leu Thr Leu Gln His Leu Pro Leu	
90 95 100	
act gta gac cat ctc aag cag aac aac aca gct aaa ctg gtg aag cag	1169
Thr Val Asp His Leu Lys Gln Asn Asn Thr Ala Lys Leu Val Lys Gln	
105 110 115	
ctg agc aag tca agt gag gat gaa gag ctc cgg aaa ttg gcc tca gtc	1217
Leu Ser Lys Ser Ser Glu Asp Glu Glu Leu Arg Lys Leu Ala Ser Val	
120 125 130 135	
ctt gtc agc gac tgg atg gct gtc atc cgc tct cag agc agt acc cag	1265
Leu Val Ser Asp Trp Met Ala Val Ile Arg Ser Gln Ser Ser Thr Gln	
140 145 150	
cct gct gag aaa gat aag aag aaa cgt aaa gat gaa gga aaa agt cga	1313
Pro Ala Glu Lys Asp Lys Lys Lys Arg Lys Asp Glu Gly Lys Ser Arg	
155 160 165	
acc acc ctt cct gag cga cct ttg aca gag gtg aag gct gag acc cgg	1361
Thr Thr Leu Pro Glu Arg Pro Leu Thr Glu Val Lys Ala Glu Thr Arg	
170 175 180	
gct gag gag gcc cca gag aag aag agg gag aag ccc aag tct ctt cgc	1409
Ala Glu Glu Ala Pro Glu Lys Lys Arg Glu Lys Pro Lys Ser Leu Arg	
185 190 195	
acc aca gca ccc agt cat gcc aag ttc cgt tcc act gga cta gag ctg	1457

Thr	Thr	Ala	Pro	Ser	His	Ala	Lys	Phe	Arg	Ser	Thr	Gly	Leu	Glu	Leu	
200					205					210					215	
gag	aca	cca	tcc	ttg	gtg	cct	gtg	aag	aag	aat	gcc	agc	aca	gtg	gtg	1505
Glu	Thr	Pro	Ser	Leu	Val	Pro	Val	Lys	Lys	Asn	Ala	Ser	Thr	Val	Val	
				220					225					230		
gtt	tct	gac	aag	tac	aac	ctt	aaa	ccc	atc	ccc	ctc	aaa	cgt	cag	agc	1553
Val	Ser	Asp	Lys	Tyr	Asn	Leu	Lys	Pro	Ile	Pro	Leu	Lys	Arg	Gln	Ser	
			235					240					245			
aac	gta	gct	gct	cca	gga	gat	gcc	act	ccc	cct	gca	gag	aag	aaa	tac	1601
Asn	Val	Ala	Ala	Pro	Gly	Asp	Ala	Thr	Pro	Pro	Ala	Glu	Lys	Lys	Tyr	
			250				255					260				
aag	cca	ctc	aac	aca	aca	cct	aat	gcc	acc	aaa	gag	atc	aaa	gtg	aag	1649
Lys	Pro	Leu	Asn	Thr	Thr	Pro	Asn	Ala	Thr	Lys	Glu	Ile	Lys	Val	Lys	
			265			270					275					
atc	atc	ccg	cca	cag	cct	atg	gag	ggc	ctg	ggc	ttt	ctg	gat	gct	ctt	1697
Ile	Ile	Pro	Pro	Gln	Pro	Met	Glu	Gly	Leu	Gly	Phe	Leu	Asp	Ala	Leu	
280					285					290					295	
aat	tca	gcc	cct	gtt	cca	ggc	atc	aaa	att	aag	aag	aaa	aaa	aaa	gta	1745
Asn	Ser	Ala	Pro	Val	Pro	Gly	Ile	Lys	Ile	Lys	Lys	Lys	Lys	Lys	Val	
				300					305						310	
ctg	tca	cct	acg	gct	gcc	aag	cca	agc	ccc	ttt	gaa	ggg	aaa	acg	agc	1793
Leu	Ser	Pro	Thr	Ala	Ala	Lys	Pro	Ser	Pro	Phe	Glu	Gly	Lys	Thr	Ser	
			315					320					325			
aca	gaa	cca	agc	aca	gcc	aaa	cct	tct	tcc	cca	gaa	cca	gca	cca	cct	1841
Thr	Glu	Pro	Ser	Thr	Ala	Lys	Pro	Ser	Ser	Pro	Glu	Pro	Ala	Pro	Pro	
			330				335						340			
tct	gag	gca	atg	gac	gca	gac	cgt	cca	ggc	acc	ccg	gtt	ccc	cct	gtt	1889
Ser	Glu	Ala	Met	Asp	Ala	Asp	Arg	Pro	Gly	Thr	Pro	Val	Pro	Pro	Val	
			345			350				355						
gaa	gtc	ccg	gag	ctc	atg	gat	aca	gcc	tct	ttg	gag	cca	gga	gct	ctg	1937
Glu	Val	Pro	Glu	Leu	Met	Asp	Thr	Ala	Ser	Leu	Glu	Pro	Gly	Ala	Leu	
360					365					370					375	
gat	gcc	aag	cca	gtg	gag	agt	cct	gga	gat	cct	aac	caa	ctg	acc	cgg	1985
Asp	Ala	Lys	Pro	Val	Glu	Ser	Pro	Gly	Asp	Pro	Asn	Gln	Leu	Thr	Arg	
				380					385					390		
aaa	ggc	agg	aag	agg	aaa	agt	gtg	aca	tgg	cct	gag	gaa	ggc	aaa	ctg	2033
Lys	Gly	Arg	Lys	Arg	Lys	Ser	Val	Thr	Trp	Pro	Glu	Glu	Gly	Lys	Leu	
			395					400					405			
aga	gaa	tat	ttc	tat	ttt	gaa	ttg	gat	gaa	act	gaa	cga	gta	aat	gtg	2081
Arg	Glu	Tyr	Phe	Tyr	Phe	Glu	Leu	Asp	Glu	Thr	Glu	Arg	Val	Asn	Val	
			410				415					420				
aat	aag	atc	aag	gac	ttt	ggt	gag	gcg	gct	aag	cga	gag	ata	ctg	tca	2129
Asn	Lys	Ile	Lys	Asp	Phe	Gly	Glu	Ala	Ala	Lys	Arg	Glu	Ile	Leu	Ser	

425	430	435	
gac cga cat gca ttt gag aca gcg cgg cgt ctg agc cat gat aac atg Asp Arg His Ala Phe Glu Thr Ala Arg Arg Leu Ser His Asp Asn Met 440 445 450 455			2177
gag gag aag gtg ccc tgg gtg tgc ccc cgg ccc ctg gtt ctg ccc tca Glu Glu Lys Val Pro Trp Val Cys Pro Arg Pro Leu Val Leu Pro Ser 460 465 470			2225
cct ctt gtc acc cct gga agc aat agt cag gag cga tat atc cag gct Pro Leu Val Thr Pro Gly Ser Asn Ser Gln Glu Arg Tyr Ile Gln Ala 475 480 485			2273
gag cgg gag aag gga atc ctt cag gag ctc ttc ctg aac aag gag agt Glu Arg Glu Lys Gly Ile Leu Gln Glu Leu Phe Leu Asn Lys Glu Ser 490 495 500			2321
cct cat gag cct gat cct gag ccc tac gag ccc ata ccc cct aaa ctc Pro His Glu Pro Asp Pro Glu Pro Tyr Glu Pro Ile Pro Pro Lys Leu 505 510 515			2369
atc ccc cta gat gag gag tgt tcc atg gat gag act ccg tat gtt gag Ile Pro Leu Asp Glu Glu Cys Ser Met Asp Glu Thr Pro Tyr Val Glu 520 525 530 535			2417
act ctg gaa cct ggg ggg tca ggt ggc tca cct gat ggg gca gga ggc Thr Leu Glu Pro Gly Gly Ser Gly Gly Ser Pro Asp Gly Ala Gly Gly 540 545 550			2465
tcc aag ttg cct cca gtt ctg gcc aat ctt atg gga agc atg ggt gct Ser Lys Leu Pro Pro Val Leu Ala Asn Leu Met Gly Ser Met Gly Ala 555 560 565			2513
gga aag ggc ccc caa ggc cct gga gga gga ggc att aat gtc caa gag Gly Lys Gly Pro Gln Gly Pro Gly Gly Gly Gly Ile Asn Val Gln Glu 570 575 580			2561
atc ctc acc tcc atc atg ggt agc cca aac agt cat cct tca gag gaa Ile Leu Thr Ser Ile Met Gly Ser Pro Asn Ser His Pro Ser Glu Glu 585 590 595			2609
cta ctg aaa caa cca gac tat tcg gac aag atc aag cag atg ctg gtg Leu Leu Lys Gln Pro Asp Tyr Ser Asp Lys Ile Lys Gln Met Leu Val 600 605 610 615			2657
cca cat gga ctc cta ggc cct ggc cca ata gcc aat ggt ttc cca cca Pro His Gly Leu Leu Gly Pro Gly Pro Ile Ala Asn Gly Phe Pro Pro 620 625 630			2705
ggg ggt cct ggg ggc ccc aag ggc atg cag cac ttt ccc cct gga cct Gly Gly Pro Gly Gly Pro Lys Gly Met Gln His Phe Pro Pro Gly Pro 635 640 645			2753
ggg gga cct atg cca ggt ccc cat gga ggc cct ggt ggg cca gtg ggt Gly Gly Pro Met Pro Gly Pro His Gly Gly Pro Gly Gly Pro Val Gly 650 655 660			2801

cca cgt ctt ctg ggt cct cca ccc cct ccc cgg gga ggt gat ccc ttc	2849
Pro Arg Leu Leu Gly Pro Pro Pro Pro Arg Gly Gly Asp Pro Phe	
665 670 675	
tgg gat ggc ccg ggc gac cct atg cgg ggt ggc cca atg cgg ggg ggt	2897
Trp Asp Gly Pro Gly Asp Pro Met Arg Gly Gly Pro Met Arg Gly Gly	
680 685 690 695	
cca gga cca ggt cct gga cca tac cat aga ggc cga ggt ggc cga gga	2945
Pro Gly Pro Gly Pro Gly Pro Tyr His Arg Gly Arg Gly Gly Arg Gly	
700 705 710	
gga aac gaa cct cct cct cct cct cct cca ttc cga ggc gcc aga gga	2993
Gly Asn Glu Pro Pro Pro Pro Pro Pro Phe Arg Gly Ala Arg Gly	
715 720 725	
ggt cgc tct gga gga gga ccc cca aat gga cga ggg ggc cct ggt ggg	3041
Gly Arg Ser Gly Gly Gly Pro Pro Asn Gly Arg Gly Gly Pro Gly Gly	
730 735 740	
ggc atg gtt gga ggt ggt ggg cat cgt cct cac gaa ggc cct ggt ggg	3089
Gly Met Val Gly Gly Gly Gly His Arg Pro His Glu Gly Pro Gly Gly	
745 750 755	
ggc atg ggc aac agc agt gga cat cgt ccc cac gaa ggc cct ggc ggt	3137
Gly Met Gly Asn Ser Ser Gly His Arg Pro His Glu Gly Pro Gly Gly	
760 765 770 775	
ggc atg gga agt ggg cat cgc ccc cat gaa ggc cct ggt ggt agc atg	3185
Gly Met Gly Ser Gly His Arg Pro His Glu Gly Pro Gly Gly Ser Met	
780 785 790	
ggt ggg ggt gga gga cat cgt ccc cac gaa ggc cct ggc ggt ggc atc	3233
Gly Gly Gly Gly Gly His Arg Pro His Glu Gly Pro Gly Gly Gly Ile	
795 800 805	
agt ggt ggc agt ggc cat cgt ccc cat gaa ggc cct ggc gga gga atg	3281
Ser Gly Gly Ser Gly His Arg Pro His Glu Gly Pro Gly Gly Gly Met	
810 815 820	
ggt gcc ggt ggt gga cat cgc ccc cac gaa ggc cct ggc gga agc atg	3329
Gly Ala Gly Gly Gly His Arg Pro His Glu Gly Pro Gly Gly Ser Met	
825 830 835	
ggt gga agt ggt gga cat cgt ccc cat gaa ggc cct gga cac ggg ggg	3377
Gly Gly Ser Gly Gly His Arg Pro His Glu Gly Pro Gly His Gly Gly	
840 845 850 855	
ccc cat ggc cac cgg cct cat gat gtc cct ggt cac cga ggc cat gac	3425
Pro His Gly His Arg Pro His Asp Val Pro Gly His Arg Gly His Asp	
860 865 870	
cat cga ggg ccg cca cct cat gag cac cgt ggc cat gat ggt cct ggc	3473
His Arg Gly Pro Pro His Glu His Arg Gly His Asp Gly Pro Gly	
875 880 885	

cac ggg gga ggg ggc cac cga ggg cac gat gga ggc cac agc cat gga	3521
His Gly Gly Gly Gly His Arg Gly His Asp Gly Gly His Ser His Gly	
890 895 900	
gga gac atg tca aac cgc cct gtc tgc cga cat ttc atg atg aag ggc	3569
Gly Asp Met Ser Asn Arg Pro Val Cys Arg His Phe Met Met Lys Gly	
905 910 915	
aac tgc cgc tat gag aac aac tgt gcc ttc tac cac ccg ggt gtc aat	3617
Asn Cys Arg Tyr Glu Asn Asn Cys Ala Phe Tyr His Pro Gly Val Asn	
920 925 930 935	
ggg ccc ccc ctg ccc tag ggacca tttgcctgcc ctgttcacac aacccctgtg	3671
Gly Pro Pro Leu Pro *	
940	
gactgcagcc tcgctctttc caccctgtta tggcttctgt gaggccatt ttcccttttc	3731
cccagctgat gaggagccgg cccctcagt tccacttgc ttgggttcct gggggttttc	3791
tgatcactgg tgcgcattga tgtacatatt ttcctccagt ctggggagga gagagactgg	3851
aaacgttcct ggactgctga agaggagacc cagttggctt cactttttga gaagattcgc	3911
cctgtacccc aaaccccttt ccagtattac ccttaatgct tgagaacctt aagctgggta	3971
tcctggcgaa caccctacc cttctattgc ggggtccccc atgcacacag aactctgaca	4031
caggatcagc tgcacttaag aaatcatccc agctaagttc attattcctc atgggggtggg	4091
gagatgctga aaggggtatt gtatatccca ctgcactgag agggctcaat cagctggatt	4151
tgagttctgg aacacacatc atccccaccc ctccccagc gtgggctcac cattcttagt	4211
cctttctcaa gtgggacctt caactttctg tgaacaccca gtctgcgtcc tgggtctgct	4271
aggttcgatg atggcgaact cgtatctgca tccggtgcaa gtttttagctg gcagaggtga	4331
gaccggtggt gctggtctgc ctttgccaac tatagccagt ctggagactt gataaaatac	4391
ttcagtgaga ccagcttctc atcaacttgg gcccggcgtg ctgggcctga aagtcacact	4451
acatgcactg cctttgggag tcagctcact cctgtctccc acctggaacc ttgccagcgt	4511
gaaggaggct tccaggtact tcaccctgtc aaccacctct gaatccccac caggcgctt	4571
cctgggtgga ttcaacaaga tgattttgcc ctttcccagt tctctccttc actttggcat	4631
cagttgtttt ctatgaaaac agtggattgg ttgggttttg tgcagggtct tgggttagag	4691
ccaaaatgga tttgaggatg agtatTTTTT tttttggttt tgtatatattt gtacattaat	4751
aataaacagt ggaaagagaa aaaaaaaaaa	4781

<210> 29
 <211> 968
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (184)..(411)

<220>
 <221> misc_feature
 <222> (1)...(968)
 <223> n = a,t,c or g

<400> 29

```

tttttttttt attattatac tttaagtttt agggtagatg tgcacaacgt gcagggtttgt      60
tacatatgta tacatgtgcc atgttagtgt gctgcaccca ttaactcgtc atttagcatt      120
aggtatatct cctacagcta tccctcccc cttccccac cccacaaaag gtcccagtgt      180
gtg  atg ttc ccc ttc ctg tgt cca tgt gtt ctc att gtt caa ttc cca      228
    Met Phe Pro Phe Leu Cys Pro Cys Val Leu Ile Val Gln Phe Pro
        1             5             10             15

cct atg agt gag aac atg tgg gtt tgg ttt ttt tgt cct tgc gat agt      276
Pro Met Ser Glu Asn Met Trp Val Trp Phe Phe Cys Pro Cys Asp Ser
           20             25             30

ttg ctg aga atg atg gtt tcc agc ttc atc cat gtc cct gcg aag gac      324
Leu Leu Arg Met Met Val Ser Ser Phe Ile His Val Pro Ala Lys Asp
           35             40             45

atg aac tca ccc ttt ttt atg gaa tac tac aca gcc ata aaa agg aat      372
Met Asn Ser Pro Phe Phe Met Glu Tyr Tyr Thr Ala Ile Lys Arg Asn
           50             55             60

gac aac aca tcc ctt gca ggg aca tgg atg gag caa tag gccattatcc      421
Asp Asn Thr Ser Leu Ala Gly Thr Trp Met Glu Gln  *
           65             70             75

gtagcaaact aatgcaggaa gagaaaaccc agtactgcat gttctcactt ataagtggga      481
gctaaatgat gagaacacat gaacacaaag aggggaacag acactagggc cgtttagaag      541
ttggcggggtg gtttgctttt ttntttagnt acangattta ttagnaatgg gtactaggct      601
gaataccgtt gtggatggta gtaatcgggt gaacaaagcg cccatgttca caatttttagc      661
ttatttttttc tggttttcgt catggtaccc cttgtgagtg tgagttatat gtttggttga      721
tatttatcgg ttattggatt tggtcctcgg gtggtgtttt cgtggagtggt tgggggttgtt      781
gtgattcttg tgtttggttg tgggtgtttaa aagaggtgta ggttgggtgac ccgggtgttg      841
tccacgttat tggctttgat gaggccccct tttgtgggtg gtgtttggtg ttgttttggt      901

```

ctggtcggtg gccctatttg tgctttttgt tggtagtggc ttccctggtgg tcggtgagcg 961
gtgtgttt 968

<210> 30
<211> 1082
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (168) .. (725)

<400> 30
ttttgaaaaa ctgggatagc agtaccacgc tggctagcgt ttaaacttaa gcttgggtacc 60
gagctcggat ccactagtcc agtgtggtgg aattccggcc aagtagagct ccgtcctgac 120
gcgccgcctc ccgtgggctc cggccggcta agccgcggcg gacaact atg ctg aaa 176
Met Leu Lys
1
gcc aag atc ctc ttc gtg ggg cct tgc gag agt gga aaa act gtt ttg 224
Ala Lys Ile Leu Phe Val Gly Pro Cys Glu Ser Gly Lys Thr Val Leu
5 10 15
gcc aac ttt ctg aca gaa tct tct gac atc act gaa tac agc cca acc 272
Ala Asn Phe Leu Thr Glu Ser Ser Asp Ile Thr Glu Tyr Ser Pro Thr
20 25 30 35
caa gga gtg agg atc cta gaa ttt gag aac ccg cat gtt acc agc aac 320
Gln Gly Val Arg Ile Leu Glu Phe Glu Asn Pro His Val Thr Ser Asn
40 45 50
aac aaa ggc acg ggc tgt gaa ttc gag cta tgg gac tgt ggt ggc gat 368
Asn Lys Gly Thr Gly Cys Glu Phe Glu Leu Trp Asp Cys Gly Gly Asp
55 60 65
gct aag ttt gag tcc tgc tgg ccg gcc ctg atg aag gat gct cat gga 416
Ala Lys Phe Glu Ser Cys Trp Pro Ala Leu Met Lys Asp Ala His Gly
70 75 80
gtg gtg atc gtc ttc aat gct gac atc cca agc cac cgg aag gaa atg 464
Val Val Ile Val Phe Asn Ala Asp Ile Pro Ser His Arg Lys Glu Met
85 90 95
gag atg tgg tat tcc tgc ttt gtc caa cag ccg tcc tta cag gac aca 512
Glu Met Trp Tyr Ser Cys Phe Val Gln Gln Pro Ser Leu Gln Asp Thr
100 105 110 115
cag tgt atg cta att gca cac cac aaa cca ggc tct gga gat gat aaa 560
Gln Cys Met Leu Ile Ala His His Lys Pro Gly Ser Gly Asp Asp Lys
120 125 130

gga agc ctg tct ttg tgc cca ccc ttg aac aag ctg aag ctg gtg cac	608
Gly Ser Leu Ser Leu Ser Pro Pro Leu Asn Lys Leu Lys Leu Val His	
135 140 145	
tca aac ctg gaa gat gac cct gag gag atc cgg atg gaa ttc ata aag	656
Ser Asn Leu Glu Asp Asp Pro Glu Glu Ile Arg Met Glu Phe Ile Lys	
150 155 160	
tat tta aaa agc ata atc aac tcc atg tct gag agc aga gac agg gag	704
Tyr Leu Lys Ser Ile Ile Asn Ser Met Ser Glu Ser Arg Asp Arg Glu	
165 170 175	
gag atg tca att atg acc tag cc agccttcacc tgggactgcc acatccccag	757
Glu Met Ser Ile Met Thr *	
180 185	
tgaaatcagc atgtttctcg gtgcagatct gaaatcacat ccagctcctg atgttttctt	817
ctccctctga ctgcagagga agtggtccta cctgcaggaa ggcacctgtc acacagggcg	877
ttcactcaga ccatctgtgc tctgccctga gttcagttga gaaaatccta ttatcaaatt	937
tggatttcct ggccccagaa cttcccaaag acctgtaaaa tggagggatt taccacctca	997
catatgtcca gttaaacagt ttgtggactt gtaaccgtcg cagcccaatg atacaacagt	1057
agtttaatca cgtgaaaaaa aaaaa	1082

```

<210> 31
<211> 1517
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (666)..(1406)

<220>
<221> misc_feature
<222> (1)...(1517)
<223> n = a,t,c or g

```

ctcgaaatcg atactttgcc ggaccgggnc ggnnnnccgg ggtcgacggg aggcaggagg	60
gccgaccag ggggtgctggc cgccctctgt gagaaaactg acaatgacat ccgggcctgc	120
atcaacaccc tgcagttcct gtacagccgg gccagcggg agctgagcgt gcgggacgtg	180
caggccacac gcgtgggcct caaggaccag cgcagagggc tcttctcggt gtggcaggag	240
gtcttccagc tgctcgagc ccagagcacc ccacctgcag gcgccgtgtg ggccaggacc	300
ccgccctgcc tgctgacaca ctctgctgg gtgacgggga cgcgggctcc ctcacctccg	360

175	180	185	190	
gtg gtc gtc agg agc aca gca gtc ccg agt gca ggg gac acg gcc ccg				1283
Val Val Val Arg Ser Thr Ala Val Pro Ser Ala Gly Asp Thr Ala Pro				
	195	200	205	
gag cag gac tca gtg gag cgg cgc atg ggc aca gcg gtg ggc agg agc				1331
Glu Gln Asp Ser Val Glu Arg Arg Met Gly Thr Ala Val Gly Arg Ser				
	210	215	220	
gag gtc tgg ttc cgc ttc aac gag ggt gtc tcc aac gcc gtg cgg cgc				1379
Glu Val Trp Phe Arg Phe Asn Glu Gly Val Ser Asn Ala Val Arg Arg				
	225	230	235	
agc ctg tac atc agg gac ttg ctc tag ttctc tgagccgcgg acatgccctc				1431
Ser Leu Tyr Ile Arg Asp Leu Leu *				
	240	245		
gcattgcttc ccgcagagtg cagagacagg aagctggaga tgtctttata aagtcacacc				1491
tttacagact gtaaaaaaaaa aaaaaa				1517
<210> 32				
<211> 618				
<212> DNA				
<213> Homo sapiens				
<220>				
<221> CDS				
<222> (498) .. (572)				
<400> 32				
ttttgcctaa gatacaatga taagtaaaaa catgttacag agtaataacg tgttgcatag				60
gacaataactg taaatgtggg tcattccatc aacagacact gagccctgcg acgtgcctgg				120
ctctattcta catctgaggg acacaagggtg aacaagacca ggccactgta ttcaacatct				180
acatttaatg gaaatttttg aaagaagact tgagagatta taacagtggg cttcagcatc				240
agggagtagg cctagaagaa gaggagggtca agaagtggct tttcttatta tcttcttaac				300
tcttcaaatt ttactatga gcaattatta ttttttatta aaatttttagg ccaggcttat				360
ggctgtaatc ctagcagttt gggaggccaa ggtgagcgga tcacttgagg ttgggagttc				420
gagaccagcc tgaccaacat ggagaaactc tgtctctact taaaaaaaaa tacaaaatta				480
gccaggcatg gtggcac atg cct gta gtc cca gct act cag gag act gag				530
	Met Pro Val Val Pro Ala Thr Gln Glu Thr Glu			
	1 5 10			
gca ggg gaa ttg cct gaa cct ggg aga cag agg ttg caa tga gccaaaga				579
Ala Gly Glu Leu Pro Glu Pro Gly Arg Gln Arg Leu Gln *				

15

20

25

tcacgcctcg tgccgaattc ttggcctcga gggccaaat

618

<210> 33

<211> 1188

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (313)..(849)

<400> 33

aaggatcctt aattaaatta atcccccccc cccggggaga aacgttctca ctcgctctct 60

gctcgtcgcg ggcgtcccc gccctctgct gccagaacct tggggatgtg cctagaccgc 120

gcgcagcaca cgtccggggc aaccgcgagc agaacaaacc tttggcgggc ggccaggagg 180

ctccctccca gccaccgccc ccctccagcg cctttttttc ccccatata atacaagatc 240

ttccttctc agttccctta aagcacagcc cagggaacc tcctcacagt tttcatccag 300

ccacggggcca gc atg tct ggg ggc aaa tac gta gac tcg gag gga cat 348

Met Ser Gly Gly Lys Tyr Val Asp Ser Glu Gly His
1 5 10

ctc tac acc gtt ccc atc cgg gaa cag ggc aac atc tac aag ccc aac 396

Leu Tyr Thr Val Pro Ile Arg Glu Gln Gly Asn Ile Tyr Lys Pro Asn
15 20 25

aac aag gcc atg gca gac gag ctg agc gag aag caa gtg tac gac gcg 444

Asn Lys Ala Met Ala Asp Glu Leu Ser Glu Lys Gln Val Tyr Asp Ala
30 35 40

cac acc aag gag atc gac ctg gtc aac cgc gac cct aaa cac ctc aac 492

His Thr Lys Glu Ile Asp Leu Val Asn Arg Asp Pro Lys His Leu Asn
45 50 55 60

gat gac gtg gtc aag att gac ttt gaa gat gtg att gca gaa cca gaa 540

Asp Asp Val Val Lys Ile Asp Phe Glu Asp Val Ile Ala Glu Pro Glu
65 70 75

ggg aca cac agt ttt gac ggc att tgg aag gcc agc ttc acc acc ttc 588

Gly Thr His Ser Phe Asp Gly Ile Trp Lys Ala Ser Phe Thr Thr Phe
80 85 90

act gtg acg aaa tac tgg ttt tac cgc ttg ctg tct gcc ctc ttt ggc 636

Thr Val Thr Lys Tyr Trp Phe Tyr Arg Leu Leu Ser Ala Leu Phe Gly
95 100 105

atc ccg atg gca ctc atc tgg ggc att tac ttc gcc att ctc tct ttc 684

Ile Pro Met Ala Leu Ile Trp Gly Ile Tyr Phe Ala Ile Leu Ser Phe

110	115	120	
ctg cac atc tgg gca gtt gta cca tgc att aag agc ttc ctg att gag			732
Leu His Ile Trp Ala Val Val Pro Cys Ile Lys Ser Phe Leu Ile Glu			
125	130	135	140
att cag tgc atc agc cgt gtc tat tcc atc tac gtc cac acc gtc tgt			780
Ile Gln Cys Ile Ser Arg Val Tyr Ser Ile Tyr Val His Thr Val Cys			
	145	150	155
gac cca ctc ttt gaa gct gtt ggg aaa ata ttc agc aat gtc cgc atc			828
Asp Pro Leu Phe Glu Ala Val Gly Lys Ile Phe Ser Asn Val Arg Ile			
	160	165	170
aac ttg cag aaa gaa ata taa at gacatttcaa ggatagaagt atacctgatt			881
Asn Leu Gln Lys Glu Ile *			
175			
ttttttcctt ttaattttcc tgggtgccaat ttcaagttcc aagttgctaa tacagcaaca			941
atztatgaat tgaattatct tgggtgaaaa taaaaagatc actttctcag ttttcataag			1001
tattatgtct cttctgagct atttcatcta tttttggcag tctgaatttt taaaacccat			1061
ttaaattttt ttccttacct ttttatttgc atgtggatca accatcgctt tattggctga			1121
gatatgaaca tattgttgaa aggtaatttg agagaaatat gaagaactga ggaggaaaaa			1181
aaaaaaa			1188
<210>	34		
<211>	920		
<212>	DNA		
<213>	Homo sapiens		
<220>			
<221>	CDS		
<222>	(515) .. (697)		
<400>	34		
taagcttgcg gccgcaattt tttttttttt ttttttgtat tttttggtag agacgggatt			60
tcactatggt ggtcaggctg gtctcgaact cccgaccgca agtgatccac ccgccttggc			120
ctcccaaagt gctgggatta caagcttgag ccaactgcacc cagcctggaa agtatattta			180
tgaaagggtt gcactccaca aaagcatctt tgctaggggtg tcaaggaaga gatcactaaa			240
ccaaccccaa cacatccata caattccagc aatctagaga gggctgggtcc ttttcctttt			300
ctggattatt ttctgttctc agtaaaacaa gtatttactg tgatactgaa aactggggaa			360
attaacactg attaagatat tttaaact gagtcttaat tataacagaa ccagttttca			420


```

tcagaatgct tttacgtcac attcagtgaa gtgttacgct aatatattct acagccctga      480

agatagaaaa aagggtttctc tccaggtatg agat      atg gta caa aaa tac att      532
                                         Met Val Gln Lys Tyr Ile
                                         1              5

ttt cca cat aca aaa gag aga aaa aaa caa aga cat gtg gcg ggt ggc      580
Phe Pro His Thr Lys Glu Arg Lys Lys Gln Arg His Val Ala Gly Gly
              10              15              20

gag ggg agg ccc aat ccc aac acc cta caa ggt tcc atg gaa tgg aga      628
Glu Gly Arg Pro Asn Pro Asn Thr Leu Gln Gly Ser Met Glu Trp Arg
              25              30              35

agg aac aaa aaa atc ccc aat tat ttt ggg gta aga tgt gcc cca gaa      676
Arg Asn Lys Lys Ile Pro Asn Tyr Phe Gly Val Arg Cys Ala Pro Glu
              40              45              50

aag gtg aaa tct atg caa taa aa cccaggtttt cttcaaattct agcatctagg      729
Lys Val Lys Ser Met Gln *
              55              60

atttctatca gagtttcaaa taatcagaat ttctatcaga atttctaccc tgaggtgaca      789

cctactaact gtaggttctt tcattaataaa tgaagacatc tttcaccaga atgtatcaag      849

ctataaaaact ggcttcagag cctacactta gccagagtgg aaaaaaaaaa tagtgcatat      909

tttcgacagc a                                                                920

<210> 35
<211> 1233
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (133)..(1122)

<400> 35
gtgatcatcg acgctgcgg taccggtccg gaattcccgg gtcgaccac gcgtccgggc      60

ggcttcctag tgagtcggcg gctgatttag aaggaggttc aggctacggt gagccgaagc      120

cacacaggag cc      atg gaa gtg gca gag ccc agc agc ccc act gaa gag      168
                  Met Glu Val Ala Glu Pro Ser Ser Pro Thr Glu Glu
                  1              5              10

gag gag gag gaa gag gag cac tcg gca gag cct cgg ccc cgc act cgc      216
Glu Glu Glu Glu Glu Glu His Ser Ala Glu Pro Arg Pro Arg Thr Arg
              15              20              25

tcc aat cct gaa ggg gct gag gac cgg gca gta ggg gca cag gcc agc      264
Ser Asn Pro Glu Gly Ala Glu Asp Arg Ala Val Gly Ala Gln Ala Ser

```

30	35	40	
gtg ggc agc cgc agc gag ggt gag ggt gag gcc gcc agt gct gat gat Val Gly Ser Arg Ser Glu Gly Glu Gly Glu Ala Ala Ser Ala Asp Asp 45 50 55 60			312
ggg agc ctc aac act tca gga gcc ggc cct aag tcc tgg cag gtg ccc Gly Ser Leu Asn Thr Ser Gly Ala Gly Pro Lys Ser Trp Gln Val Pro 65 70 75			360
ccg cca gcc cct gag gtc caa att cgg aca cca agg gtc aac tgt cca Pro Pro Ala Pro Glu Val Gln Ile Arg Thr Pro Arg Val Asn Cys Pro 80 85 90			408
gag aaa gtg att atc tgc ctg gac ctg tca gag gaa atg tca ctg cca Glu Lys Val Ile Ile Cys Leu Asp Leu Ser Glu Glu Met Ser Leu Pro 95 100 105			456
aag ctg gag tgc ttc aac ggc tcc aaa acc aac gcc ctc aat gtc tcc Lys Leu Glu Ser Phe Asn Gly Ser Lys Thr Asn Ala Leu Asn Val Ser 110 115 120			504
cag aag atg att gag atg ttc gtg cgg aca aaa cac aag atc gac aaa Gln Lys Met Ile Glu Met Phe Val Arg Thr Lys His Lys Ile Asp Lys 125 130 135 140			552
agc cac gag ttt gca ctg gtg gtg gtg aac gat gac acg gcc tgg ctg Ser His Glu Phe Ala Leu Val Val Val Asn Asp Asp Thr Ala Trp Leu 145 150 155			600
tct ggc ctg acc tcc gac ccc cgc gag ctc tgt agc tgc ctc tat gat Ser Gly Leu Thr Ser Asp Pro Arg Glu Leu Cys Ser Cys Leu Tyr Asp 160 165 170			648
ctg gag acg gcc tcc tgt tcc acc ttc aat ctg gaa gga ctt ttc agc Leu Glu Thr Ala Ser Cys Ser Thr Phe Asn Leu Glu Gly Leu Phe Ser 175 180 185			696
ctc atc cag cag aaa act gag ctt ccg gtc aca gag aac gtg cag acg Leu Ile Gln Gln Lys Thr Glu Leu Pro Val Thr Glu Asn Val Gln Thr 190 195 200			744
att ccc ccg cca tat gtg gtc cgc acc atc ctt gtc tac agc cgt cca Ile Pro Pro Pro Tyr Val Val Arg Thr Ile Leu Val Tyr Ser Arg Pro 205 210 215 220			792
cct tgc cag ccc cag ttc tcc ttg acg gag ccc atg aag aaa atg ttc Pro Cys Gln Pro Gln Phe Ser Leu Thr Glu Pro Met Lys Lys Met Phe 225 230 235			840
cag tgc cca tat ttc ttc ttt gac gtt gtt tac atc cac aat ggc act Gln Cys Pro Tyr Phe Phe Phe Asp Val Val Tyr Ile His Asn Gly Thr 240 245 250			888
gag gag aag gag gag gag atg agt tgg aag gat atg ttt gcc ttc atg Glu Glu Lys Glu Glu Glu Met Ser Trp Lys Asp Met Phe Ala Phe Met 255 260 265			936

ggc agc ctg gat acc aag ggt acc agc tac aag tat gag gtg gca ctg 984
 Gly Ser Leu Asp Thr Lys Gly Thr Ser Tyr Lys Tyr Glu Val Ala Leu
 270 275 280

gct ggg cca gcc ctg gag ttg cac aac tgc atg gcg aaa ctg ttg gcc 1032
 Ala Gly Pro Ala Leu Glu Leu His Asn Cys Met Ala Lys Leu Leu Ala
 285 290 295 300

cac ccc ctg cag cgg cct tgc cag agc cat gct tcc tac agc ctg ctg 1080
 His Pro Leu Gln Arg Pro Cys Gln Ser His Ala Ser Tyr Ser Leu Leu
 305 310 315

gag gag gag gat gaa gcc att gag gtt gag gcc act gtc tga accatcc 1129
 Glu Glu Glu Asp Glu Ala Ile Glu Val Glu Ala Thr Val *
 320 325 330

ctgtacatct gcaccttctt gtgcaaggaa gtccttggcc taaagccttg gttctcaaac 1189

tgggttcctt gggacctccg ggggtgggggg gttccaggag gcat 1233